

# LIMIT CYCLES FOR CUBIC SYSTEMS WITH A SYMMETRY OF ORDER 4 AND WITHOUT INFINITE CRITICAL POINTS

M.J. ÁLVAREZ, A. GASULL AND R. PROHENS

ABSTRACT. In this paper we study the cubic systems which are invariant under a rotation of  $2\pi/4$  radians. They write as  $\dot{z} = \varepsilon z + p z^2 \bar{z} - \bar{z}^3$ , where  $z$  is complex, the time is real and  $\varepsilon = \varepsilon_1 + i\varepsilon_2$ ,  $p = p_1 + ip_2$  are complex parameters. When they have some critical points at infinity, *i.e.*  $|p_2| \leq 1$ , it is well-known that they can have at most one (hyperbolic) limit cycle which surrounds the origin. On the other hand when they have no critical points at infinity, *i.e.*  $|p_2| > 1$ , there are examples exhibiting at least two limit cycles surrounding nine critical points. In this paper we give two criteria for proving in some cases uniqueness and hyperbolicity of the limit cycle that surrounds the origin. Our results apply to systems having a limit cycle which surrounds either 1, 5 or 9 critical points, being the origin one of these points. The key point of our approach is the use of Abel equations.

## 1. INTRODUCTION AND MAIN RESULTS

This paper deals with the equation

$$\dot{z} = \frac{dz}{dt} = \varepsilon z + p z^2 \bar{z} - \bar{z}^3, \quad (1.1)$$

where  $z$  is a point in the complex plane,  $t$  is real,  $\varepsilon = \varepsilon_1 + i\varepsilon_2$ ,  $p = p_1 + ip_2$  are complex parameters and  $p_2 \geq 0$ . In Section 2 we will see that this last inequality can be assumed without loss of generality. We are interested in the number and location of the limit cycles of this equation that surround the origin and, eventually, other critical points.

The previous equation (1.1) is the particular case for  $q = 4$  of a general family, the one with a rotational invariance of  $2\pi/q$  radians ( $q \in \mathbb{N}$ ,  $q > 2$ ). In [1] it is proved that the differential equations which are invariant for a rotation of an angle of  $2\pi/q$  radians have the normal form

$$\dot{z} = z\mathcal{A}(|z|^2) + \mathcal{B}\bar{z}^{q-1} + O(|z|^{q+1}), \quad q > 2, \quad (1.2)$$

where  $\mathcal{A}$  is a complex polynomial of degree  $[(q-1)/2]$ . The above equation without the remainder term is called the *principal equation*. For instance,

the principal equations for the particular cases  $q = 3, 4$  are, respectively,

$$\dot{z} = a_0 z + a_1 z |z|^2 + \mathcal{B} \bar{z}^2, \quad \dot{z} = a_0 z + a_1 z |z|^2 + \mathcal{B} \bar{z}^3.$$

The cases  $q \leq 4$  are called *strong resonances* while the cases  $q \geq 5$  are called *weak resonances*. The flow for all the cases of the family except  $q = 4$  is studied in several places (see [8] for  $q < 4$  and [1] for  $q > 4$ ). The study of the periodic orbits of case  $q = 4$  turns out to be especially difficult. In the following we summarize several well-known partial results given in [1], [4], [10], [11] (here it can be found a complete study in the whole family (1.2)) and [13]. The study of equation (1.1) can be split in the following three cases:

- **Case 1:** Equation (1.1) has a unique critical point, the origin.
- **Case 2:** Equation (1.1) has five critical points, the origin and four saddle-nodes.
- **Case 3:** Equation (1.1) has nine critical points, the origin, four saddle points and four critical points of index +1.

Case 1 is totally understood. In [4] it is proved that in this case, there are only two possibilities for the global phase portrait of the equation: either there exists a unique periodic orbit, and when it exists it is hyperbolic, or the origin is a global attractor. In that article the authors also completely solve the case where the Poincaré compactification of equation (1.1) has critical points at infinity. As we will see in Lemma 2.2, it corresponds to the condition  $p_2 \leq 1$ . In this case, there exists at most one limit cycle encircling the origin and, when it exists, it is hyperbolic. Moreover, if there exist nonzero singular points they can not be surrounded by a periodic orbit.

The problem of the number of limit cycles not surrounding the origin corresponds to Case 3 and it is also totally solved. In [13] it is proved that either there are no limit cycles or that there are exactly four hyperbolic ones, each one of them surrounding exactly one of the critical points of index +1.

Thus, the remaining problem is to determine, in case that the infinity has no critical points, the number of limit cycles that can exist surrounding the origin, and eventually surrounding also the other 4 or 8 critical points, and whether they coexist or not with the 4 limit cycles that do not surround the origin. In [10] and [11] it is proved that, in the remaining case  $p_2 > 1$ , at least two limit cycles can exist surrounding the 9 critical points. Both works consider the Hamiltonian system,

$$\dot{z} = i\varepsilon_2 z + ip_2 z^2 \bar{z} - \bar{z}^3,$$

which phase portrait has 9 critical points (5 centers and 4 saddles) and a ring full of periodic orbits surrounding the 9 critical points. When this

equation is perturbed to get a complete equation (1.1) it is proved that, at least, two of the periodic orbits of that ring give rise to two hyperbolic limit cycles for the perturbed equation. As far as we know, the last results appeared for this case are mainly numeric or have been proved assisted by the computer, see [12] and [7] respectively.

Our main result is given in the following theorem.

**Theorem A.** (a) Consider equation (1.1) with  $\varepsilon_2 \neq 0$ ,  $p_2 > 1$  and define the following four numbers:

$$\begin{aligned}\Sigma_A^- &= \frac{\varepsilon_2 p_1 p_2 - \sqrt{\varepsilon_2^2 (p_1^2 + p_2^2 - 1)}}{p_2^2 - 1}, & \Sigma_A^+ &= \frac{\varepsilon_2 p_1 p_2 + \sqrt{\varepsilon_2^2 (p_1^2 + p_2^2 - 1)}}{p_2^2 - 1}, \\ \Sigma_B^- &= \frac{\varepsilon_2 p_1 p_2 - \sqrt{\varepsilon_2^2 (p_1^2 + 9p_2^2 - 9)}}{2(p_2^2 - 1)}, & \Sigma_B^+ &= \frac{\varepsilon_2 p_1 p_2 + \sqrt{\varepsilon_2^2 (p_1^2 + 9p_2^2 - 9)}}{2(p_2^2 - 1)}.\end{aligned}$$

If one of the conditions

$$(i) \quad \varepsilon_1 \notin (\Sigma_A^-, \Sigma_A^+), \quad (ii) \quad \varepsilon_1 \notin (\Sigma_B^-, \Sigma_B^+),$$

is satisfied then it has at most one limit cycle surrounding the origin. Furthermore, when it exists it is hyperbolic.

(b) There are equations (1.1) under condition (i) having exactly one hyperbolic limit cycle surrounding either 1 or 5 critical points and equations under condition (ii) having exactly one limit cycle surrounding either 1, 5 or 9 critical points.

*Remark 1.1.* In the case that  $\varepsilon_2 = 0$  equation (1.1) can be integrated and it is not difficult to see that it has no periodic orbits.

A key point for our study is that equation (1.1) can be transformed into an Abel equation. Some of the known results about the number of limit cycles surrounding the origin of equation (1.1) can also be reproduced by using our approach, that as far as we know has not been used before to study equation (1.1).

We will see in Proposition 2.5 that the results of Theorem A when condition (i) is satisfied are only useful for bounding the number of limit cycles surrounding the origin of equation (1.1) in the Cases 1 and 2 defined above. Condition (ii) also applies for some cases of the more interesting and difficult situation, namely Case 3, where the limit cycles that surround the origin can also surround the 9 critical points.

**Acknowledgement.** We are grateful to A. Zegeling for suggesting the study of this problem to us.

## 2. PRELIMINARY RESULTS

A reduction in the principal equation associated to equation (1.2) when  $q = 4$  is that if  $\mathcal{B} \neq 0$  it can be chosen to be real; in the literature it is found  $\mathcal{B} = 1$  or  $\mathcal{B} = -1$ . We have set this last possibility getting equation (1.1). Furthermore, as we have said before, we can always assume  $p_2 \geq 0$ , by changing  $t$  by  $-t$  and  $\theta$  by  $\theta + \pi$  if necessary. Our first result reduces the study of the periodic orbits of equation (1.1) that surround the origin to the study of the number of solutions that satisfy  $x(0) = x(2\pi)$  of an Abel equation

$$\frac{dx}{d\theta} = A(\theta)x^3 + B(\theta)x^2 + C(\theta)x, \quad (2.1)$$

with  $A, B$  and  $C$  smooth,  $2\pi$ -periodic functions. For these solutions their multiplicity can be defined as the usual multiplicity as zeroes of the displacement map defined by the flow of equation (2.1) between  $\theta = 0$  and  $\theta = 2\pi$ . We prove next result, which essentially follows from the results of [5].

**Lemma 2.1.** *Each periodic orbit  $\gamma$  of equation (1.1), with  $\varepsilon_2 \neq 0$ , that surrounds the origin corresponds to a non-zero solution  $\Gamma$  that satisfies  $x(0) = x(2\pi)$  of the Abel equation*

$$\frac{dx}{d\theta} = A(\theta)x^3 + B(\theta)x^2 + C(\theta)x, \quad (2.2)$$

where  $C(\theta) = 2\varepsilon_1/\varepsilon_2$ ,

$$\begin{aligned} A(\theta) &= \frac{4}{\varepsilon_2}(p_2 + \sin(4\theta))\left((\cos(4\theta) - p_1)\varepsilon_2 + (p_2 + \sin(4\theta))\varepsilon_1\right) \quad \text{and} \\ B(\theta) &= -\frac{2}{\varepsilon_2}\left(2\varepsilon_1p_2 - \varepsilon_2p_1 + 3\varepsilon_2\cos(4\theta) + 2\varepsilon_1\sin(4\theta)\right). \end{aligned}$$

Furthermore the multiplicity of the periodic orbit  $\gamma$  coincides with the multiplicity of  $\Gamma$ .

*Proof.* We can transform equation (1.1) in the cylinder  $\mathbb{R} \times [0, 2\pi]$  through the change of variables  $z = \sqrt{r}(\cos(\theta) + i\sin(\theta))$ , that is always well-defined because we are interested in  $r \geq 0$ . We obtain

$$\begin{cases} \dot{r} &= 2\varepsilon_1r + 2(p_1 - \cos(4\theta))r^2, \\ \dot{\theta} &= \varepsilon_2 + (p_2 + \sin(4\theta))r. \end{cases} \quad (2.3)$$

Note that the limit cycles surrounding the origin of (1.1) become non-contractible periodic orbits of system (2.3) in the cylinder with  $r > 0$  and, as it is proved in [3], they do not intersect the set  $\{(r, \theta) : \dot{\theta} =$

$\varepsilon_2 + (p_2 + \sin(4\theta))r = 0$ . So they are solutions of

$$\frac{dr}{d\theta} = \frac{2\varepsilon_1 r + 2(p_1 - \cos(4\theta))r^2}{\varepsilon_2 + (p_2 + \sin(4\theta))r}, \quad (2.4)$$

that satisfy  $r(0) = r(2\pi)$ . As  $\varepsilon_2 \neq 0$ , by using Cherkas' transformation,  $x = r/(\varepsilon_2 + (p_2 + \sin(4\theta))r)$ , see [5], the previous equation is transformed into the Abel equation (2.2). Moreover, all the non-contractible periodic orbits of system (2.3) are transformed into solutions of the Abel equation that satisfy  $x(0) = x(2\pi)$ . It is also clear that Cherkas' transformation keeps the multiplicity of the solutions. Thus the result follows.  $\square$

As we are mainly interested in the case that the infinity in the Poincaré compactification has no critical points (the other cases are already solved), we prove a lemma that gives which is this condition in terms of the parameters of equation (1.1). Moreover, we get the stability of the infinity and of the origin.

**Lemma 2.2.** *Consider equation (1.1).*

- (i) *The infinity in the Poincaré compactification has no critical points and only if  $p_2 > 1$ .*
- (ii) *When  $p_2 > 1$ , the infinity is an attractor (resp. a repellor) when  $p_1 > 0$  (resp.  $p_1 < 0$ ).*
- (iii) *The origin is an attractor (resp. a repellor) when  $\varepsilon_1 < 0$  (resp.  $\varepsilon_1 > 0$ ).*

*Proof.* If we make the change of variable  $R = 1/r$  and the reparametrization  $\frac{dt}{ds} = R$  we get a polar version of the Poincaré compactification.

$$\begin{cases} R' &= -2(p_1 - \cos(4\theta))R - 2\varepsilon_1 R^2, \\ \theta' &= p_2 + \sin(4\theta) + \varepsilon_2 R. \end{cases} \quad (2.5)$$

Then, the previous system has no critical points on the infinity, which is given by the invariant set  $\{R = 0\}$ , if and only if  $p_2 > 1$ .

If we want to compute the stability of infinity in the case that  $p_2 > 1$ , applying the results in [9] we have that the stability of  $\{R = 0\}$  in system (2.5) is given by the sign of

$$\int_0^{2\pi} \frac{-2(p_1 - \cos(4\theta))}{p_2 + \sin(4\theta)} d\theta = \frac{-4\pi p_1}{\sqrt{p_2^2 - 1}},$$

and then the result follows. The study of the stability of the origin is trivial because it is a nondegenerate critical point.  $\square$

We will need the following general result about Abel equations proved in [6], that we state for the sake of completeness.

**Theorem 2.3.** *Consider the Abel equation (2.1) and assume that either  $A(\theta) \not\equiv 0$  or  $B(\theta) \not\equiv 0$  does not change sign. Then it has at most three solutions satisfying  $x(0) = x(2\pi)$ , taking into account their multiplicities.*

Next lemma gives explicit conditions on  $p$  and  $\epsilon$  to ensure that either the function  $A$  or the function  $B$  associated to equation (1.1) does not change sign.

**Lemma 2.4.** *When  $p_2 > 1$ , consider the values  $\Sigma_A^+, \Sigma_A^-, \Sigma_B^+$  and  $\Sigma_B^-$  defined in Theorem A. Let  $A(\theta)$  and  $B(\theta)$  be the functions given in Lemma 2.1. Then*

- (i)  $\epsilon_1 \notin (\Sigma_A^-, \Sigma_A^+)$  if and only if  $A(\theta)$  does not change sign,
- (ii)  $\epsilon_1 \notin (\Sigma_B^-, \Sigma_B^+)$  if and only if  $B(\theta)$  does not change sign.

*Proof.* (i) It is not difficult to check that  $A(\theta)$  does not change sign in the interval for  $\epsilon_1$  stated above. A sketch of the verification is as follows. We compute the maxima and the minima of the function  $A(\theta)$ . As  $p_2 > 1$  it is equivalent to compute the extrema of  $(\cos(4\theta) - p_1)\epsilon_2 + (p_2 + \sin(4\theta))\epsilon_1$ . We apply the change of variables  $\cos(4\theta) = x$  and  $\sin(4\theta) = y$ , and we optimize the function subject to the restriction  $x^2 + y^2 = 1$ . We get that the value of the function at both extrema is  $A_{\pm} = \epsilon_1 p_2 - \epsilon_2 p_1 \pm \sqrt{\epsilon_1^2 + \epsilon_2^2}$ . They are both positive or negative if and only if  $\epsilon_1 \in (-\infty, \Sigma_A^-] \cup [\Sigma_A^+, +\infty)$ .

(ii) In the expression of the function  $B(\theta)$  we substitute  $\cos(4\theta)$  by  $x$  and  $\sin(4\theta)$  by  $y$  subject to the restriction  $x^2 + y^2 = 1$ . We get that the only two extrema are on  $(x_+, y_+)$  and  $(x_-, y_-)$  where  $x_{\pm} = \frac{\pm 3\epsilon_2}{\sqrt{4\epsilon_1^2 + 9\epsilon_2^2}}$  and  $y_{\pm} = \frac{\pm 2\epsilon_1}{\sqrt{4\epsilon_1^2 + 9\epsilon_2^2}}$ . The values of the function  $B(\theta)$  on them are  $B_{\pm} = 2 \left( \epsilon_2 p_1 - 2\epsilon_1 p_2 \mp \sqrt{4\epsilon_1^2 + 9\epsilon_2^2} \right)$ . Hence, if they are both positive or negative then  $B(\theta)$  will not change sign. These are precisely the conditions of the statement of the lemma.  $\square$

Next result gives a geometric interpretation of the behaviour of the function  $A(\theta)$  appearing in the Abel equation (2.2) in terms of the number of critical points of equation (1.1).

**Proposition 2.5.** *Consider equation (1.1) and assume that  $\epsilon_2 \neq 0$  and  $p_2 > 1$ . Set  $\Theta := \{(r, \theta) : \dot{\theta} = \epsilon_2 + (p_2 + \sin(4\theta))r = 0\}$ . Then*

- (i) *If  $\Theta = \emptyset$  then equation (1.1) has only one critical point, the origin.*
- (ii) *When  $\Theta \neq \emptyset$ , the set  $\Theta$  is a closed curve that surrounds the origin.*

*Furthermore:*

- (a) *If  $\epsilon_2 A(\theta) > 0$  (resp.  $< 0$ ) then equation (1.1) has only one critical point, the origin, and the interior of  $\Theta$  is negatively (resp. positively) invariant.*

(b) If  $\varepsilon_2 A(\theta) \geq 0$  (resp.  $\leq 0$ ) then equation (1.1) has five critical points and the vector field associated to the differential equation (1.1) points towards the exterior (resp. interior) of the region delimited by  $\Theta$  at all points of this curve except at the four critical points of the vector field over it.

(c) If  $A(\theta)$  changes sign then equation (1.1) has nine critical points.

*Proof.* Following the proof of Lemma 2.1 we write equation (1.1) as (2.3). The proof of (i) is trivial.

(ii) The non-zero critical points are the points  $(r, \theta)$ , with  $r > 0$ , that belong to  $\{\dot{\theta} = 0\} \cap \{\dot{r} = 0\}$ . Since  $p_2 + \sin(4\theta) \neq 0$ , the first set,  $\Theta$ , is given by the closed curve  $r = -\varepsilon_2 / (p_2 + \sin(4\theta))$ . If we compute  $\dot{r}$  on  $\Theta$  we obtain

$$\dot{r}|_{\dot{\theta}=0} = r \frac{\varepsilon_2 A(\theta)}{(p_2 + \sin(4\theta))^2}.$$

From the above equality the lemma follows.  $\square$

### 3. PROOF OF THEOREM A

Let us define the function  $c(\theta) = p_2 + \sin(4\theta)$ . Observe that, as  $p_2 > 1$ ,  $c(\theta) > 0$  for all  $\theta \in [0, 2\pi]$ .

(a) We start assuming that condition (i) is satisfied. By using Lemma 2.1 we know that the study of the periodic orbits of equation (1.1) can be reduced to the study of the Abel equation (2.2). Furthermore, since  $\varepsilon_1 \notin (\Sigma_A^-, \Sigma_A^+)$ , by condition (i) of Lemma 2.4 we know that the function  $A(\theta)$  appearing in the Abel equation does not change sign. Hence, from Theorem 2.3, the maximum number of solutions satisfying  $x(0) = x(2\pi)$  in system (2.2), taking into account their multiplicity, is three. One of them is  $x = 0$ . Since  $c(\theta) \neq 0$ , by some simple calculations it is easy to prove that the curve  $x = 1/c(\theta)$  is a second solution satisfying this condition. If we undo the Cherkas' transformation used to convert equation (2.4) into the Abel equation it turns out that  $x = 1/c(\theta)$  is mapped into the infinity of the differential equation. Then, by Lemma 2.1, the maximum number of limit cycles of equation (1.1) is one. The fact that this limit cycle, when exists, is hyperbolic, also follows from the same lemma.

The proof of (a) when condition (ii) is satisfied follows the same steps, but changing the role of  $A(\theta)$  by the role of  $B(\theta)$ .

(b) We fix the parameters of equation (1.1) as:  $\varepsilon_2 = -1$ ,  $p_1 = \frac{-1}{2}$  and  $p_2 = \frac{5}{4}$ . In this case,  $\Sigma_A^+ = \frac{2}{9}(5 + 2\sqrt{13})$ , and  $\Sigma_B^+ = \frac{1}{9}(5 + 2\sqrt{85}) < \Sigma_A^+$ .

We study the family for  $\varepsilon_1 > \Sigma_A^+$ . Applying Lemma 2.2, the stability of the origin is given by the sign of  $\varepsilon_1 > 0$ , hence it is a strong unstable focus. By the same lemma, we get that the infinity in the Poincaré compactification is also unstable.

As  $\varepsilon_1 > \Sigma_A^+$  then  $A(\theta)$  does not vanish and there are no other critical points but the origin. Furthermore since  $\varepsilon_2 A(\theta) > 0$ , by using Lemma 2.5.(b) we know that the exterior of the closed curve  $\Theta$  is positively invariant. Hence by using the Poincaré-Bendixson Theorem, the behaviour at infinity and part (a) of the theorem, we know that the differential equation has exactly one limit cycle which surrounds the curve  $\Theta$  and that it is hyperbolic and stable.

When  $\varepsilon_1 = \Sigma_A^+$  there appear four semi-elementary critical points on  $\Theta$ , that are saddle-nodes. We are going to prove that, at this value of the parameter  $\varepsilon_1$ , the periodic orbit still exists and surrounds the five critical points. Since the infinity is always a repellor, the only way for this limit cycle to disappear is going, when  $\varepsilon_1$  goes to  $\Sigma_A^+$ , to a limit periodic set formed by the four saddle-nodes and their separatrices. To prevent this situation we will construct a polygonal line without any contact point with the flow of the differential equation, which will help us to control the behaviour of the separatrices of the four saddle-nodes. More concretely, this polygonal line forces the unstable separatrices of the saddle-nodes to have as  $\omega$ -limit a periodic orbit surrounding  $\Theta$ , see Figure 1.

Because of the symmetry of the system it is enough to study one of the saddle-nodes, for instance the one in the first quadrant, namely  $(x_0, y_0)$ , in cartesian coordinates.

Let  $x_1$  be the unique real zero of the equation

$$\begin{aligned} & \frac{5}{4}x^3 + \frac{3}{2}(-2 + \sqrt{10})x^2 + \left(\frac{53}{10} - 9\sqrt{\frac{2}{5}}\right)x + \\ & + \frac{1}{375}(7762 - 2423\sqrt{10} + 100\sqrt{13}(-2 + \sqrt{10})) = 0, \end{aligned}$$

that is, approximately,  $x_1 \simeq 0.56$ , and consider the polygonal line defined as

$$(x(t), y(t)) = \begin{cases} (t, t) & \text{if } 0 \leq t < \sqrt{\frac{2}{5}} \\ (t, 3(t - \sqrt{\frac{2}{5}}) + \sqrt{\frac{2}{5}}) & \text{if } \sqrt{\frac{2}{5}} \leq t < 1 \\ ((t-1)(x_1-1) + 1, 3 - 2\sqrt{\frac{2}{5}}) & \text{if } 1 \leq t < 2 \\ ((t-3)(x_0-x_1) + x_0, \\ (t-3)(y_0 - 3 + 2\sqrt{\frac{2}{5}}) + y_0) & \text{if } 2 \leq t < 3. \end{cases}$$

The flow associated to equation (2.3) is transversal to the previous polygonal line, that is, the polygonal has not any contact point. We give the details for  $\sqrt{\frac{2}{5}} \leq t < 1$ . The other cases follow by using analogous arguments.



Let us consider the explicit expression of the second segment of the polygonal line as  $y = 3(x - \sqrt{\frac{2}{5}}) + \sqrt{\frac{2}{5}}$ . The scalar product of the vector field associated to equation (2.3) with the normal vector to former segment,  $(-3, 1)$ , is given by

$$-2\sqrt{\frac{2}{5}} \frac{(77 + 20\sqrt{13})}{45} + \frac{62x}{5} - 59\sqrt{\frac{2}{5}}x^2 + 29x^3,$$

when it is evaluated on the segment. This cubic equation has only one real root  $x_2 > 1$ , which is simple. Hence, for  $x \in [\sqrt{\frac{2}{5}}, 1]$  the scalar product is negative and, consequently, the flow crosses the considered segment transversally.

By using the previous arguments and the fact that the infinity is a repeller, by the Poincaré-Bendixson Theorem the only possible  $\omega$ -limit for the unstable separatrix of the saddle-node is a periodic orbit which has to surround the four saddle-nodes, see again Figure 1.




FIGURE 1. The polygonal curve without any contact and the separatrices of the saddle-node of system (2.3).

Since for the value  $\varepsilon_1 = \Sigma_A^+$  the function  $A(\theta)$  does not change its sign, from part (a) of this theorem, we get the hyperbolicity of the limit cycle.

Now we move  $\varepsilon_1$  a little more towards zero but very close to  $\Sigma_A^+$ . In this case,  $B(\theta)$  is strictly positive because  $\Sigma_A^+ > \Sigma_B^+$  and we have 9 critical points: four saddles and four critical points of index +1 on  $\Theta$  and a focus at the origin. Applying again part (a), but now condition (ii), we know that the maximum number of limit cycles surrounding the origin is one. As for  $\varepsilon_1 = \Sigma_A^+$  the limit cycle was hyperbolic, it still exists. Then we have a vector field with  $B(\theta)$  not changing sign, 8 non-zero critical points and a limit cycle surrounding them and the origin.

□

**3.1. Some Remarks on the proof of Theorem A.** (i) We note that in the proof of the above theorem, when  $\varepsilon_1 = \Sigma_A^+$ , the vector field associated to equation (1.1) points towards the exterior of the region delimited by  $\Theta$  at all points of this curve except at the four saddle-nodes. Nevertheless one of the stable separatrices of each one of the saddle-nodes lies outside of the region bounded by  $\Theta$ . Hence, the behaviour of the vector field on this curve is not enough to conclude that the limit cycle still exists for this value of  $\varepsilon_1$  and we have needed to introduce the polygonal line used in the proof.

(ii) The fact that when  $c(\theta)$  does not vanish the curve  $x = 1/c(\theta)$  is a periodic orbit of the Abel equation (2.2) is not exclusive for this particular equation. In fact, it is valid for all the Abel equations obtained from the Cherkas' transformation applied to planar polynomial vector fields with homogeneous non-linearities, see for instance [2].

(iii) The same result stated as condition (i) of our main theorem but only when the function  $A(\theta)$  does not vanish has also been proved in [4] by using different methods. Dynamically there is a big difference, because in our result allowing  $A(\theta)$  to vanish but not changing sign means that equation (2.2) has exactly 5 critical points. On the other hand the results of [4] apply only when the critical point is unique.

(iv) Observe that the values of the parameters considered in the example given in (b) of the previous result are not close to any Hamiltonian system. Moreover, the concrete set of values of our example are in the region C of [11] and in the region iii) of [4], the most difficult region to study, as said by Arnold in [1].

**Acknowledgements.** The first two authors are partially supported by grants MTM2005-06098-C02-1 and 2005SGR-00550. This paper is also supported by the CRM Research Program: On Hilbert's 16th Problem.

#### REFERENCES

1. ARNOLD, V. *Chapitres supplémentaires de la théorie des équations différentielles ordinaires*. Éditions Mir-Moscou, 1980.

2. CARBONELL, M.; LLIBRE, J. *Limit cycles of polynomial systems with homogeneous nonlinearities*. J. Math. Anal. Appl., **142** (1989), pp. 573–590.
3. COLL, B.; GASULL, A.; PROHENS, R. *Differential equations defined by the sum of two quasi-homogeneous vector fields*. Can. J. Math., **49** (1997), pp. 212–231.
4. CHENG, C.Q.; SUN, Y.S. *Metamorphoses of phase portraits of vector field in the case of symmetry of order 4*. J. of Diff. Equations, **95** (1992), pp. 130–139.
5. CHERKAS, L.A. *Number of limit cycles of an autonomous second-order system*. Diff. Equations, **5** (1976), pp. 666–668.
6. GASULL, A.; LLIBRE, J. *Limit cycles for a class of Abel Equation*. SIAM J. Math. Anal., **21** (1990), 1235–1244.
7. GUCKENHEIMER, J. *Phase portraits of planar vector fields: computer proofs*. Exp. Mathematics, **4** (1995), pp. 153–165.
8. HOROZOV, E. *Versal deformations of equivariant vector field for the case of symmetries of order 2 and 3*. Trudy Sem. Pet., **5** (1979), pp. 163–192 (in russian).
9. LLOYD, N.G. *A note on the number of limit cycles in certain two-dimensional systems*. J. London Math. Soc., **20** (1979), 277–286.
10. NEISHTADT, A.I. *Bifurcation of the phase pattern of an equation system arising in the problem of stability loss of self-oscillations close to 1:4 resonance* (in russian). J. Appl. Math. Mech., **42** (1978), n. 5, pp. 830–840.
11. CHOW, S.N.; LI, C.; WANG, D. *Normal Forms and Bifurcation of Planar Vector Fields*. Cambridge University Press (1994).
12. YU, P.; HAN, M.; YUAN, Y. *Analysis on limit cycles of  $Z_q$ -equivariant polynomial vector fields with degree 3 or 4*. To appear in J. Math. Anal. Appl. (2006).
13. ZEGELING, A. *Equivariant unfoldings in the case of symmetry of order 4*. Bulgaricae Mathematicae Pub., **19** (1993), pp. 71–79.

M.J. Álvarez, R. Prohens  
Departament de Matemàtiques i Informàtica,  
Universitat de les Illes Balears, 07122, Palma de Mallorca, Spain.  
*E-mail*: chus.alvarez@uib.es; rafel.prohens@uib.es

A. Gasull  
Departament de Matemàtiques, Edifici Cc  
Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain.  
*E-mail*: gasull@mat.uab.es