

# PG SNPs Template (2 pages maximum)

*Required subheadings for the report are indicated below in bold and italics*

## ***TRIM63 (MuRF-1) Gene Polymorphism is Associated with Biomarkers of Exercise-Induced Muscle Damage***

### **Authors and Institutions:**

Baumert P<sup>1</sup>, Lake M<sup>1</sup>, Drust B<sup>1</sup>, Stewart CE<sup>1</sup>, Erskine RM<sup>1,2</sup>

<sup>1</sup>*Research Institute for Sport & Exercise Sciences, Liverpool John Moores University, Liverpool, UK;*

<sup>2</sup>*Institute of Sport, Exercise & Health, University College London, London, UK,*

### **ABSTRACT**

Unaccustomed strenuous exercise can lead to muscle strength loss, inflammation and delayed onset muscle soreness, which may be influenced by genetic variation. We investigated if a missense single nucleotide polymorphism (A>G, rs2275950) within the *TRIM63* gene (encoding MuRF-1 and potentially affecting titin mechanical properties) was associated with the variable response to unaccustomed eccentric exercise. Sixty-five untrained, healthy participants (genotyped for rs2275950: AA, AG and GG) performed 120 maximal eccentric knee extensions (ECC) to induce muscle damage. Isometric and isokinetic maximal voluntary knee extension contractions (MVCs) and muscle soreness were assessed before, immediately after, and 48h after ECC. AA homozygotes were consistently stronger [baseline isometric MVC: 3.23±0.92 Nm/kg (AA) vs. 2.09±0.67 Nm/kg (GG); p=0.006] and demonstrated less muscle soreness over time (p=0.022) compared to GG homozygotes. This may be explained by greater titin stiffness in AA homozygotes, leading to intrinsically stronger muscle fibers that are more resistant to eccentric damaging contractions.

### **Keywords**

Tripartite Motif Containing 63 (TRIM63) Single Nucleotide Polymorphism (SNP), E3 ubiquitin-protein ligase, Exercise-Induced Muscle Damage (EIMD), Delayed onset muscle soreness (DOMS) phenotype, MURF1 Gene

### **BACKGROUND/MOTIVATION FOR THE STUDY**

Unaccustomed strenuous exercise can lead to muscle strength loss, inflammation and delayed onset muscle soreness, which are biomarkers of exercise-induced muscle damage (EIMD) (3).

Insufficient recovery from EIMD may increase injury risk, and there is evidence that people recover at different rates (3), which may be influenced by genetic variation (1).

Muscle RING finger-1 (MuRF-1) is known to be associated with cardiomyopathy and muscle atrophy (6) and it plays a crucial role in the ubiquitin proteasome system by attaching ubiquitin polymers to damaged myofibrillar proteins following EIMD, thus enabling the 26S-proteasome complex to degrade these ubiquitin-marked proteins (4). MuRF-1 is localized at the Z- and also the M-line of the sarcomere, where it also interacts with the strain sensing kinase domain of the sarcomeric protein, titin (2). This kinase domain is thought to modulate the mechanical properties of titin and the gene expression of other muscle-encoding proteins (5).

MuRF-1 is encoded by the human tripartite motif containing 63 (*TRIM63*) gene, in which a missense single nucleotide polymorphism (SNP, A>G, rs2275950) at amino acid 237 causes a change from lysine to glutamate. This may decrease MuRF-1's affinity for the titin strain sensing kinase domain, thus reducing muscle fiber stiffness and increasing its susceptibility to EIMD. To our knowledge, however, this SNP has not yet been linked to any phenotypic trait. We therefore aimed to investigate an association between this SNP and EIMD in untrained young men and women. We hypothesized that (i) the minor G-allele would be associated with lower muscle strength, and (ii) G-allele carriers would exhibit a greater damage response following maximal eccentric exercise compared to AA homozygotes.

### ***PHENOTYPE***

To induce quadriceps femoris muscle damage, familiarized participants performed 12 sets of 10 maximal eccentric unilateral contractions of the knee extensors. Maximum isometric (at 80° knee flexion) and isokinetic (60°/s) voluntary knee-extension contraction (MVC) torque (Humac Norm, CSI Inc, Stoughton, MA, and Biodex Multi-Joint System 3 Pro Inc, Shirley, NY) normalized to body mass, muscle soreness (measured by visual analogue scale in conjunction with a three repetition bodyweight squat) were assessed before, immediately after and 48 h after the EIMD intervention. Venous blood samples were collected from an antecubital vein for genotypic (1x10 ml; BD EDTA vacutainer) and serum analysis (3x10 ml; BD serum collection vacutainer). Serum tubes were centrifuged at 1,300 g for 15 min (4°C) and all samples were stored at -80°C. Serum creatine kinase (CK) activity (Catachem Inc., Connecticut, NE) was analyzed according to manufacturer protocols.

### **Cohort details**

Following ethical approval from Liverpool John Moores University Ethics Committee, and in accordance with the Declaration of Helsinki, written informed consent was obtained from our population cohort, which comprised young, untrained, healthy (identified by physical activity and health questionnaires) female (n=39) and male (n=26) Caucasians (mean  $\pm$  SD: age =  $22.5 \pm 4.0$  year; height =  $1.71 \pm 0.09$  m; body mass =  $70.9 \pm 14.4$  kg). Inclusion criteria were: aged between 18-35 years and no history of (i) leg strength training in the past 6 months, (ii) muscle-tendon injury in the last 12 months, and (iii) bone fracture in the lower limbs. Participants were requested to maintain their normal daily routine and to refrain from consuming alcohol or participating in any exercise for two days prior to the study and throughout the intervention.

### **Type of study**

For this candidate SNP study, extraction of the DNA was carried out using QIAamp DNA Blood Kit (Qiagen, Crawley, UK) and following the spin protocol for DNA purification from whole blood. Real-time polymerase chain reaction (PCR) was performed with Rotor-Gene Q (Qiagen) to determine *TRIM63* (A>G, rs2275950) genotype in each participant. Reactions were completed on a 72-well rotor-disc. Each 10  $\mu$ L reaction volume contained: 5  $\mu$ L Genotyping Master Mix, 0.5  $\mu$ L genotyping assay mix (Applied Biosystems, Foster City, CA), 3.5  $\mu$ L nuclease-free H<sub>2</sub>O (Qiagen) and 1  $\mu$ L DNA. For control wells, 1  $\mu$ L nuclease-free H<sub>2</sub>O replaced the DNA template. The following PCR protocol was used: 40 cycles of incubation at 92°C for 15s (denaturation), then annealing and extension at 60°C for 1min. Lastly, genotype was determined in all samples (analyzed in duplicate), using Rotor-Gene Q Software 2.3.1 (Qiagen).

### **Details of the SNP(s) studied**

The investigated SNP (rs2275950) is located in the Tripartite Motif Containing 63 gene (*TRIM63*), on chromosome 1 at position 26,058,512 (dbSNP Build 150). This missense A>G (adenine to guanine) SNP has not previously been associated with the response to exercise. The frequency of the effect (major) A allele is 0.77. Linkage disequilibrium (LD) calculations for the *TRIM63* SNP were performed using the LDlink suite and data from the 1000 Genomes Project European ancestry populations (7, 8).

### ***Analysis model***

Data for each parameter were assessed for normal distribution with the Shapiro-Wilk test and by inspection of the Q-Q plots. Hardy-Weinberg equilibrium was determined for the *TRIM63* SNP using a  $\chi^2$  test. Two-way mixed ANCOVAs [within-subjects factor: time (pre, post- and 48h post; between-subjects factor: genotype (AA, AG, GG); covariate: sex] with Tukey post-hoc tests were used to detect associations between the *TRIM63* SNP and isometric and isokinetic MVC knee-extension torque (normalized to body mass), muscle soreness, and serum CK activity over time. All data are presented as mean ( $\pm$  SD), and statistical significance was identified when  $P < 0.05$ . All MVC data were recorded in AcqKnowledge software (Version 4.2, Biopac-Systems Inc., Goleta, USA) and statistical analysis was performed with SPSS (v23, IBM Inc., Armonk, NY).

### ***RESULTS***

Genotype frequency distribution for the *TRIM63* (rs2275950;  $\chi^2=1.156$ ,  $P=0.282$ ) SNP was in Hardy-Weinberg equilibrium. Forty participants were AA, 20 AG, and five were GG genotype. Normalized isometric ( $p<0.001$ ) and isokinetic ( $p<0.001$ ) MVC, leg muscle soreness ( $p<0.001$ ) and CK activity ( $p=0.01$ ) all showed a main effect of time, indicating EIMD. There was a main effect for genotype regarding normalized isometric ( $p=0.006$ , Fig. 1) and isokinetic ( $p=0.031$ ) MVC, with AA stronger than GG in both cases [isometric MVC normalized to body mass at pre-EIMD:  $3.23\pm 0.92$  Nm/kg (AA) vs.  $2.67\pm 1.09$  Nm/kg (AG) vs.  $2.09\pm 0.67$  Nm/kg (GG)]. There was a genotype x time interaction regarding muscle soreness ( $p=0.022$ ), with AA homozygotes showing attenuated muscle soreness 48 h post-EIMD compared to AG and GG genotypes ( $2.9\pm 2.2$  cm vs.  $4.5\pm 2.3$  cm and  $5.4\pm 2.2$  cm, respectively; Fig. 2).

### ***INTERPRETATION***

Homozygotes of the major *TRIM63* A-allele were stronger (relative to body mass) and recovered quicker following strenuous exercise compared to G-allele carriers. It is possible that the A-allele is linked to a higher affinity for titin's strain sensing kinase domain, leading to greater kinase activation. This may improve the mechanical properties of titin, enabling a stiffer muscle fiber to transmit more force laterally and longitudinally, potentially making the fiber intrinsically stronger and more resistant to acute damaging eccentric contractions. However, rs2275950 is in LD with other *TRIM63* SNPs, and with SNPs within the exostosin-like 1 (*EXTL1*) and the solute carrier family 30 member 2 (*SLC30A2*) genes. Nevertheless, given MuRF-1's

role in muscle protein degradation and with titin's strain sensing kinase domain, rs2275950 is a likely genetic determinant of the inter-individual variability in the human response to EIMD. Further work is necessary to elucidate if (i) other *TRIM63*, *EXTL1* and/or *SLC30A2* SNPs (either in combination or isolation) are of functional significance regarding EIMD; (ii) this quicker recovery following an acute bout of intense exercise in AA homozygotes compared to G-allele carriers translates into greater muscle hypertrophy and/or strength gains following chronic resistance training.

## **ACKNOWLEDGMENTS**

## **GRANTS**

## **DISCLOSURES**

No conflicts of interest, financial or otherwise, are declared by the author(s).

## **AUTHOR CONTRIBUTIONS**

R.M.E conception and design of study; P.B., A.B, J.A.C.,V.E, and K.O.J. performed experiments; P.B, and R.M.E analyzed data; R.M.E., P.B, and C.E.S. interpreted results of experiments; P.B. prepared figures; P.B. drafted manuscript; R.M.E., P.B., M.L., B.D. and C.E.S edited and revised manuscript; R.M.E., P.B., M.L., B.D. and C.E.S. approved final version of manuscript.

## **REFERENCES**

1. **Baumert P, Lake MJ, Stewart CE, Drust B, and Erskine RM.** Genetic variation and exercise-induced muscle damage: implications for athletic performance, injury and ageing. *Eur J Appl Physiol* 1-31, 2016.
2. **Centner T, Yano J, Kimura E, McElhinny AS, Pelin K, Witt CC, Bang M-L, Trombitas K, Granzier H, and Gregorio CC.** Identification of muscle specific ring finger proteins as potential regulators of the titin kinase domain. *J Mol Biol* 306: 717-726, 2001.
3. **Clarkson PM, and Hubal MJ.** Exercise-induced muscle damage in humans. *American journal of physical medicine & rehabilitation* 81: S52-S69, 2002.
4. **Kerksick CM, Roberts MD, Dalbo VJ, Kreider RB, and Willoughby DS.** Changes in skeletal muscle proteolytic gene expression after prophylactic supplementation of EGCG and NAC and eccentric damage. *Food Chem Toxicol* 61: 47-52, 2013.
5. **Kötter S, Andresen C, and Krüger M.** Titin: central player of hypertrophic signaling and sarcomeric protein quality control. *Biol Chem* 395: 1341-1352, 2014.

Appendix 1 Figure 1




Figure 1. Isometric knee extension maximal voluntary contraction torque normalized to body mass for individuals of the three TRIM63 (A>G, rs2275950) genotypes before (Pre), directly after (Post) and 48h after (48 h Post) a strenuous exercise intervention. \* indicates significant differences ( $p < 0.05$ ) between the AA and GG genotypes. MVC = Maximal voluntary contraction. Values are mean  $\pm$  SD.

Appendix 1 Figure 2




Figure 2. Muscle Soreness measured via Visual Analogue Scale for individuals of the three TRIM63 (A>G, rs2275950) genotypes assessed before (Pre), directly after (Post) and 48h after (48 h Post) 120 a strenuous exercise intervention. \* indicates significant differences ( $p < 0.05$ ) between AA and GG genotypes. † indicates significant differences ( $p < 0.05$ ) between AA and AG genotypes. Values are mean  $\pm$  SD.

## Appendix 3

### List of the G-REX consortium members:

Philipp Baumert<sup>1</sup>, Bethany Adams<sup>1</sup>, Josephine A Cabot<sup>1</sup>, Victoria Edwards<sup>1</sup>, Kelsie O Johnson<sup>1</sup>, Mark Lake<sup>1</sup>, Barry Drust<sup>1</sup>, Claire E Stewart<sup>1</sup>, Robert M Erskine<sup>1,2</sup>

<sup>1</sup>*Research Institute for Sport & Exercise Sciences, Liverpool John Moores University, Liverpool, UK;*

<sup>2</sup>*Institute of Sport, Exercise & Health, University College London, London, UK,*

### Additional References

6. **Watanabe M, and Hatakeyama S.** TRIM proteins and diseases. *The Journal of Biochemistry* 161: 135-144, 2017.
7. **Consortium GP.** An integrated map of genetic variation from 1,092 human genomes. *Nature* 491: 56-65, 2012.
8. **Machiela MJ, and Chanock SJ.** LDlink: a web-based application for exploring population-specific haplotype structure and linking correlated alleles of possible functional variants. *Bioinformatics* 31: 3555-3557, 2015.