

Publ. Mat. **53** (2009), 231–244**OPERATOR VALUED *BMO* AND COMMUTATORS**

OSCAR BLASCO

*Abstract*


---

If  $E$  is a Banach space,  $b \in BMO(\mathbb{R}^n, \mathcal{L}(E))$  and  $T$  is a  $\mathcal{L}(E)$ -valued Calderón-Zygmund type operator with operator-valued kernel  $k$ , we show the boundedness of the commutator  $T_b(f) = bT(f) - T(bf)$  on  $L^p(\mathbb{R}^n, E)$  for  $1 < p < \infty$  whenever  $b$  and  $k$  verify some commuting properties. Some endpoint estimates are also provided.

---

**1. Introduction and notation**

We shall work on  $\mathbb{R}^n$  endowed with the Lebesgue measure  $dx$  and use the notation  $|A| = \int_A dx$ . Given a Banach space  $(X, \|\cdot\|)$  and  $1 \leq p < \infty$  we shall denote by  $L^p(\mathbb{R}^n, X)$  the space of Bochner  $p$ -integrable functions endowed with the norm  $\|f\|_{L^p(\mathbb{R}^n, X)} = (\int_{\mathbb{R}^n} \|f(x)\|^p dx)^{1/p}$ , by  $L_c^\infty(\mathbb{R}^n, X)$  the closure of the compactly supported functions in  $L^\infty(\mathbb{R}^n, X)$  and by  $L_{\text{weak}, \alpha}(\mathbb{R}^n, X)$  the space of measurable functions such that  $|\{x \in \mathbb{R}^n : \|f(x)\| > \lambda\}| \leq \alpha(\lambda)$  where  $\alpha: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a non increasing function. We write  $H^1(\mathbb{R}^n, X)$  for the Hardy space defined by  $X$ -valued atoms, that is the space of integrable functions  $f = \sum_k \lambda_k a_k$  where  $\lambda_k \in \mathbb{R}$ ,  $\sum_k |\lambda_k| < \infty$  and  $a_k$  belong to  $L_c^\infty(\mathbb{R}^n, X)$ ,  $\text{supp}(a_k) \subset Q_k$  for some cube  $Q_k$ ,  $\int_{Q_k} a(x) dx = 0$  and  $\|a(x)\| \leq \frac{1}{|Q_k|}$ . We also write, for a positive function  $\phi$  defined on  $\mathbb{R}^+$ ,  $BMO_\phi(\mathbb{R}^n, X)$  for the space of locally integrable functions such that there exists  $C > 0$  such that for all cube  $Q$

$$\frac{1}{|Q|} \int_Q \|f(x) - f_Q\| dx \leq C\phi(|Q|)$$

---

2000 *Mathematics Subject Classification*. Primary: 42B20; Secondary: 42B25.

*Key words*. Bounded mean oscillation, commutator, operator-valued singular integrals.

Partially supported by Proyecto MTM 2005-08350.

where  $f_Q = \frac{1}{|Q|} \int_Q f(x) dx$ . For  $\phi(t) = 1$  we denote the space  $BMO(\mathbb{R}^n, X)$  and the above condition is equivalent to

$$\text{osc}_p(f, Q) = \left( \frac{1}{|Q|} \int_Q \|f(x) - f_Q\|^p dx \right)^{1/p} < \infty$$

for each (equivalently for all)  $1 \leq p < \infty$ . The infimum of the constants satisfying the above inequalities define the “norm” in the space.

Let us denote by  $f^\#$  and  $M(f)$  the sharp and the Hardy-Littlewood maximal functions of  $f$  defined by

$$f^\#(x) = \sup_{x \in Q} \text{osc}_1(f, Q) \quad \text{and} \quad M(f)(x) = \sup_{x \in Q} \frac{1}{|Q|} \int_Q \|f(x)\| dx.$$

We write  $M_q(f) = M(\|f\|^q)^{1/q}$  for  $1 \leq q < \infty$ .

It is well known that

$$(1) \quad f^\#(x) \approx \sup_{x \in Q} \inf_{c_Q \in X} \frac{1}{|Q|} \int_Q \|f(x) - c_Q\| dx$$

and that  $f^\# \in L^p(\mathbb{R}^n)$  implies that  $f \in L^p(\mathbb{R}^n, X)$  for  $1 < p < \infty$ .

Recall also that  $M_q$  maps  $L^q(\mathbb{R}^n, X)$  into  $L_{\text{weak}, 1/t^q}$  and

$$(2) \quad M_q: L^p(\mathbb{R}^n, X) \rightarrow L^p(\mathbb{R}^n)$$
 is bounded for  $q < p \leq \infty$ .

Throughout the paper  $E$  denotes a Banach space and  $\mathcal{L}(E)$  denotes the space of bounded linear operators on  $E$ .

**Definition 1.1.** We shall say that  $T$  is a  $\mathcal{L}(E)$ -Calderón-Zygmund type operator if the following properties are fulfilled:

$$(3) \quad T: L^p(\mathbb{R}^n, E) \rightarrow L^p(\mathbb{R}^n, E)$$
 is bounded for some  $1 < p < \infty$ ,

there exists a locally integrable function  $k$  from  $\mathbb{R}^n \times \mathbb{R}^n \setminus \{(x, x)\}$  into  $\mathcal{L}(E)$  such that

$$(4) \quad Tf(x) = \int k(x, y)f(y) dy$$

for every  $E$ -valued bounded and compactly supported function  $f$  and  $x \notin \text{supp } f$ , and there exists  $\varepsilon > 0$  such that

$$(5) \quad \|k(x, y) - k(x', y)\| \leq C \frac{|x - x'|^\varepsilon}{|x - y|^{n+\varepsilon}}, \quad |x - y| \geq 2|x - x'|.$$

**Remark 1.2.** *It is well known (see [RRT] or [GR]) that in such a case  $T$  is bounded on  $L^q(\mathbb{R}^n, E)$  for any  $1 < q < \infty$ .*

Throughout the literature, after the result on commutators in [CRW], many results appeared in connection with the boundedness of commutators of Calderón-Zygmund type operators and multiplication by a function  $b$  given by  $T_b(f) = bT(f) - T(bf)$  on many different function spaces and on their weighted and vector-valued versions (see [ST1], [ST2], [ST3], [ST4], [ST5]). Also endpoint estimates for the commutator was a topic that attracted several people on different directions (see [CP], [HST], [PP], [P1], [P2], [PT1], [PT2]).

We shall deal in this paper with the unweighted but operator-valued version of the commutators and will give some results about its boundedness on  $L^p(\mathbb{R}^n, E)$  and produce some endpoint estimates.

The following result was shown by C. Segovia and J. L. Torrea (even with some weights and two different Banach spaces).

**Theorem 1.3** ([ST1, Theorem 1]). *Let  $T$  be an  $\mathcal{L}(E)$ -valued Calderón-Zygmund type operator and let  $\ell \rightarrow \tilde{\ell}$  be a correspondence from  $\mathcal{L}(E)$  to  $\mathcal{L}(E)$  such that*

$$(6) \quad \tilde{\ell}T(f)(x) = T(\ell f)(x)$$

and

$$(7) \quad k(x, y)\ell = \tilde{\ell}k(x, y).$$

*If  $b$  is an  $\mathcal{L}(E)$ -valued function such that  $b$  and  $\tilde{b}$  belong to  $BMO(\mathbb{R}^n, \mathcal{L}(E))$  then*

$$T_b(f) = bT(f) - T(bf)$$

*is bounded from  $L^p(\mathbb{R}^n, E) \rightarrow L^p(\mathbb{R}^n, E)$  for all  $1 < p < \infty$ .*

The endpoint estimates of that result were later studied by E. Harboure, C. Segovia and J. L. Torrea (see Theorem A and Theorem 3.1 in [HST]) when  $b$  was assumed to be scalar-valued. From their results one concludes that non-constant scalar valued *BMO* functions do not define bounded commutators from  $L^\infty(\mathbb{R}^n, E)$  to  $BMO(\mathbb{R}^n, F)$  when the kernel of the Calderón-Zygmund type operators are  $\mathcal{L}(E, F)$ -valued. Also it was shown that, in general,  $T_b$  does not map  $H^1(\mathbb{R}^n, E)$  into  $L^1(\mathbb{R}, F)$ .

The aim of this note is to use the techniques developed in the papers [ST1], [HST] to get some extensions for operator-valued  $BMO$ -functions having some commuting properties with the kernel. In particular we show that if  $\|k(x, y)\| \leq \psi(|x - y|^n)$  for certain function  $\psi$  then the commutators of operator-valued  $BMO$  functions and operator-valued Calderón-Zygmund operators map  $L_c^\infty(\mathbb{R}^n, E)$  into  $BMO_\phi(\mathbb{R}^n, E)$  for a function  $\phi$  depending on  $\psi$ . Also we shall see that the commutator is bounded from  $H^1(\mathbb{R}^n, E)$  into  $L_{\text{weak}, \alpha}(\mathbb{R}^n, E)$  for a suitable  $\alpha$  defined from  $\psi$ .

Throughout the paper  $b: \mathbb{R}^n \rightarrow \mathcal{L}(E)$  is locally integrable and  $T$  is a Calderón-Zygmund type operator defined on  $L^p(\mathbb{R}^n, E)$  with a kernel  $k$  satisfying (3), (4) and (5). We write

$$T_b(f)(x) = b(x)(T(f)(x)) - T(bf)(x)$$

where we understand the product  $bf$  as the  $E$ -valued function  $b(y)(f(y))$ .

We shall use the notation  $Q$  for a cube in  $\mathbb{R}^n$ ,  $x_Q$  for its center,  $\ell(Q)$  for the side length,  $\lambda Q$  for a cube centered at  $x_Q$  with side length  $\lambda\ell(Q)$  and  $Q^c = \mathbb{R}^n \setminus Q$ . Finally, as usual,  $C$  stands for a constant that may vary from line to line.

## 2. The results

We improve Theorem 1.3 by realizing that conditions (6) and (7) are not of independent nature. Our basic assumptions throughout the paper are the following ones:

$$\mathbf{(A1)} \quad b(z)k(x, y) = k(x, y)b(z), \quad x, y, z \in \mathbb{R}^n, x \neq y.$$

$$\mathbf{(A2)} \quad b_Q T(e\chi_A)(x) = T(b_Q e\chi_A)(x), \quad x \in Q, A \subseteq Q \text{ measurable, } e \in E.$$

We would like to point out that **(A1)** produces the following cancellation property.

**Lemma 2.1.** *Let  $b$  satisfy **(A1)**, let  $Q, Q'$  be cubes in  $\mathbb{R}^n$  and  $f_1$  and  $f_2$  be compactly supported  $E$ -valued with  $\text{supp } f_1 \subset Q'$  and  $\text{supp } f_2 \subset (Q')^c$ . Then*

$$(8) \quad b_Q T(f_2)(x) = T(b_Q f_2)(x), \quad x \in Q'.$$

$$(9) \quad b_Q T(f_1)(x) = T(b_Q f_1)(x), \quad x \in (Q')^c.$$

*Proof:* Let us show (8). Recall that if  $F \in L^1(\mathbb{R}^n, X)$  and  $\Phi \in \mathcal{L}(X)$  for a given Banach space  $X$  then  $\Phi(\int F(x) dx) = \int \Phi F(x) dx$ . Hence, considering  $X = \mathcal{L}(E)$  and  $\Phi(T) = Tb_Q$  or  $\Phi(T) = b_Q T$  one gets, for  $x \in Q'$ ,

$$\begin{aligned} b_Q T(f_2)(x) &= b_Q \left( \int_{(Q')^c} k(x, y) f_2(y) dy \right) \\ &= \int_{(Q')^c} b_Q k(x, y) f_2(y) dy \\ &= \int_{(Q')^c} \left( \frac{1}{|Q|} \int_Q b(z) dz \right) k(x, y) f_2(y) dy \\ &= \int_{(Q')^c} \left( \frac{1}{|Q|} \int_Q b(z) k(x, y) dz \right) f_2(y) dy \\ &= \int_{(Q')^c} k(x, y) \left( \frac{1}{|Q|} \int_Q b(z) dz \right) f_2(y) dy \\ &= T(b_Q f_2)(x). \end{aligned}$$

(9) follows similarly and it is left to the reader. □

The assumptions **(A1)** and **(A2)** hold true, for instance, in the following cases.

**Example 2.2.** Let  $T, S$  be operators in  $\mathcal{L}(E)$  with  $ST = TS$ . Let  $b(x) = b_0(x)T$  and  $k(x, y) = k_0(x, y)S$  for scalar valued functions  $b_0$  and  $k_0$ .

Hence our results will apply whenever either  $b$  or  $k$  are scalar-valued.

**Example 2.3.** Let  $E$  be a Banach space,  $b_0(x) \in E^*$  and let  $k(x, y)$  be scalar valued function such that  $T$  is bounded on  $L^p(\mathbb{R}^n, E)$ . The case  $T_{b_0}(f) = \langle b_0, T(f) \rangle - T(\langle b_0, f \rangle)$  follows from the operator-valued case by selecting  $e \in E$  and  $b(x)(f) = \langle b_0(x), f \rangle e$  in  $\mathcal{L}(E)$ .

We formulate now the results of the paper. The first one is just a modification of a similar result from **[ST1]** but stated here under slightly weaker assumptions.

**Theorem 2.4.** *Let  $b \in BMO(\mathbb{R}^n, \mathcal{L}(E))$  and let  $T$  be a Calderón-Zygmund type operator defined on  $L^p(\mathbb{R}^n, E)$  where the kernel and  $b$  satisfy **(A1)** and **(A2)**. Then  $T_b$  is bounded on  $L^p(\mathbb{R}^n, E)$  for any  $1 < p < \infty$ .*

Next we analyze the endpoint estimates. We construct a function  $\phi$  for the commutator  $T_b$  to be bounded from  $L_c^\infty(\mathbb{R}^n, E)$  into  $BMO_\phi(\mathbb{R}^n, \mathcal{L}(E))$ .

**Theorem 2.5.** *Let  $T$  be a Calderón-Zygmund type operator with operator-valued kernel  $k$  and assume that*

$$(10) \quad \|k(x, y)\| \leq \psi(|x - y|^n), \quad x \neq y$$

for some  $\psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that  $\int_s^\infty \psi(u) du = \phi(s) < \infty$  for all  $s > 0$ .

If  $b \in BMO(\mathbb{R}^n, \mathcal{L}(E))$  satisfies **(A1)** and that  $T_b$  is bounded on some  $L^p(\mathbb{R}^n, E)$  then  $T_b$  is bounded from  $L_c^\infty(\mathbb{R}^n, E)$  into  $BMO_{1+\phi}(\mathbb{R}^n, E)$ .

We also find a function  $\alpha$  such that the commutator of a function  $b$  in  $BMO(\mathbb{R}^n, \mathcal{L}(E))$  with a Calderón-Zygmund type operator  $T_b$  maps the space  $H^1(\mathbb{R}^n, E)$  into  $L_{\text{weak}, \alpha}(\mathbb{R}^n, E)$ .

**Theorem 2.6.** *Let  $T$  be a Calderón-Zygmund type operator with operator-valued kernel  $k$ . Assume that*

$$(11) \quad \|k(x, y)\| \leq \gamma(|x - y|^n), \quad x \neq y$$

for some decreasing function  $\gamma: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  and

$$(12) \quad \|k(x, y) - k(x, y')\| \leq C \frac{|y - y'|^\varepsilon}{|x - y|^{n+\varepsilon}}, \quad |x - y| \geq 2|y - y'|.$$

If  $b \in BMO(\mathbb{R}^n, \mathcal{L}(E))$  satisfies **(A1)** and  $T_b$  is bounded on some  $L^p(\mathbb{R}^n, E)$  then  $T_b$  is bounded from  $H^1(\mathbb{R}^n, E)$  into  $L_{\text{weak}, \alpha}^1(\mathbb{R}^n, E)$  for  $\alpha(\lambda) = \gamma^{-1}(\|b\|_{BMO}^{-1} \lambda)$ .

As corollaries of these results one obtains the following applications.

**Corollary 2.7** (see [ST1]). *Let  $H$  be the Hilbert transform*

$$H(f)(x) = p.v. \int \frac{f(y)}{x - y} dy,$$

and  $E$  be a UMD space (see [GR]). If  $b \in BMO(\mathbb{R}, \mathcal{L}(E))$  then

- (i)  $H_b$  maps  $L^p(\mathbb{R}, E)$  to  $L^p(\mathbb{R}, E)$  for  $1 < p < \infty$  and
- (ii)  $H_b$  maps  $H^1(\mathbb{R}, E)$  to  $L_{\text{weak}, 1/t}(\mathbb{R}, E)$ .

Although our results are stated in  $\mathbb{R}$ , similar ones work in  $\mathbb{T}$ . In this case we can obtain

**Corollary 2.8** (see [HST]). *Let  $\tilde{H}$  be the conjugate function in the torus*

$$\tilde{H}(f)(x) = p.v. \frac{1}{2\pi} \int \cot\left(\frac{x-y}{2}\right) f(y) dy, \quad x \in [-\pi, \pi]$$

and  $E$  be a UMD space. If  $b \in BMO(\mathbb{R}, \mathcal{L}(E))$  then

- (i)  $H_b$  maps  $L^p(\mathbb{T}, E)$  to  $L^p(\mathbb{T}, E)$  for  $1 < p < \infty$ ,
- (ii)  $H_b$  maps  $H^1(\mathbb{T}, E)$  to  $L_{\text{weak},1/t}(\mathbb{T}, E)$  and
- (iii)  $H_b$  maps  $L^\infty(\mathbb{T}, E)$  to  $BMO_{|\log t|^{-1}}(\mathbb{T}, E)$ .

### 3. Proof of the results

Let us start by showing some consequences from **(A1)** and **(A2)**.

**Lemma 3.1.** *Let  $b$  satisfy **(A1)** and **(A2)**,  $Q$  be a cube in  $\mathbb{R}^n$  and  $f$  be simple  $E$ -valued function. Then*

$$(13) \quad b_Q T(f)(x) = T(b_Q f)(x), \quad x \in Q.$$

*Proof:* Take  $f_1 = f\chi_Q$  and  $f_2 = f - f_1$ . Using Lemma 2.1 one obtains  $b_Q T(f_2)\chi_Q = T(b_Q f_2)\chi_Q$  and **(A2)** shows that  $b_Q T(f_1)\chi_Q = T(b_Q f_1)\chi_Q$ . □

The following useful lemma is essentially included in [HST].

**Lemma 3.2.** *Let  $Q$  be a cube, denote  $Q_j = 2^j Q$  and let  $f$  be compactly supported  $E$ -valued with  $\text{supp } f \subset (2Q)^c$ . Then there exists  $C > 0$  such that*

$$(14) \quad \|T(f)(x) - T(f)(x')\| \leq C \frac{|x-x'|^\varepsilon}{\ell(Q)^\varepsilon} \sum_{j=2}^\infty \frac{2^{-j\varepsilon}}{|Q_j|} \int_{Q_j} \|f(y)\| dy, \quad x, x' \in Q.$$

*Proof:* Using (4) and (5) one has

$$\begin{aligned}
\|T(f)(x) - T(f)(x')\| &\leq \int_{(2Q)^c} \|k(x, y) - k(x', y)\| \|f(y)\| dy \\
&\leq C|x - x'|^\varepsilon \int_{(2Q)^c} \frac{\|f(y)\|}{|x - y|^{n+\varepsilon}} dy \\
&\leq C|x - x'|^\varepsilon \sum_{j=1}^{\infty} \int_{Q_{j+1} - Q_j} \frac{\|f(y)\|}{|x - y|^{n+\varepsilon}} dy \\
&\leq C|x - x'|^\varepsilon \sum_{j=2}^{\infty} \frac{1}{\ell(Q_j)^{n+\varepsilon}} \int_{Q_j} \|f(y)\| dy \\
&\leq C \frac{|x - x'|^\varepsilon}{\ell(Q)^\varepsilon} \sum_{j=2}^{\infty} 2^{-j\varepsilon} \frac{1}{|Q_j|} \int_{Q_j} \|f(y)\| dy. \quad \square
\end{aligned}$$

*Proof of Theorem 2.4:* Let  $f$  be a simple  $E$ -valued function. Let  $Q$  be a cube,  $f_1 = f\chi_{2Q}$  and  $f_2 = f - f_1$ . Put  $c_Q = T((b_Q - b)f_2)(x_Q)$ .

For each  $x \in Q$  one has, applying Lemma 3.1,

$$T_b f(x) - c_Q = \sum_{i=1}^3 \sigma_i(x)$$

where

$$\sigma_1(x) = (b - b_Q)Tf(x),$$

$$\sigma_2(x) = T((b_Q - b)f_1)(x)$$

and

$$\sigma_3(x) = T((b_Q - b)f_2)(x) - T((b_Q - b)f_2)(x_Q).$$

Observe that for  $1 < q < \infty$  and  $1/q + 1/q' = 1$  we can write

$$\frac{1}{|Q|} \int_Q \|\sigma_1(x)\| dx \leq \text{osc}_{q'}(b, Q) \left( \frac{1}{|Q|} \int_Q \|Tf(x)\|^q dx \right)^{1/q}.$$



For any  $q > q_1 > 1$  one can use Remark 1.2, for  $1/r + 1/q = 1/q_1$ , to obtain

$$\begin{aligned} \frac{1}{|Q|} \int_Q \|\sigma_2(x)\| dx &\leq \left( \frac{1}{|Q|} \int_Q \|T(b_Q - b)f_1(x)\|^{q_1} dx \right)^{1/q_1} \\ &\leq C \|T\|_{\mathcal{L}(L^{q_1}(\mathbb{R}^n, E))} \left( \frac{1}{|Q|} \int_Q \|(b - b_Q)f_1(x)\|^{q_1} dx \right)^{1/q_1} \\ &\leq C \|T\|_{\mathcal{L}(L^{q_1}(\mathbb{R}^n, E))} \operatorname{osc}_r(b, Q) \left( \frac{1}{|Q|} \int_Q \|f(x)\|^q dx \right)^{1/q}. \end{aligned}$$

Using Lemma 3.2, and taking into account that  $\|b_Q - b_{2Q}\| \leq C \operatorname{osc}_{q_1}(b, 2Q)$ , we also can estimate

$$\begin{aligned} \|\sigma_3(x)\| &\leq C \sum_{j=2}^{\infty} 2^{-j\varepsilon} \frac{1}{|Q_j|} \int_{Q_j} \|(b(y) - b_Q)f(y)\| dy \\ &\leq C \sum_{j=2}^{\infty} 2^{-j\varepsilon} \left( \frac{1}{|Q_j|} \int_{Q_j} \|b(y) - b_Q\|^{q'} dy \right)^{1/q'} \left( \frac{1}{|Q_j|} \int_{Q_j} \|f(y)\|^q dy \right)^{1/q} \\ &\leq C \sum_{j=2}^{\infty} 2^{-j\varepsilon} \left( \sum_{k=2}^j \operatorname{osc}_{q'}(b, Q_k) \right) \left( \frac{1}{|Q_j|} \int_{Q_j} \|f(y)\|^q dy \right)^{1/q} \\ &\leq C \sup_{j \geq 2} \left( \frac{1}{|Q_j|} \int_{Q_j} \|f(y)\|^q dy \right)^{1/q} \left( \sum_{j=2}^{\infty} 2^{-j\varepsilon} \left( \sum_{k=2}^j \operatorname{osc}_{q'}(b, Q_k) \right) \right) \\ &\leq C \|b\|_{BMO} \sup_{j \geq 2} \left( \frac{1}{|Q_j|} \int_{Q_j} \|f(y)\|^q dy \right)^{1/q} \sum_j j 2^{-j\varepsilon}. \end{aligned}$$

Hence, combining the previous estimates, one obtains

$$T_b(f)^\#(x) \leq C \|b\|_{BMO} (M_q(Tf)(x) + M_q(f)(x)).$$

Now, for a given  $1 < p < \infty$ , select  $1 < q < p$  and apply (2), which, combined with the boundedness of  $T$  on  $L^p(\mathbb{R}^n, E)$ , shows that  $\|T_b(f)^\#\|_{L^p(\mathbb{R}^n)} \leq C \|f\|_{L^p(\mathbb{R}^n, E)}$ . Now use the vector-valued analogue of Fefferman-Stein's result (see [FS], [RRT]) to obtain that  $\|T_b(f)\|_{L^p(\mathbb{R}^n, E)} \leq C \|f\|_{L^p(\mathbb{R}^n, E)}$ . □

*Proof of Theorem 2.5:* As in the previous theorem, let  $f$  be a simple  $E$ -valued function. Let  $Q$  be a cube,  $f_1 = f\chi_{2Q}$ ,  $f_2 = f - f_1$  and  $c_Q = T((b_Q - b)f_2)(x_Q)$ . Now, using Lemma 2.1, we write

$$T_b f(x) = T_b(f_1)(x) + (b(x) - b_Q)T(f_2)(x) + T((b_Q - b)f_2)(x).$$

Denote now

$$\begin{aligned}\sigma_1(x) &= T_b(f_1)(x), \\ \sigma_2(x) &= (b(x) - b_Q)T(f_2)(x), \\ \sigma_3(x) &= T((b_Q - b)f_2)(x) - T((b_Q - b)f_2)(x_Q).\end{aligned}$$

Hence  $T_b f - c_Q = \sum_{i=1}^3 \sigma_i$ . Note that the boundedness of  $T_b$  on  $L^p(\mathbb{R}^n, E)$  gives

$$\frac{1}{|Q|} \int_Q \|\sigma_1(x)\| dx \leq C \|T_b\|_{\mathcal{L}(L^p)} \left( \frac{1}{|2Q|} \int_{2Q} \|f(x)\|^p dx \right)^{1/p} \leq C \|f\|_\infty.$$

On the other hand

$$\begin{aligned}\frac{1}{|Q|} \int_Q \|\sigma_2(x)\| dx &\leq \frac{1}{|Q|} \int_Q \|b(x) - b_Q\| \left\| \int_{(2Q)^c} k(x, y) f(y) dy \right\| dx \\ &\leq C \frac{1}{|Q|} \int_Q \|b(x) - b_Q\| \left( \int_{(2Q)^c} \psi(|x-y|^n) \|f(y)\| dy \right) dx \\ &\leq C \|f\|_\infty \frac{1}{|Q|} \int_Q \|b(x) - b_Q\| \left( \int_{|u|>\ell(Q)} \psi(|u|^n) du \right) dx \\ &\leq C \|f\|_\infty \left( \frac{1}{|Q|} \int_Q \|b(x) - b_Q\| dx \right) \left( \int_{\ell(Q)}^\infty r^{n-1} \psi(r^n) dr \right) \\ &\leq C \|f\|_\infty \|b\|_{BMO} \left( \int_{|Q|}^\infty \psi(t) dt \right).\end{aligned}$$

Finally Lemma 3.2 gives immediately

$$\frac{1}{|Q|} \int_Q \|\sigma_3(x)\| dx \leq C \|b\|_{BMO} \|f\|_\infty.$$

This allows us to conclude the estimate

$$\frac{1}{|Q|} \int_Q \|T_b f(x) - c_Q\| dx \leq C \|f\|_\infty (1 + \phi(|Q|)).$$

This shows that  $T_b$  maps  $L_c^\infty(\mathbb{R}^n, E)$  into  $BMO_{1+\phi}(\mathbb{R}^n, E)$ .  $\square$

*Proof of Theorem 2.6:* Let  $a$  be an  $E$ -valued atom supported on  $Q$ . Using Lemma 2.1 again we can write

$$T_b a(x) = \chi_{2Q}(x)T_b(a)(x) + \chi_{(2Q)^c}(x)(b(x) - b_Q)T(a)(x) + \chi_{(2Q)^c}(x)T((b_Q - b)a)(x).$$

Denote now

$$\begin{aligned} \sigma_1(x) &= \chi_{2Q}(x)T_b(a)(x), \\ \sigma_2(x) &= \chi_{(2Q)^c}(x)(b(x) - b_Q)T(a)(x), \\ \sigma_3(x) &= \chi_{(2Q)^c}(x)T((b_Q - b)a)(x). \end{aligned}$$

Now, using the boundedness of  $T_b$  on  $L^p(\mathbb{R}^n, E)$ ,

$$\begin{aligned} \int_{\mathbb{R}^n} \|\sigma_1(x)\| dx &\leq C|Q|^{1/p'} \|T_b(a)\|_{L^p(\mathbb{R}^n, E)} \\ &\leq C\|T_b\|_{\mathcal{L}(L^p)}|Q| \left( \frac{1}{|Q|} \int_Q \|a(x)\|^p dx \right)^{1/p} \\ &\leq C\|T_b\|_{\mathcal{L}(L^p)}. \end{aligned}$$

Also we have

$$\begin{aligned} \int_{\mathbb{R}^n} \|\sigma_2(x)\| dx &\leq \int_{(2Q)^c} \|b(x) - b_Q\| \left\| \int_Q k(x, y)a(y) dy \right\| dx \\ &\leq \int_{(2Q)^c} \|b(x) - b_Q\| \left\| \int_Q (k(x, y) - k(x, x_Q))a(y) dy \right\| dx \\ &\leq C \int_{(2Q)^c} \|b(x) - b_Q\| \left( \int_Q \frac{|y - x_Q|^\varepsilon}{|x - y|^{n+\varepsilon}} \|a(y)\| dy \right) dx \\ &\leq C \frac{\ell(Q)^\varepsilon}{|Q|} \int_Q \left( \int_{(2Q)^c} \frac{\|b(x) - b_Q\|}{|x - y|^{n+\varepsilon}} dx \right) dy \\ &\leq C \frac{\ell(Q)^\varepsilon}{|Q|} \int_Q \left( \sum_{j=2}^\infty \frac{1}{\ell(Q_j)^{n+\varepsilon}} \int_{Q_j - Q_{j-1}} \|b(x) - b_Q\| dx \right) dy \\ &\leq C \left( \sum_{j=2}^\infty 2^{-j\varepsilon} \frac{1}{|Q_j|} \int_{Q_j} \|b(x) - b_Q\| dx \right) \leq C\|b\|_{BMO}. \end{aligned}$$

Now decompose  $\sigma_3 = \sigma_{3,1} + \sigma_{3,2}$  where

$$\sigma_{3,1}(x) = \chi_{(2Q)^c}(x) \int_Q (k(x, y) - k(x, x_Q))(b_Q - b(y))a(y) dy,$$

$$\sigma_{3,2}(x) = \chi_{(2Q)^c}(x)k(x, x_Q) \int_Q b(y)a(y) dy.$$

Therefore

$$\begin{aligned} \int_{\mathbb{R}^n} \|\sigma_{3,1}(x)\| dx &\leq \int_{(2Q)^c} \int_Q \|k(x, y) - k(x, x_Q)\| \|b_Q - b(y)\| \|a(y)\| dy dx \\ &\leq \int_{(2Q)^c} \frac{\ell(Q)^\varepsilon}{|Q|} \left( \int_Q \frac{\|b_Q - b(y)\|}{|x - y|^{n+\varepsilon}} dy \right) dx \\ &\leq \frac{\ell(Q)^\varepsilon}{|Q|} \int_Q \|b_Q - b(y)\| \left( \int_{(2Q)^c} \frac{dx}{|x - y|^{n+\varepsilon}} \right) dy \\ &\leq \frac{\ell(Q)^\varepsilon}{|Q|} \int_Q \|b_Q - b(y)\| \left( \int_{|x|>\ell(Q)} \frac{dx}{|x|^{n+\varepsilon}} \right) dy \leq C \|b\|_{BMO}. \end{aligned}$$

Since  $\|\int_Q b(y)a(y) dy\| \leq \frac{1}{|Q|} \int_Q \|b(y) - b_Q\| dy$  we can estimate

$$\begin{aligned} \sigma_{3,2}(x) &\leq \chi_{(2Q)^c}(x) \|k(x, x_Q)\| \|b\|_{BMO} \\ &\leq \|b\|_{BMO} \chi_{(2Q)^c}(x) \gamma(|x - x_Q|^n). \end{aligned}$$

Therefore one gets

$$\begin{aligned} |\{x : \sigma_{3,2}(x) > \lambda\}| &\leq |\{x \in (2Q)^c : \gamma(|x - x_Q|^n) > \|b\|_{BMO}^{-1} \lambda\}| \\ &= |\{x \in (2Q)^c : |x - x_Q| < [\gamma^{-1}(\|b\|_{BMO}^{-1} \lambda)]^{1/n}\}|. \end{aligned}$$

This gives the estimate  $|\{x : \sigma_{3,2}(x) > \lambda\}| \leq \psi^{-1}(\|b\|_{BMO}^{-1} \lambda) = \alpha(\lambda)$ .  
The proof is then easily concluded.  $\square$

### References

- [B] S. BLOOM, A commutator theorem and weighted BMO, *Trans. Amer. Math. Soc.* **292**(1) (1985), 103–122.
- [CRW] R. R. COIFMAN, R. ROCHBERG, AND G. WEISS, Factorization theorems for Hardy spaces in several variables, *Ann. of Math. (2)* **103**(3) (1976), 611–635.

- [CP] D. CRUZ-URIBE AND C. PÉREZ, Two-weight, weak-type norm inequalities for fractional integrals, Calderón-Zygmund operators and commutators, *Indiana Univ. Math. J.* **49(2)** (2000), 697–721.
- [FS] C. FEFFERMAN AND E. M. STEIN,  $H^p$  spaces of several variables, *Acta Math.* **129(3–4)** (1972), 137–193.
- [GR] J. GARCÍA-CUERVA AND J. L. RUBIO DE FRANCIA, “*Weighted norm inequalities and related topics*”, North-Holland Mathematics Studies **116**, Notas de Matemática [Mathematical Notes] 104, North-Holland Publishing Co., Amsterdam, 1985.
- [HST] E. HARBOURE, C. SEGOVIA, AND J. L. TORREA, Boundedness of commutators of fractional and singular integrals for the extreme values of  $p$ , *Illinois J. Math.* **41(4)** (1997), 676–700.
- [P1] C. PÉREZ, Endpoint estimates for commutators of singular integral operators, *J. Funct. Anal.* **128(1)** (1995), 163–185.
- [P2] C. PÉREZ, Sharp estimates for commutators of singular integrals via iterations of the Hardy-Littlewood maximal function, *J. Fourier Anal. Appl.* **3(6)** (1997), 743–756.
- [PP] C. PÉREZ AND G. PRADOLINI, Sharp weighted endpoint estimates for commutators of singular integrals, *Michigan Math. J.* **49(1)** (2001), 23–37.
- [PT1] C. PÉREZ AND R. TRUJILLO-GONZÁLEZ, Sharp weighted estimates for multilinear commutators, *J. London Math. Soc. (2)* **65(3)** (2002), 672–692.
- [PT2] C. PÉREZ AND R. TRUJILLO-GONZÁLEZ, Sharp weighted estimates for vector-valued singular integral operators and commutators, *Tohoku Math. J. (2)* **55(1)** (2003), 109–129.
- [RRT] J. L. RUBIO DE FRANCIA, F. J. RUIZ, AND J. L. TORREA, Calderón-Zygmund theory for operator-valued kernels, *Adv. in Math.* **62(1)** (1986), 7–48.
- [ST1] C. SEGOVIA AND J. L. TORREA, Vector-valued commutators and applications, *Indiana Univ. Math. J.* **38(4)** (1989), 959–971.
- [ST2] C. SEGOVIA AND J. L. TORREA, A note on the commutator of the Hilbert transform, *Rev. Un. Mat. Argentina* **35** (1989), 259–264 (1991).
- [ST3] C. SEGOVIA AND J. L. TORREA, Weighted inequalities for commutators of fractional and singular integrals, Conference on Mathematical Analysis (El Escorial, 1989), *Publ. Mat.* **35(1)** (1991), 209–235.
- [ST4] C. SEGOVIA AND J. L. TORREA, Commutators of Littlewood-Paley sums, *Ark. Mat.* **31(1)** (1993), 117–136.

- [ST5] C. SEGOVIA AND J. L. TORREA, Higher order commutators for vector-valued Calderón-Zygmund operators, *Trans. Amer. Math. Soc.* **336(2)** (1993), 537–556.
- [SW] E. M. STEIN AND G. WEISS, “*Introduction to Fourier analysis on Euclidean spaces*”, Princeton Mathematical Series **32**, Princeton University Press, Princeton, N.J., 1971.

Departament de Matemàtiques  
Universitat de València  
Burjassot 46100, València  
Spain  
*E-mail address:* oscar.blasco@uv.es

Rebut el 22 de gener de 2008.