

CATEGORICAL METHODS IN GRADED RING THEORY

ANGEL DEL RÍO*

A la memoria de Pere Menal

Abstract

Let G be a group, R a G -graded ring and X a right G -set. We study functors between categories of modules graded by G -sets, continuing the work of [M]. As an application we obtain generalizations of Cohen-Montgomery Duality Theorems by categorical methods. Then we study when some functors introduced in [M] (which generalize some functors occurring in [D1], [D2] and [NRV]) are separable. Finally we obtain an application to the study of the weak dimension of a group graded ring.

Introduction and Notation

The study of group-graded rings has been the ground for the research of many authors last years. Different methods have been elaborated to investigate properties of these rings. One of the most successful tools is that given by Cohen-Montgomery Duality Theorems [CM]. Furthermore, some categorical methods have been introduced by other authors which have been very useful. For instance the study of separable functors, introduced in [NVV], has caused nice theorems in the field.

Quite often the methods invented have been introduced in the particular case when the grading group is finite. This is the case of the two methods mentioned above. Then some efforts have produced different approaches to the general case. See [Q], [B1], [B2], [AN] and [NRV] for some extensions of Cohen-Montgomery Duality Theorem and [Ra] for separable functors.

In [NRV] the authors introduce a category $gr - (R, X, G)$ associated to a G -graded ring R (G being a group) and a right G -set X . The most

*Partially supported by CICYT PB90-0300.

important example appears when X is the set G/H of right H -cosets. This category appears to be useful in the study of the ring R .

Let G and G' be two groups, R a G -graded ring, R' a G' -graded ring, X a right G -set and X' a right G' -set. In [M] covariant adjoint functors between the categories $gr - (R, X, G)$ and $gr - (R', X', G')$ are studied, generalizing some results of [Rí2].

Let $\rho : R \rightarrow R'$ be a ring homomorphism, $\xi : X \rightarrow X'$ a map and $\gamma : G \rightarrow G'$ a group homomorphism and assume that for every $g \in G$ and $x \in X$, $\rho(R_g) \subseteq R_{\gamma(g)}$ and $\xi(x)\gamma(g) = \xi(xg)$. Associated to $T = (\rho, \xi, \gamma)$ there is a pair of adjoint functors $T^* = gr - (R, X, G) \rightarrow gr - (R', X', G')$ and $T_* : gr - (R', X', G') \rightarrow gr - (R, X, G)$ for which many functors occurring in Graded Clifford Theory (cf. [D1], [D2], [GN], [NV2]) as well as induced and coinduced functors (cf. [MN]) are particular cases.

After some definitions and notations we recall in Section 1 results of [M] on covariant adjoint functors and point out some similar fact on contravariant adjoint functors which extend some results of [MR]. We finish Section 1 with some applications to the functors T^* and T_* associated to a context $T = (\rho, \xi, \gamma)$ as in the previous paragraph.

The results of Section 1 set up the framework to give a description of all the equivalences and Morita dualities (in the sense of Colby-Fuller [CF]) between categories of the type $gr - (R, X, G)$ using similar methods as in [Rí2] and [MR]. This is done without proofs in Section 2.

In Section 3 we apply the tools of Section 1 to extend Cohen-Montgomery Duality Theorems to our setting by using categorical methods. Namely, we show how the category $gr - (R, X, G)$ can be seen as the full subcategory of a category of right modules generated by an ideal. Then we prove new versions of Cohen-Montgomery Duality Theorems for actions and coactions. All the previous versions of these Duality Theorems can be obtained as particular cases. Furthermore, some equivalences of categories are given which will be useful in applications.

Section 4 is devoted to study when T^* and T_* are separable. We give some arithmetical tests for the matter. When $R = R'$ and ρ is the identity map on R , T^* and T_* are isomorphic to the functors T_ξ and S^ξ introduced in [NRV] and, in this case, T_ξ is always separable while the test of separability for S^ξ can be simplified.

We finish the paper with an application to the weak dimension of a group graded ring.

The author would like to express his gratitude to Claudia Menini for her very interesting suggestions.

All rings are supposed to be associative with unit and all modules are

unital.

For a ring R , $\text{mod-}R$ will denote the category of right R -modules and $R\text{-mod}$ the category of left R -modules.

The notation M_R (resp. ${}_R M$) will be used to emphasize that M is a right (resp. left) R -module.

In the sequel G will stand for a multiplicative group and $R = \bigoplus_{g \in G} R_g$ for a G -graded ring. The unit of G will be denoted by e while 1 will denote the unit of R .

Let X be a right (resp. left) G -set. For any $g \in G$ and any $x \in X$, xg (resp. gx) denotes the action of g on x .

Following the notation introduced in [NRV], $gr - (R, X, G)$ will denote the category of X -graded right R -module. That is, the objects of $gr - (R, X, G)$ are right R -modules M with a decomposition into additive subgroups $M = \bigoplus_{x \in X} M_x$ such that for every $g \in G$ and every $x \in X$, $M_x R_g \subseteq M_{xg}$. Given M, N objects in $gr - (R, X, G)$, the set of morphisms from M to N in $gr - (R, X, G)$ is

$$\begin{aligned} \text{Hom}_{gr-(R,X,G)}(M, N) &= \\ &= \{f \in \text{Hom}_R(M, N) \mid f(M_x) \subseteq N_x \text{ for every } x \in X\}. \end{aligned}$$

Similarly, if X is a left G -set then $(G, X, R) - gr$ will denote the category of X -graded left R -modules. If $M \in (G, X, R) - gr$ and $x \in X$, then the x -th component of M will be denote by ${}_x M$.

If G is considered as a right G -set by regular action, then $gr - (R, G, G)$ is just $gr - R$. If X is a singleton, then $gr - (R, X, G)$ is $\text{mod-}R$.

Let H be a subgroup of G and consider G acting on $G/H = \{Ha \mid a \in G\}$ by $(Ha)g = Hag$, then $gr - (R, G/H, G)$ is denoted by $gr - (R, G/H)$.

Given $M \in gr - (R, X, G)$ (resp. $M \in (G, X, R) - gr$), $m \in M$ and $x \in X$, m_x (resp. ${}_x m$) will denote the x -th homogeneous component of m . That is, m_x is defined by $m = \sum_{x \in X} m_x$ with $m_x \in M_x$ for every $x \in X$. If F is a subset of X , then $M_F = \bigoplus_{x \in F} M_x$ and $m_F = \sum_{x \in F} m_x$.

The support of an X -graded right R -module is defined to be the set $\text{Supp}(M) = \{x \in X \mid M_x \neq 0\}$. Also the support of $m \in M$ is the set $\text{Supp}(m) = \{x \in X \mid m_x \neq 0\}$.

When we refer to the support or the homogeneous component of degree $g \in G$ of R (or of an element of R) we will consider R as a G -graded right (or left) R -module.

If X is a right G -set then X has a canonical structure of left G -set as follows: For any $g \in G$, $x \in X$, $gx = xg^{-1}$. We will refer to both right

and left action without specific mention on the side when it is clear from the context.

Given $M \in gr - R$ and $x \in X$, the $x - th$ suspension of M is defined as the X -graded right R -module $M(x)$ which is equal to M as right R -module with the grading $M(x)_y = \bigoplus_{xg=y} M_g$. If $M \in R - gr$, X is a left G -set and $x \in X$, then the $x - th$ suspension of M is the X -graded left R -module $(x)M$ which is equal to M as a left R -module with the X -grading ${}_y(x)M = \bigoplus_{gx=y} M_g$.

1. Adjoint functors

Throughout this section G and G' will be two groups, X a right G -set and X' a right G' -set. Moreover R will be a G -graded ring and R' a G' -graded ring.

In this first section we will give an explicit description of the covariant functors between $gr - (R, X, G)$ and $gr - (R', X', G')$ and the contravariant functors between $(G, X, R) - gr$ and $gr - (R', X', G')$. This description has been obtained in [Rf2], for the covariant case, when $X = G$ and $X' = G'$ with regular action, and generalized for arbitrary G -sets in [M]. For the contravariant case, when $X = G$ and $X' = G'$, see [MR].

In the last part of the section we consider some instances of adjoint functors introduced in [M].

Definition. Given $M \in gr - (R, X, G)$ and $N \in (G, X, R) - gr$, $M \hat{\otimes}_R N$ will denote the additive subgroup of $M \otimes_R N$ generated by the elements of the form $m \otimes n$ where $x \in X$, $m \in M_x$ and $n \in {}_xN$.

Lemma 1.1. (a) $R(x) \hat{\otimes}_R N \cong {}_xN$ for every $N \in (G, X, R) - gr$. Moreover this isomorphism is natural in both variables.

(b) Let $\{M_i | i \in I\}$ be a family of objects in $gr - (R, X, G)$ and $N \in (G, X, R) - gr$, then $(\bigoplus_{i \in I} M_i) \hat{\otimes}_R N \cong \bigoplus_{i \in I} (M_i \hat{\otimes}_R N)$.

Proof: (a) Let $\Phi : R(x) \otimes_R N \rightarrow N$ be the canonical isomorphism. If $r \in R_g \cap R(x)_y$ and $n \in {}_yN$ then $xg = y$ and hence $x = gy$. Therefore $\Phi(r \otimes n) = rn \in {}_xN$. Thus $\Phi(R(x) \hat{\otimes}_R N) \subseteq {}_xN$. On the other hand, if $n \in {}_xN$, then $n = \Phi(1 \otimes n)$ and $1 \otimes n \in R(x) \hat{\otimes}_R N$ because $1 \in R(x)_x$.

(b) Straightforward. ■

Definition. Let Z be a G - G' -set. A Z -graded R - R' -bimodule is an R - R' -bimodule M with a Z -grading, $M = \bigoplus_{z \in Z} M_z$ such that $R_g M_z R'_{g'} \subseteq$

$M_{gzg'}$ for every $g \in G, z \in Z$ and $g' \in G'$.

The set $X \times X'$ has a canonical structure of G - G' -set given by:

$$g(x, x')g' = (gx, x'g') \quad (g \in G, x \in X, x' \in X', g' \in G').$$

Let $P = \bigoplus_{(x, x') \in X \times X'} P_{(x, x')}$ be an $X \times X'$ -graded R - R' -bimodule.

For every $x \in X, {}_xP = \bigoplus_{x' \in X'} P_{(x, x')}$ is a submodule of $P_{R'}$ and $({}_xP)_{x'} = P_{(x, x')}$ ($x' \in X'$) defines an X' -grading in ${}_xP$ such that ${}_xP \in gr - (R', X', G')$. Moreover P is the coproduct of the family $\{{}_xP | x \in X\}$ in $gr - (R', X', G')$.

Similarly for every $x' \in X', P_{x'} = \bigoplus_{x \in X} P_{(x, x')} \in (G, X, R) - gr$ and P is the coproduct of $\{P_{x'} | x' \in X'\}$ in $(G, X, R) - gr$.

We are going to define three functors associated to the $X \times X'$ -graded R - R' -bimodule P .

The functor $-\hat{\otimes}_R P : gr - (R, X, G) \rightarrow gr - (R', X', G')$.

It associates to $M \in gr - (R, X, G)$ the right R' -module $(-\hat{\otimes}_R P)(M) = M \hat{\otimes}_R P$, considered as a submodule of $(M \oplus_R P)_R$, with the X' -grading $M \hat{\otimes}_R P_{x'} = M \hat{\otimes}_R P_{x'} (x' \in X')$.

The functor $-\hat{\otimes}_R P$ associates to $f \in \text{Hom}_{gr - (R, X, G)}(M, N)$ the map $f \hat{\otimes}_R P : m \otimes p \mapsto f(m) \otimes p$.

The functor $H(P_{R'}, -) : gr - (R', X', G') \rightarrow gr - (R, X, G)$.

It associates to $M \in gr - (R', X', G')$ the right R -module

$$H(P_{R'}, M) = \{f \in \text{Hom}_{gr - (R', X', G')}(P, M) | f({}_xP) = 0 \text{ for almost all } x \in X\}$$

considered as a submodule of $\text{Hom}_{R'}(P, M)_R$ with the X -grading:

$$H(P_{R'}, M)_x = \{f \in \text{Hom}_{gr - (R', X', G')}(P, M) | f({}_yP) = 0 \text{ for all } y \in X - \{x\}\} (x \in X).$$

The functor $H(P_R, -)$ associates to $f \in \text{Hom}_{gr - (R', X', G')}(M', N')$, the map $H(P_R, f) : H(P_{R'}, M) \rightarrow H(P_{R'}, N)$ given by: $H(P_R, f)(\varphi) = f \circ \varphi$ for all $\varphi \in H(P_{R'}, M)$.

The functor $H(-, {}_R P) : (G, X, R) - gr \rightarrow gr - (R', X', G')$.

It associates to $M \in (G, X, R) - gr$, the right R' -module

$$H(M, {}_R P) = \\ = \{f \in \text{Hom}_{(G, X, R) - gr}(M, P) \mid f(m) \subseteq P_F \text{ for some finite subset } F \subseteq X'\}$$

considered as a submodule of $\text{Hom}_R(M, P)_{R'}$ with the X' -grading:

$$H(M, {}_R P)_{x'} = \{f \in \text{Hom}_{(G, X, R) - gr}(M, P) \mid f(m) \subseteq P_{x'}\} \quad (x' \in X').$$

The functor $H(-, {}_R P)$ associates to $f \in \text{Hom}_{(G, X, R) - gr}(M, N)$, the map $H(f, {}_R P) : H(N, {}_R P) \rightarrow H(M, {}_R P)$, given by: $H(f, {}_R P)(\varphi) = \varphi \circ f$ for all $\varphi \in H(N, {}_R P)$.

It is left to the reader to check that the above three functors are well defined. Similarly there are two covariant functors $P \hat{\otimes}_{R'} - : (G', X', R') - gr \rightarrow (G, X, R) - gr$ and $H({}_R P, -) : (G, X, R) - gr \rightarrow (G', X', R') - gr$ and a contravariant functor $H(-, P_{R'}) : gr - (R', X', G') \rightarrow (G, X, R) - gr$.

If \mathcal{C} is a category, then \mathcal{C}^{op} will denote the dual category.

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ be two contravariant functors. Then we can consider $F : \mathcal{C} \rightarrow \mathcal{D}^{\text{op}}$ and $G : \mathcal{D}^{\text{op}} \rightarrow \mathcal{C}$ as covariant functors. We say that (F, G) is a pair of adjoint contravariant functors on the right if $F : \mathcal{C} \rightarrow \mathcal{D}^{\text{op}}$ is left adjoint of $G : \mathcal{D}^{\text{op}} \rightarrow \mathcal{C}$.

Proposition 1.2. *Let P be an R - R' -graded bimodule.*

- (a) [M] *The functor $- \hat{\otimes}_R P$ is left adjoint of $H(P_{R'}, -)$.*
- (b) *$(H(-, {}_R P), H(-, P_{R'}))$ is a pair of adjoint contravariant functors on the right.*

Proof: See [M] for (a). In [MR] there is a proof of (b) in the particular case $G = X$ and $G' = X'$ with regular actions. The same method works in the general case. ■

For every $r \in R$ let $\lambda_r : R \rightarrow R$ be the left multiplication-by- r map and $\chi_r : R \rightarrow R$ the right multiplication-by- r map. The following lemma is obvious:

Lemma 1.3. *Let $r \in R_g$ and $x \in X$. Then*

- (a) $\lambda_r \in \text{Hom}_{gr - (R, X, G)}(R(x), R(gx))$.
- (b) $\chi_r \in \text{Hom}_{(G, X, R) - gr}((gx)R, (x)R)$.

Definition 1.4. Let $F : gr - (R, X, G) \rightarrow gr - (R', X', G')$ be a covariant functor. The $X \times X'$ -graded R - R' -bimodule associated to F is the right R' -module $P = \bigoplus_{x \in X} F(R(x))$ with the following structure of left R -module:

$$rp = [F(\lambda_r)](p) \quad (r \in R_g, p \in F(R(x)))$$

and the $X \times X'$ -grading:

$$P_{(x,x')} = F(R(x))_{x'} \quad (x \in X, x' \in X').$$

Let now $F : (G, X, R) - gr \rightarrow gr - (R', X', G')$ be a contravariant functor. The $X \times X'$ -graded R - R' -bimodule associated to F is the right R' -module $P = \bigoplus_{x \in X} F((x)R)$ with the following structure of left R -module

$$rp = [F(\chi_r)](p) \quad (r \in R_g, p \in F((x)R))$$

and the $X \times X'$ -grading

$$P_{(x,x')} = F((x)R)_{x'} \quad (x \in X, x' \in X').$$

Lemma 1.5. *Let $M \in (G, X, R) - gr$. Then $\text{Hom}_{(G, X, R) - gr}((x)R, M) \cong {}_xM$ and this isomorphism is natural in each variable.*

Proof: For any $m \in M$ let $\chi_m : R \rightarrow M$ be the right-multiplication-by- r map. For every $m \in M$, $\chi_m \in \text{Hom}_{(G, X, R) - gr}((x)R, {}_R M)$ if and only if $m \in {}_xM$. Moreover, for every $f \in \text{Hom}_{(G, X, R) - gr}(R(x), P)$, $f = \chi_{f(1)}$. Therefore the map $m \mapsto \chi_m$ is an isomorphism between ${}_xM$ and $\text{Hom}_{(G, X, R) - gr}((x)R, M)$. The naturality is left to the reader. ■

Now we are ready to describe all covariant and contravariant functors between categories of modules graded by G -sets.

Proposition 1.6. (a) [M] $F : gr - (R, X, G) \rightarrow gr - (R', X', G')$ is a left adjoint functor if and only if there exists an $X \times X'$ -graded R - R' -bimodule P such that $F \cong - \hat{\otimes}_R P$. Moreover, in this case P is isomorphic to the $X \times X'$ -graded R - R' -bimodule associated to F .

(b) $F : gr - (R', X', G') \rightarrow gr - (R, X, G)$ is a right adjoint functor if and only if there exists an $X \times X'$ -graded R - R' -bimodule P such that $F \cong H(P_{R'}, -)$.

(c) $F : (G, X, R) - gr \rightarrow gr - (R', X', G')^{op}$ is a left adjoint functor if and only if there exists an $X \times X'$ -graded R - R' -bimodule P such that

$F \cong H(-, {}_R P)$. Moreover, in this case, P is isomorphic to the $X \times X'$ -graded R - R' -bimodule associated to F .

Proof: (a) and (b) have been proved in [M]. For a proof of (c), an argument as in the proof of Proposition 2.1 of [MR] works, by using Lemma 1.5. ■

Now we focus our attention on some particular functors introduced in [M]. Some examples of these functors can be found in [D1], [D2], [NRV] and [MN].

Let $\rho : R \rightarrow R'$ be a homomorphism of rings, $\xi : X \rightarrow X'$ a map and $\gamma : G \rightarrow G'$ a group homomorphism. Assume that the following conditions hold:

$$\rho(R_g) \subseteq R_{\gamma(g)} \quad \xi(xg) = \xi(x)\gamma(g) \quad (x \in X, g \in G).$$

Set $T = (\rho, \xi, \gamma)$ (see [M]). For any $M \in gr - (R, X, G)$, let $T^*(M) = M \otimes_R R'$ and for every $x' \in X'$, let $T^*(M)_{x'}$ be the additive subgroup of $T^*(M)$ generated by the elements of the form $m \otimes r'$ with $m \in M_{x'}$, $r \in R'_g$ and $\xi(x)g' = x'$.

For every $f \in \text{Hom}_{gr - (R, X, G)}(M, N)$ set $T^*(f) = f \otimes R' : T^*(M) \rightarrow T^*(N)$.

Proposition 1.7. *Let $T = (\rho, \xi, \gamma)$ and T^* as above. The following assertions hold:*

- (a) T^* defines a left adjoint functor from $gr - (R, X, G)$ to $gr - (R', X', G')$.
- (b) The right adjoint functor of T^* is isomorphic to the functor $T_* : gr - (R', X', G') \rightarrow gr - (R, X, G)$ given as follows:

$$T_*(M') = \bigoplus_{x \in X} T_*(M')_x \quad (M' \in gr - (R', X', G'))$$

where the map $m' \mapsto m'^x$ is an isomorphism of additive groups from $M'_{\xi(x)}$ to $T_*(M')_x$ and the structure of right R -module on $T_*(M)$ is given by the following rule:

$$(m'^x)r = (m'\rho(r))^{xg} \quad (m' \in M', r \in R_g).$$

If $f \in \text{Hom}_{gr - (R', X', G')}(M', N')$ and $m' \in M'$ then $T_*(f)(m'^x) = f(m')^x$.

- (c) T_* is a left adjoint functor.

Proof: (a) Clearly $T^*(M) = \sum_{x' \in X'} T^*(M)$ and $T^*(M)_{x'} R'_g \subseteq T^*(M)_{x'g'}$, for every $x' \in X'$ and $g' \in G'$.

Let $P = \bigoplus_{x \in X} R'(\xi(x)) \in gr - (R', X', G')$. For any $x \in X$, let $u_x : R'(\xi(x)) \rightarrow P$ be the canonical monomorphism.

P becomes an $X \times X'$ graded R - R' -bimodule after the following rules:

$$\begin{aligned} ru_x(r') &= u_{gx}(rr') & (g \in G, r \in G_g, r' \in R', x \in X) \\ P_{(x,x')} &= R'(\xi(x))_{x'} & (x \in X, x' \in X'). \end{aligned}$$

Furthermore, for every $M \in gr - (R, X, G)$, the map

$$\begin{aligned} \phi_M : T^*(M) &\longrightarrow M \hat{\otimes}_R P \\ m \otimes r' &\mapsto \sum_{x \in X} m_x \otimes u_x(r') \end{aligned}$$

is an isomorphism of right R' -modules and for every $x' \in X'$, $\phi_M(T^*(M)_{x'}) = M \hat{\otimes}_R P_{x'}$. Therefore, $T^*(M) = \bigoplus_{x' \in X'} T^*(M)_{x'}$ and hence T^* defines a functor from $gr - (R, X, G)$ to $gr - (R', X', G')$. Moreover T^* is a left adjoint, because $T^* \cong - \hat{\otimes}_R P$.

(b) For every $M' \in gr - (R', X', G')$,

$$\begin{aligned} H(P_{R'}, M') &= \bigoplus_{x \in X} \text{Hom}_{gr - (R', X', G')}(xP, M') \\ &= \bigoplus_{x \in X} \text{Hom}_{gr - (R', X', G')}(R'(\xi(x)), M') \cong \bigoplus_{x \in X} M'_{\xi(x)}. \end{aligned}$$

Using this isomorphism it is straightforward to see that $H(P_{R'}, -) \cong T_*$ and (b) follows from Proposition 1.2.

(c) Let Q be the $X' \times X$ -graded R' - R -bimodule associated to T_* .

For any $M' \in gr - (R', X', G')$ we define

$$\begin{aligned} \Phi_{M'} : M' \hat{\otimes}_{R'} Q &\longrightarrow T_*(M') \\ m' \otimes q &\mapsto (m'q)^x \quad (m' \in M'_{x'}, q \in Q_{(x',x)}). \end{aligned}$$

Then $\Phi_{M'}$ is bijective and Φ defines an isomorphism of functors between $- \hat{\otimes}_{R'} Q$ and T_* . Therefore T_* is a left adjoint and $H(Q_{R'} -)$ is its right adjoint. ■

Lemma 1.8. *Let $T = (\rho, \xi, \gamma)$ as above and assume that $R' = R$ and ρ is the identity map. Consider X' as a right G -set by $x'g = x'\gamma(g)$ ($x' \in X', g \in G$). Then*

- (a) $gr - (R, X', G') = gr - (R, X', G)$.
 (b) $T^* = S^*$ and $T_* = S_*$ where $S = (1_R, \xi, 1_G)$.

Proof: (a) Clearly an $M \in gr - (R, X', G')$ is an object of $gr - (R, X', G)$ because for every $x' \in X'$, $M_{x'}R_g \subseteq M_{x'}R_{\gamma(g)} \subseteq M_{x'\gamma(g)} = M_{x'g}$.

On the other hand, let $M \in gr - (R, X', G)$. Let $g' \in G'$ and $0 \neq r \in R_{g'}$. If $g \in \text{Supp}_R(r)$, then $r_g = \rho(r_g) \in R_{\gamma(g)}$. Thus $r_g = (r_g)_{\gamma(g)}$. If $\gamma(g) \neq g'$, then

$$0 = r_{\gamma(g)} = \sum_{h \in \text{Supp}_R(r) \cap \gamma^{-1}(\gamma(g))} (r_h)_{\gamma(g)} = \sum_{h \in \text{Supp}_R(r) \cap \gamma^{-1}(\gamma(g))} r_h.$$

Therefore $r_g = 0$. Thus $\text{Supp}_R(r) \subseteq \gamma^{-1}(g')$. Let now $x' \in X'$. Then $M_{x'}r = M_{x'} \sum_{g \in \text{Supp}_R(r)} r_g \subseteq \sum_{g \in \text{Supp}_R(r)} M_{x'g} = \sum_{g \in \text{Supp}_R(r)} M_{x'\gamma(g)} = M_{x'g'}$.

(b) Is obvious. ■

Let R be a G -graded ring and $\xi : X \rightarrow X'$ a morphism of right G -sets. If $T = (1_R, \xi, 1_G)$, then the functors T^* and T_* are the functors T_ξ and S^ξ defined in [NRV]. In [NRV] the authors prove that if for every $x' \in X'$, $\xi^{-1}(x')$ is finite, then S^ξ is left adjoint of T_ξ . But the converse is also true as the following proposition shows.

Proposition 1.9. *Let R be a G -graded ring and $\xi : X \rightarrow X'$ a morphism of right G -sets. Let F be the right adjoint functor of S^ξ . F is a left adjoint functor if and only if for every $x' \in X'$, $\xi^{-1}(x')$ is finite. Moreover, in this case F is isomorphic to T_ξ .*

Proof: Let Q be the $X' \times X$ -graded R - R -bimodule associated to S^ξ . By Proposition 1.6, $S^\xi \cong - \hat{\otimes}_R Q$ and $F \cong H(Q_{R'} -)$.

Assume that $\xi^{-1}(x')$ is finite for every $x' \in X'$. For each $M \in gr - (R, X, G)$, let

$$\begin{aligned} \mu_M : H(Q_R, M) &\longrightarrow T_\xi(M) \\ \mu_M(f) &= \sum_{\xi(x)=x'} (f(1^x))_x \quad (f \in H(Q_{R'}M)_x). \end{aligned}$$

The claim follows by proving that $\mu : H(Q_R, -) \rightarrow T_\xi$ is a functorial isomorphism. This is left to the reader.

Conversely, if $H(Q_R, -)$ is a left adjoint functor, then it preserves direct sums. Let $x' \in X'$. Consider the functor $(-)_x' : gr - (R', X', G') \rightarrow \text{mod} - R'_c$ which associates to $M' \in gr - (R', X', G')$ the right R'_c -module $M'_{x'}$, and to $f \in \text{Hom}_{gr - (R', X', G')}(M', N')$ the map $f_{x'} : m' \mapsto f(m'_{x'})$. This functor preserves direct sums and hence $\text{Hom}_{gr - (R, X, G)}(x'Q, -) \cong (-)_{x'} \circ H(Q_R, -)$ preserves direct sums too. This implies that ${}_x'Q = S_\xi(R'(x'))$ is finitely generated (cf. [S]).

Let $\alpha = \sum_{\xi(x)=x'} (1^x)r_x = 0$ with $r_x \in R$. For every $y \in X$, $\alpha_y = \sum_{\xi(x)=x'} (1^x)(\sum_{xg=y} (r_x)g) = 0$. Therefore, we can assume, without loss of generality, that there exists $y \in Y$ such that for every $g \in \text{Supp}(r_x)$, $xg = y$. Thus $(\sum_{\xi(x)=x'} r_x)^y = \alpha_y = 0$ and hence $\sum_{\xi(x)=x'} r_x = 0$. But r_x has degree x in $(y)R$. We conclude that $r_x = 0$ for every $x \in X$. Therefore the sum $\sum_{x \in X} 1^x R$ is direct.

On the other hand, if $r \in R_g \cap Q_{(x',x)}$, then $x'g = \xi(x)$. Therefore $r^x = 1^x g^{-1} r$. We conclude that ${}_x'Q = \bigoplus_{\xi(x)=x'} 1^x R$. Since ${}_x'Q$ is finitely generated, the sum is finite, i.e. $\xi^{-1}(x')$ is finite. ■

2. Equivalences and Morita dualities

Propositions 1.2 and 1.6 set up the framework to give a description of all the equivalences and Morita dualities (in the sense of Colby-Fuller [CF]) between categories of modules graded by G -sets. This section is dedicated to present that description. Since the methods are similar to those given in [Rf2] and [MR] we shall skip the proofs.

Let \hat{R} be the $X \times X$ -graded R - R -bimodule associated to the identity functor in $gr - (R, G, X)$.

Let P be an $X \times X'$ -graded R - R' -bimodule and Q an $X' \times X''$ -graded R' - R'' -bimodule. Then $P \hat{\otimes}_{R'} Q$ is a subbimodule of $P \otimes_R Q$ and $(P \hat{\otimes}_{R'} Q)_{(x,x'')} = {}_x P \hat{\otimes}_R Q_{x''}$ ($x \in X, x'' \in X''$) defines an $X \times X''$ -grading on ${}_R P \hat{\otimes}_{R'} Q_{R''}$. Furthermore the functors $(- \hat{\otimes}_{R'} Q) \circ (- \hat{\otimes}_R P)$ and $- \hat{\otimes}_R (P \hat{\otimes}_{R'} Q)$ are isomorphic.

Proposition 2.1. *The following assertions are equivalent:*

- (a) *The categories $gr - (R, X, G)$ and $gr - (R', X', G')$ are equivalent.*
- (b) *There are an $X \times X'$ -graded R - R' -bimodule P and an $X' \times X$ -graded R' - R -bimodule Q such that $\hat{R} \cong P \hat{\otimes}_{R'} Q$ and $\hat{R} \cong Q \hat{\otimes}_R P$ as bigraded bimodules.*

Moreover, if P and Q are as in (b), then $-\hat{\otimes}_R P$ and $-\hat{\otimes}_{R'} Q$ are inverse equivalences of categories.

Corollary 2.2. (a) The categories $gr-(R, X, G)$ and $gr-(R', X', G')$ are equivalent if and only if the categories $(G, X, R) - gr$ and $(G', X', R') - gr$ are equivalent.

(b) Let P be an $X \times X'$ -graded R - R' -bimodule. $-\hat{\otimes}_R P : gr-(R, X, G) \rightarrow gr-(R', X', G')$ is an equivalence if and only if $P \hat{\otimes}_{R'} - : (G', X', R') - gr \rightarrow (G, X, R) - gr$ is an equivalence.

Definition. Let P be an $X \times X'$ -graded R - R' -bimodule and Q an $X \times X''$ -graded R - R'' -bimodule. We define

$$H({}_R P, {}_R Q) = H({}_R P, Q) \cap H(P, {}_R Q).$$

If we consider $\text{Hom}_R(P, Q)$ as an R' - R'' -bimodule by

$$(r' f r'')(p) = f(p r') r'' \quad (f \in \text{Hom}_R(P, Q), p \in P, r' \in R', r'' \in R'')$$

then $H_R(P, Q)$ is a subbimodule of ${}_R \text{Hom}_R(P, Q)_{R''}$, and it is an $X' \times X''$ -graded R' - R'' -bimodule with the following grading:

$$H({}_R P, {}_R Q)_{(x', x'')} = {}_{x'} H({}_R P, Q) \cap H(P, {}_R Q)_{x''} \quad (x' \in X', x'' \in X'').$$

Similarly, if P is an $X' \times X$ -graded R' - R -bimodule and Q is an $X'' \times X$ -graded R'' - R -bimodule, then $H(P_R, Q_R) = H(P_R, Q) \cap H(P, Q_R)$ has a natural structure of $X'' \times X'$ -graded R'' - R' -bimodule.

Let P be an $X \times X'$ -graded R - R' -bimodule. Consider the following map:

$$\begin{aligned} \chi^P : \hat{R}' &\longrightarrow H({}_R P, {}_R P) \\ \chi^P(r)(p) &= (p r')_{y'} \quad (r' \in R'(x')_{y'}, p \in P) \end{aligned}$$

χ^P is a homomorphism of $X' \times X'$ -graded $R' \times R'$ -bimodules.

Similarly there is an $X \times X$ -graded homomorphism of R - R -bimodules

$$\begin{aligned} \lambda^P : \hat{R} &\longrightarrow H(P_{R'}, P_{R'}) \\ \lambda^P(r)(p) &= {}_x (r p) \quad (r \in R(x), p \in P). \end{aligned}$$

Theorem 2.3. Let P be an $X \times X'$ -graded R - R' -bimodule. The following conditions are equivalent:

(a) The functor $-\hat{\otimes}_R P : gr-(R, X, G) \rightarrow gr-(R', X', G')$ is an equivalence of categories.

- (b) The functor $H(P_{R'}, -) : gr - (R', X', G''') \rightarrow gr - (R, X, G)$ is an equivalence of categories.
- (c) $P_{R'}$ is a projective generator in $gr - (R', X', G')$, $\lambda^P : \hat{R} \rightarrow H(P_{R'}, P_{R'})$ is an isomorphism and for every $x \in X$, ${}_x P_{R'}$ is finitely generated.
- (d) ${}_R P$ is a projective generator in $(G, X, R) - gr$, $\chi^P : \hat{R}' \rightarrow H({}_R P, {}_R P)$ is an isomorphism and, for every $x' \in X'$, ${}_R P_{x'}$ is finitely generated.

Let A be a ring and G a group. For any G -grading in A , the category $\text{mod-}A$ can be seen as a category of X -graded right A -modules where X is a singleton. Last theorem yields to the following two corollaries.

Corollary 2.4. *Let R be a G -graded ring, X a right G -set and A a ring. The following are equivalent.*

- (a) $gr - (R, X, G)$ is equivalent to $\text{mod-}A$.
- (b) $(G, X, R) - gr$ is equivalent to $A\text{-mod}$.
- (c) There exists a finitely generated projective generator P in $gr - (R, X, G)$ such that A is isomorphic to $\text{End}_{gr - (R, X, G)}(P)$.
- (d) There exists a finite subset F of X such that $U = \bigoplus_{x \in F} R(x)$ is a generator of $gr - (R, X, G)$ and $\text{End}_{gr - (R, X, G)}(U)$ is Morita equivalent to A .

Corollary 2.5. *Let R be a G -graded ring and X a right G -set. The following are equivalent.*

- (a) There exists a ring A such that $gr - (R, X, G)$ is equivalent to $\text{mod-}A$.
- (b) There exists a ring A such that $(G, X, R) - gr$ is equivalent to $A\text{-mod}$.
- (c) $gr - (R, X, G)$ has a finitely generated (projective) generator.
- (d) There exists a finite subset F of X such that $\bigoplus_{x \in F} R(x)$ is a generator of $gr - (R, X, G)$.

Example 2.6. Consider $T = (\rho, \xi, \gamma)$ as in Section 1. The functor T^* is isomorphic to $-\hat{\otimes}_R P$ where $P = \bigoplus_{x \in X} R'(\xi(x))$ (see the proof of Proposition 1.7). Since for every $x \in X$, ${}_x P_{R'} = R'(\xi(x))$ is finitely generated and projective, T^* is an equivalence if and only if $P_{R'}$ generates $gr - (R', X', G')$ and $\lambda^P : \hat{R} \rightarrow H(P_{R'}, P_{R'})$ is an isomorphism. We claim that $T^* \cong -\hat{\otimes}_R R' : gr - (R, X, G) \rightarrow gr - (R', X', G')$ is an equivalence if and only if the following conditions hold

- (a) ρ is an isomorphism between R and $\bigoplus_{\text{Im}(\xi)g' \cap \text{Im}(\xi) \neq \emptyset} R'_{g'}$.

- (b) For every $x' \in X'$ there exists $r'_1 \in R'_{g'_1}, r'_2 \in R'_{g'_2}, \dots, r'_n \in R'_{g'_n}; s'_1 \in R'_{g'_1-1}, s'_2 \in R'_{g'_2-1}, \dots, s'_n \in R'_{g'_n-1}$, such that $x'g'_i \in \text{Im}(\xi)$ for every $i = 1, 2, \dots, n$ and $\sum_{i=1}^n r'_i s'_i = 1$.

Proof: If $r' \in R'$ and $x \in X$, then r'^x will denote r' considered in $R'(\xi(x))$.

Assume T^* is an equivalence. Then $\lambda^P : \hat{R} \rightarrow H(P_{R'}, P_{R'})$ is an isomorphism and $P_{R'}$ is a generator. Let $r \in \text{Ker}(\rho)$. For every $p \in P$, $[\lambda^P(r^x)](p) = {}_x(rp) = {}_x(\rho(r)p) = 0$. Thus $r^x \in \text{Ker}(\lambda^P) = 0$. Therefore ρ is injective. If $r \in R_g$ and $x \in X$, then $\rho(r) \in R_{\gamma(g)}$ and $\xi(x)\gamma(g) = \xi(xg)$. Therefore $\rho(R) \subseteq \bigoplus_{\text{Im}(\xi)g' \cap \text{Im}(\xi) \neq \emptyset} R'_{g'}$. On the other hand, let $r' \in R'_{g'}$ and $x, y \in X$ such that $\xi(x)g' = \xi(y)$. Consider the following homomorphism of right R' -modules

$$L : P \longrightarrow P$$

$$L(s'^z) = \begin{cases} (r's')^x & \text{if } z = y \\ 0 & \text{if } z \neq x \end{cases} \quad (s' \in R', z \in X).$$

Plainly $L \in H(P_{R'}, P_{R'})(x, y)$, thus there exists $r \in R(x)_y$ such that $L = \lambda^P(r^x)$. But $r'^x = L(1^y) = \lambda^P(r)(1^y) = {}_x(\rho(r)^x) = \rho(r)^x$. This proves (a).

Let $x' \in X'$. Since $P_{R'} = \bigoplus_{x \in X} R'(\xi(x))$ is a generator of $gr - (R', X', G')$, then $\{R'(\xi(x)) | x \in X\}$ is a set of generators of $gr - (R', X', G')$. Therefore there exist $x_1, x_2, \dots, x_n \in X$ and an X' -graded epimorphism $\bigoplus_{i=1}^n R'(\xi(x_i)) \rightarrow R'(x')$. Thus there exist $r'_1 \in R'(x')_{\xi(x_1)}, r'_2 \in R'(x')_{\xi(x_2)}, \dots, r'_n \in R'(x')_{\xi(x_n)}$ and $s'_1 \in R'(\xi(x_1))_{x'}, s'_2 \in R'(\xi(x_2))_{x'}, \dots, s'_n \in R'(\xi(x_n))_{x'}$ such that $\sum_{i=1}^n r'_i s'_i = 1$. If we replace the s'_i 's and the r'_i 's by its homogeneous components we can assume they are homogeneous. Moreover, if we take the homogeneous component of degree 1 we may assume that r'_1 and s'_1 have inverse degrees and if r'_1 has degree g'_1 , then $x'g'_1 = \xi(x_1)$. This proves (b).

Conversely, assume that (a) and (b) hold. Since λ^P is an $X \times X$ -graded homomorphism, to prove that it is injective it is enough to prove that if $0 \neq r \in R(x)_y = P_{(x,y)}$, then $\lambda^P(r^x) \neq 0$. But $xg = y$ for every $g \in \text{Supp}(r)$. Therefore $r1^y = \sum_{g \in \text{Supp}(r)} r^{gy} = r^x$. Thus $[\lambda^P(r^x)](1^y) = {}_x(r1^y) = {}_x(r^x) \neq 0$. Because λ^P is $X \times X$ -graded, to prove that it is surjective it is enough to see that $H(P_{R'}, P_{R'})(x, y) \subseteq \text{Im}(\lambda^P)$ for every

$(x, y) \in X \times X$. Let $f \in H(P_{R'}, P_{R'})(x, y)$ and $r' = f(1^y) \in R'(\xi(x))_{\xi(y)}$. For every $g' \in \text{Supp}(r')$, $\xi(x)g' = \xi(y)$. Therefore, by (a) there exists $r_{g'} \in R(x)_y$ such that $\rho(r_{g'}) = r'_{g'}$. Then it is easy to see that $f = \lambda^P(r)$ where $r = \sum_{g' \in \text{Supp}(r')} r_{g'}$.

Let $x' \in X'$. Let r'_i, s'_i and $g'_i (i = 1, 2, \dots, n)$ as in (b). Let $L : \bigoplus_{i=1}^n R'(x'_i g'_i) \rightarrow R'(x')$ given by $L(a'_1, a'_2, \dots, a'_n) = \sum_{i=1}^n r'_i a'_i$. L is an homomorphism of X' -graded modules and $1 = L(r'_1, r'_2, \dots, r'_n)$. But $x'_i g'_i \in \text{Im}(\xi)$. Therefore $\{R'(\xi(x)) | x \in X\}$ is a set of generators of $gr - (R', X', G')$. Therefore $P_{R'}$ is a generator of $gr - (R', X', G')$. This finish the proof. ■

Example 2.7. As a particular case of Example 2.6, let R be a G -graded ring, X a right G -set, $x \in X$ and G_x the stabilizer of x in G . Consider $Y = \{x\}$ as a right G_x -set. Set $T = (\rho, \xi, \gamma)$ where $\rho : R_{G_x} \rightarrow R$, $\xi : Y \rightarrow X$ and $\gamma : H \rightarrow G$ are inclusion maps.

In this case condition (a) in the above example always holds. On the other hand $gr - (R_H, Y, H) = \text{mod-}R_H$ and the functor T_* is isomorphic to the functor $(-)_x$ which associates $M \in gr - (R, X, G)$ to M_x and acts on morphisms by restricting on the domains. In this situation the result proved in last example translates into:

The following conditions are equivalent:

- (a) $- \otimes_{R_{G_x}} R : \text{mod-}R_{G_x} \rightarrow gr - (R, X, G)$ is an equivalence.
- (b) $(-)_x$ is an equivalence.
- (c) $R(x)$ is a generator in $gr - (R, X, G)$.
- (d) For every $y \in Y$, $R_{x^{-1}x} = R_{x^{-1}y} \cdot R_{y^{-1}x}$ (where $R_{x^{-1}y} = \bigoplus_{xg=y} R_g$).

Notice that condition (d) implies that G acts transitively on X .

When $X = G$ with regular right action, the above equivalent conditions do not depend on x . Namely, if $R(g)$ is a generator in $gr - R$, then for every $x \in G$, $R(x)$ is a generator too. Graded rings satisfying these condition are the strongly graded rings. When X is not G the situation is not so symmetric, as the following example shows.

Example 2.8. Let A be a ring and B a proper overring such that the inclusion map $j : A \rightarrow B$ splits as a morphism of $A - A$ -bimodules, i.e. there is an $A - A$ -bimodule homomorphism $\varepsilon : A \rightarrow A$ such that $\varepsilon \circ j = 1_A$.

Let G be the permutation group on three elements. Let $a \in G$ be a transposition and $b \in G$ a 3-cycle. Let K be the subgroup of G generated by b and H the subgroup of G generated by a .

Let $M = A \times B \times B$ considered as an A - A -bimodule. If we define

$$\left. \begin{aligned} b(x, y, z) &= (\varepsilon(z), \varepsilon(x), \varepsilon(y)) \\ (x, y, z)b &= (\varepsilon(y), \varepsilon(z), \varepsilon(x)) \end{aligned} \right\} (x, y, z) \in M$$

then M becomes an $A[K] \times A[K]$ -bimodule. Consider $R = A[K] \ltimes M$ the trivial extension of $A[K]$ by M . The elements of R are pairs (α, m) with $\alpha \in A[K]$ and $m \in M$ with the ring structure given by

$$\left. \begin{aligned} (\alpha, m) + (\beta, n) &= (\alpha + \beta, m + n) \\ (\alpha, m)(\beta, n) &= (\alpha\beta, \alpha n + m\beta) \end{aligned} \right\} \alpha, \beta \in A[K], m, n \in M.$$

Consider in R the following G -grading:

$$\begin{array}{lll} R_e = A \times 0 & R_b = Ab \times 0 & R_{b^2} = Ab^2 \times 0 \\ R_a = 0 \times (A \times 0 \times 0) & R_{ab} = 0 \times (0 \times B \times 0) & R_{ab^2} = 0 \times (0 \times 0 \times B). \end{array}$$

Let X be the set of right H -cosets and consider G acting on X by right translation. Let $x = H$ and $y = Hb$. Then

$$\begin{aligned} R_{x^{-1}x} &= R_H = R_{Hb} \cdot R_{b^{-1}H} = R_{x^{-1}y} \cdot R_{y^{-1}x} \\ R_{y^{-1}x} \cdot R_{x^{-1}y} &= R_{b^{-1}H} \cdot R_{Hb} = [Ab^2 \times (0 \times B \times 0)][Ab \times (0 \times B \times 0)] = \\ &= A \times (0 \times 0 \times A) \neq A \times (0 \times 0 \times B) = R_{b^{-1}Hb} = R_{y^{-1}y}. \end{aligned}$$

Therefore, by Example 2.7, $R(x)$ is a generator in $gr-(R, X, G)$ while $R(y)$ is not.

In [CF] the authors extend the concept of Morita duality to Grothendieck categories. Let $(F : \mathcal{C} \rightarrow \mathcal{D}, G : \mathcal{D} \rightarrow \mathcal{C})$ be a pair of adjoint functors on the right. Let us denote by $\sigma : 1_{\mathcal{C}} \rightarrow GF$ and $\tau : 1_{\mathcal{D}} \rightarrow FG$ the units of the adjunction (they can be interpreted as unit and counit of a pair of covariant functors). Let

$$\begin{aligned} \mathcal{R}_{\mathcal{C}} &= \{C \in \mathcal{C} \mid \sigma_C \text{ is an isomorphism}\} \\ \mathcal{R}_{\mathcal{D}} &= \{D \in \mathcal{D} \mid \tau_D \text{ is an isomorphism}\}. \end{aligned}$$

Then F and G induce a duality between $\mathcal{R}_{\mathcal{C}}$ and $\mathcal{R}_{\mathcal{D}}$.

We say that the pair (F, G) is a Morita Duality if F and G are exact and the subcategories $\mathcal{R}_{\mathcal{C}}$ and $\mathcal{R}_{\mathcal{D}}$ contain a set of generators of \mathcal{C} and \mathcal{D} , respectively, and both are closed under subobjects and quotient objects.

In [AW] the authors define an alternative version of Morita duality for Grothendieck categories which is not equivalent to the Colby-Fuller's

one in general but it is so in the case of categories of modules graded by G -sets (see [GG]).

Note that if (F, G) is a Morita duality, then \mathcal{R}_C and \mathcal{R}_D are closed under finite coproducts, therefore \mathcal{R}_C and \mathcal{R}_D contain any finitely generated object of C and D , respectively.

Let P be a $X \times X'$ -graded R - R' -bimodule. Let $u : 1_{gr-(R, X, G)} \rightarrow H(P_{R'}, -) \circ (- \hat{\otimes}_R P)$ and $c : (- \hat{\otimes}_R P) \circ H(P_{R'}, -) \rightarrow 1_{gr-(R', X', G')}$ the unit and the counit of the adjunction pair $(H(P_{R'}, -), (- \hat{\otimes}_R P))$ and $\sigma : 1_{(G, X, R)-gr} \rightarrow H(-, P_{R'}) \circ H(-, {}_R P)$ and $\tau : 1_{gr-(R', X', G')} \rightarrow H(-, {}_R P) \circ H(-, P_{R'})$ the unit and counit of the adjunction pair $(H(-, {}_R P), H(-, P_{R'}))$.

An object $M \in (G, X, R) - gr$ (resp. $N \in gr - (R', X', G')$) is said to be P -reflexive if σ_M (resp. τ_N) is an isomorphism.

Lemma 2.9. *Let P an $X \times X'$ -graded R - R' -bimodule. The following are equivalent:*

- (a) $\lambda^P : \hat{R} \rightarrow H(P_{R'}, P_{R'})$ (resp. $\chi^P : \hat{R}' \rightarrow H({}_R P, {}_R P)$ is an) is an isomorphism.
- (b) For every $x \in X$ (resp. $x' \in X'$), $u_{R(x)}^P$ (resp. $c_{R'(x')}^P$) is an isomorphism.
- (c) For every $x \in X$ (resp. $x' \in X'$), $(x)R$ (resp. $R'(x')$) is P -reflexive.

P is said to be faithfully balanced if both λ^P and χ^P are isomorphisms.

Theorem 2.10. *Let P be an $X \times X'$ -graded R - R' -bimodule. The following conditions are equivalent:*

- (a) $(H(-, {}_R P), H(-, P_{R'}))$ define a Morita duality between $(G, X, R) - gr$ and $gr - (R', X', G')$.
- (b) For every $x \in X$ and $x' \in X'$, every graded quotient of $(x)R, R'(x'), {}_x P$ and $P_{x'}$ is P -reflexive.
- (c) P is graded faithfully balanced, $\{P_{x'} | x' \in X'\}$ is a set of injective cogenerators in the category $(G, X, R) - gr$ and $\{{}_x P | x \in X\}$ is a set of injective cogenerators in the category $gr - (R', X', G')$.

3. Duality Theorems

The main goal of this section is to extend the Duality Theorems of Cohen and Montgomery [CM] to our setting. There are other alternative extensions which can be found in [Q], [B1], [B2] and [AN] (see also [A]).

The originality of our methods is that they are purely categorical. We give some equivalences of categories which will be useful in applications.

Let R be a G -graded ring and X a left G -set. X will be considered as a right G -set by $xg = g^{-1}x$ ($x \in X, g \in G$).

Let $P = \bigoplus_{x \in X} R(x)$. For every $x \in X$ let $u_x : R(x) \rightarrow P$ be the canonical monomorphism and $\pi_x : P \rightarrow R(x)$ the canonical projection. P becomes an $X \times X$ -graded R - R -bimodule by

$$ru_x(a)s = u_{gx}(ras) \quad (r \in R_g, a, s \in R)$$

and the $X \times X$ -grading

$$P_{(x,y)} = u_x(R(x)_y) \quad (x, y \in X).$$

Moreover ${}_R P$ and P_R are faithful modules.

Note that P is the $X \times X$ -graded R - R -bimodule associated to the identity functor in $gr - (R, X, G)$. (See Definition 1.4).

Given $x, y \in X$ we write $R_{xy^{-1}} = \sum_{x=gy} R_g$. Also if $r \in R$, then $r_{xy^{-1}}$ will denote $\sum_{x=gy} r_g$. Note that $R_{xy^{-1}}$ (resp. $r_{xy^{-1}}$) is both the component of degree x of $(y)R$ (resp. of r considered in $(y)R$) and the component of degree y of $R(x)$ (resp. of r in $R(x)$).

Remark. Let $FCM_X(R)$ be the ring of all finite column square X -matrices with entries in R . Since ${}_R P$ is free and $\{u_x(1) | x \in X\}$ is a basis of ${}_R P$, then the map

$$\begin{aligned} \Phi : \text{End}({}_R P) &\longrightarrow FCM_X(R) \\ f &\longmapsto M_f \end{aligned}$$

(being M_f the square X -matrix having $\pi_y f u_x(1)$ at the (x, y) -th entry) is a ring isomorphism. Moreover, it is quite easy to see that $\Phi(\text{End}_{gr-(R, X, G)}(P)) = \{a \in FCM_X(R) | \text{for every } x, y \in X, \text{ the } (x, y)\text{-th entry belongs to } R_{xy^{-1}}\}$. Thus Φ gives a matricial description of $(\text{End}_{gr-(R, X, G)}(P))$. This matricial description has been obtained by T. Albu [A] for the particular case $X = G$.

Let $S = \text{End}_{gr-(R, X, G)}(P)$. If we consider S trivially graded by $G' = \{e\}$ and $X' = G'$ as a left G' -set, then P is naturally an $X' \times X$ -graded S - R -bimodule.

Since P is an $X \times X$ -graded R - R -bimodule, then the map $r \mapsto \Lambda_r$ which associates to an element $r \in R$ the left multiplication-by- r map in P is a ring homomorphism $\Lambda : R \rightarrow S$. Since ${}_R P$ is faithful, Λ is injective. Note that for every $r \in R_{xy^{-1}}$, $\pi_y \Lambda_r u_x = \lambda_r : R(x) \rightarrow R(y)$.

Let $I = H({}_R P, P)$. Clearly $I = \{\alpha \in S | \alpha u_x = 0 \text{ for almost all } x \in X\}$.

Lemma 3.1. *I is an S-R-bisubmodule of S and it is isomorphic to sP_R .*

Proof: Let $\varphi : P \rightarrow S$ be the additive homomorphism given by

$$\varphi(u_x(r))u_y = u_x\lambda_{r_{xy^{-1}}}, \quad (r \in R, x, y \in X).$$

Note that for every $x, y \in X, u_x\lambda_{r_{xy^{-1}}} = \Lambda_{r_{xy^{-1}}}u_y$.

Let $\alpha \in S, r \in R$ and $x, y, z \in X$. Then $\pi_y\varphi(\alpha u_x(r))u_z = \pi_y(\sum_{w \in X} \varphi u_w \pi_w \alpha u_x(r))u_z = \pi_y\varphi(u_y \pi_y \alpha u_x(r))u_z = \lambda_{(\pi_y \alpha u_x(r))_{yz^{-1}}}$. If $g \in G$, then $r_g \in R(x)_{xg}$, therefore $\pi_y \alpha u_x(r_g) \in R(y)_{xg}$. Thus $(\pi_y \alpha u_x(r))_{yz^{-1}} = \sum_{xg=z} \pi_y \alpha u_x(r_g) = \pi_y \alpha u_x(r_{xz^{-1}})$. Then $\pi_y\varphi(\alpha u_x(r))u_z = \lambda_{\pi_y \alpha u_x(r_{xz^{-1}})} = \pi_y \alpha u_x \lambda_{r_{xz^{-1}}} = \pi_y \alpha \varphi(u_x(r))u_z$. We conclude that for every $x \in X, r \in R$ and $\alpha \in S, \varphi(\alpha u_x(r)) = \alpha \varphi(u_x(r))$ and hence φ is S-linear.

Let now $r, s \in R, x, y \in X$. Then $\varphi(u_x(r)s)u_y = \varphi(u_x(rs))u_y = u_x\lambda_{(rs)_{xy^{-1}}}$. But $(rs)_{xy^{-1}} = \sum_{w \in X} r_{xw^{-1}}s_{wx^{-1}}$. Thus $\varphi(u_x(r)s)u_y = \sum_{w \in X} u_x\lambda_{r_{xw^{-1}}} \lambda_{s_{wy^{-1}}} = \sum_{w \in X} \varphi(u_x(r))u_w \lambda_{s_{wy^{-1}}} = \varphi(u_x(r)) \sum_{w \in X} u_w \lambda_{s_{wy^{-1}}} = \varphi(u_x(r)) \sum_{w \in X} \Lambda_{s_{wy^{-1}}} u_y = \varphi(u_x(r))\Lambda_s u_y$. Therefore $\varphi(u_x(r)s) = \varphi(u_x(r))s$, that is, φ is an homomorphism of right R-modules.

Let $p = \sum_{x \in X} u_x(r_x)$ be an element of $\text{Ker}(\varphi)$. Then for all $x, y \in X, 0 = \pi_x\varphi(p)u_y = u_x\lambda_{(r_x)_{xy^{-1}}}$. Thus $r_x = \sum_{y \in X} (r_x)_{xy^{-1}} = 0$. We conclude that φ is injective.

Let $r \in R$ and $x \in X$. Then for every $y \in X, \varphi(u_x(r))u_y = 0$ unless $x \in \text{Supp}(r)y$. Therefore $\varphi(P) \subseteq I$.

Finally, let $\alpha \in I$. For every $x \in X$ let $r_x = \sum_{y \in X} \pi_x \alpha u_y(1)$. Since $u_y(1) \in R(y)_y, \pi_x \alpha u_y(1) \in R(x)_y$ and hence $(r_x)_{xy^{-1}} = \pi_x \alpha u_y(1)$. Let $p = \sum_{x \in X} u_x(r_x)$. Then, for every $y \in Y, \varphi(p)u_y(r') = \sum_{x \in X} u_x((r_x)_{xy^{-1}}r') = \sum_{x \in X} u_x(\pi_x \alpha u_y(1)r') = \alpha u_y(r')$. Thus $\varphi(p) = \alpha$. We conclude that $\varphi(P) = I$. ■

For every subset F of X let $p_F \in S$ given by $p_F(m) = \sum_{x \in F} u_x \pi_x(m)$.

Plainly $p_F p_{F'} = p_{F \cap F'}$, for every two subset F and F' of X . Therefore, p_F is an idempotent for every subset F of X . Moreover $p_F + p_{X-F} = 1$ for every $F \subseteq X$.

For every $x \in X$ let $p_x = p_{\{x\}} (= u_x \pi_x)$. $\{p_x | x \in X\}$ is a set of orthogonal idempotents of S . For every subset F of X , $P_F \in I$ if and only if F is finite.

Lemma 3.2. (a) Let $\alpha \in S$, $\alpha \in I$ if and only if there exists a finite subset F of X such that $\alpha = \alpha p_F$.

(b) I is idempotent.

(c) $sI = \bigoplus_{x \in X} Sp_x$ and $Sp_x = Rp_x$ for every $x \in X$.

Proof. Let $\alpha \in I$. There exists a finite subset F of X such that $\alpha u_x = 0$ for all $x \in X - F$. Then $\alpha = \alpha p_F$. This proves (a) and (b).

The sum $\sum_{x \in X} Sp_x$ is direct because $\{p_x | x \in X\}$ is a set of orthogonal idempotents. Moreover, for every $x \in X$ and $\alpha \in S$, $\alpha p_x = \Lambda_r p_x$ where $r = \sum_{y \in X} \pi_y \alpha u_x(1)$. Indeed, let $m \in P$, then $\alpha p_x(m) = \alpha u_x \pi_x(m) = \sum_{y \in X} u_y \pi_y(\alpha u_x(1)) \pi_x(m) = \sum_{y \in X} u_y (r_{y x^{-1}}) \pi_x(m) = \sum_{y \in X} u_y \lambda_{r_{y x^{-1}}}(\pi_x(m)) = \sum_{y \in X} \Lambda_{r_{y x^{-1}}} u_x \pi_x(m) = \Lambda_r p_x(m)$. ■

Given a left A -module M , then $\text{Gen}(A M)$ denotes the full subcategory of $A\text{-mod}$ formed by the right A -modules generated by $A M$. Next Lemma is obvious.

Lemma 3.3. Let I be an idempotent left ideal of a ring A . Then $\text{Gen}(A I) = \{M \in A\text{-mod} | IM = M\}$.

Theorem 3.4. The pair of adjoint functors $(G, X, R) - gr \begin{matrix} P \hat{\otimes}_R \\ \mathbb{E} \\ H(S P, -) \end{matrix} S\text{-mod}$, defines an equivalence of categories between $(G, X, R) - gr$ and $\text{Gen}(S I)$.

Proof. Let $T = P \hat{\otimes}_R -$ and $H = H(S P, -)$. Let u be the unit of the adjunction pair (T, H) and c the counit. Let us remark that u and c are given by:

$$\begin{aligned} u_M : M &\longrightarrow HT(M) && (M \in (G, X, R) - gr) \\ [u_M(m)](p) &= \sum_{x \in X} m_x \otimes_x p && (m \in M, p \in P) \\ c_N : TH(N) &\longrightarrow N && (N \in \text{mod-}S) \\ c_N(f \otimes p) &= f(p) && (x \in X, p \in {}_x P, f \in H(P_S, N)_x) \end{aligned}$$

For every $M \in (G, X, R) - gr$, u_M is nothing but the composition of the following canonical isomorphisms: $M \cong \bigoplus_{x \in X} P(x) \hat{\otimes}_R M$ (see Lemma

1.1) = $IP \hat{\otimes}_R M \cong \text{Hom}({}_S I, P \hat{\otimes}_R M) = HT(M)$. Therefore u_M is an isomorphism.

Let now $N \in S\text{-mod}$. Let $\varphi_N : TH(N) \rightarrow IN$ be the composition of the following canonical isomorphisms: $\varphi_N : P \hat{\otimes}_R \text{Hom}({}_S I, N) \cong \bigoplus_{x \in X} R(x) \hat{\otimes}_R \text{Hom}({}_S I, N) \cong \text{Hom}({}_S I, N) \cong IN$. Then the following diagram is commutative:

$$\begin{array}{ccc} TH(N) & \xrightarrow{1} & TH(N) \\ \varphi_N \uparrow & & \uparrow c_N \\ IN & \xrightarrow{j} & N \end{array}$$

where j is the inclusion map.

Thus c_N is an isomorphism if and only if $N = IN$. Now Lemma 3.3 applies. ■

Remark. Theorem 3.4 implies [CM, Theorem 2.2] and [NRV, Theorem 2.13].

Recall that a ring A (not necessarily with unit) is said to have local units if for every finite subset F of A there exists an idempotent e of A such that $F \subseteq eAe$. We will say that a left ideal I of S has local units if it has local units when it is considered as a ring without unit.

Lemma 3.5. *Let A be a ring and I a left ideal of A which has local units. Let B a subring of A which contains I . The restriction of scalar functor induces an equivalence between $\text{Gen}_{(A)I}$ and $\text{Gen}_{(B)I}$.*

Proof: Let $F : A\text{-mod} \rightarrow B\text{-mod}$ be the restriction of scalars functor. F is obviously faithful. Let $M, N \in \text{Gen}_{(A)I}$ and $f \in \text{Hom}_B(M, N)$. Let $m \in M$ and $a \in A$. Let e be an idempotent in I such that $em = m$. Then $f(am) = f(aem) = aef(em) = af(m)$. Therefore $f \in \text{Hom}_A(M, N)$ and hence F is full.

Let now $M \in \text{Gen}_{(B)I}$. For every $m \in M$ and $a \in A$ let $am = (ae)m$ where e is an idempotent in I such that $m = em$. This definition does not depend on the choice of e because if $em = m = e'm$ there exists e'' idempotent in I such that $e, e' \in e''Ae''$. Thus $(ae)m = (ae'')em = (ae'')m = (ae'')e'm = (ae')m$. This gives a structure of left A -module on M and $F(M) = M$. ■

Corollary 3.6. *For every subring A of S which contains I , $P \hat{\otimes}_R - : (G, X, R) - gr \rightarrow A\text{-mod}$ defines an equivalence of categories between $(G, X, R) - gr$ and $\text{Gen}(AI)$.*

Remark. Last corollary generalizes [A, Theorem 2.5].

Let G, X, R, P and S as above. Let H be a group and assume that X has an structure of G - H -set. Let $\overline{\text{END}}(P_R)_H = \{\alpha \in \text{End}(P_R) | \alpha(P_x) \subseteq P_{xH} \text{ for every } x \in X\}$. $\overline{\text{END}}(P_R)$ is a subring of $\text{End}(P_R)$.

For every $h \in H$ let $\text{END}(P_R)_h = \{\alpha \in \text{End}(P_R) | \alpha(P_x) \subseteq P_{xh^{-1}} \text{ for every } x \in X\}$. For every $h, h' \in H$, $\text{END}(P_R)_h \cdot \text{END}(P_R)_{h'} \subseteq \text{END}(P_R)_{hh'}$. Therefore $\text{END}(P_R)_H = \sum_{h \in H} \text{END}(P_R)_h$ is a subring of $\overline{\text{END}}(P_R)_H$.

Moreover for every $h \in H$ the homomorphism $\bar{h} : P \rightarrow P$ given by $\bar{h}u_x = u_{xh^{-1}}(x \in X)$ belongs to $\text{END}(P_R)_h$ and $\bar{h}\bar{h}' = \bar{h}h'$ for every $h, h' \in H$. Therefore $\text{END}(P_R)_H$ is an H -Clifford system.

Lemma 3.7. *The sum $\sum_{h \in H} \text{END}(P_R)_h$ is direct if and only if H acts freely on X . In this case $\text{END}(P_R)$ is a skew group ring over H with underground ring S .*

Proof: Let $h \in H$ and $0 \neq \alpha \in \text{END}(P_R)_h \cap \sum_{h' \neq h} \text{END}(P_R)_{h'}$. Let $x \in X$ such that $\alpha u_x \neq 0$. Then $\alpha u_x(1) \neq 0$. Let $y \in X$ such that $\pi_y \alpha u_x(1) \neq 0$. Since $u_x(1) \in P_x$, then $\pi_y \alpha u_x(1) \in R(y)_{xh^{-1}}$. Let $g \in \text{Supp}(\pi_y \alpha u_x(1))$. Then $xh^{-1} = g^{-1}y$. Let $h_1, h_2, \dots, h_n \in H - \{h\}$, and $\alpha_1 \in \text{END}(P_R)_{h_1}, \alpha_2 \in \text{END}(P_R)_{h_2}, \dots, \alpha_n \in \text{END}(P_R)_{h_n}$, such that $\alpha = \sum_{i=1}^n \alpha_i$. Let $i = 1, 2, \dots, n$ such that $\alpha_i u_x(1) \neq 0$. By repeating the same argument, one has that $xh_i^{-1} = g^{-1}y = xh^{-1}$. Thus X is not a free left H -set.

Conversely, assume that the sum $\sum_{h \in H} \text{END}(P_R)_h$ is direct. Let $x \in X$ and $h \in H$ such that $xh = x$. Then $0 \neq P_x \in \text{END}(P_R)_e \cap \text{END}(P_R)_{h^{-1}}$ and hence $h = e$. ■

Let X be an G - H -set. Consider the following relation of equivalence on X :

$$x \sim y \iff y \in xH.$$

Then $X/H = \{xH | x \in X\}$ is the set of equivalent classes of X by the relation of equivalence \sim . G acts on X/H by $g(xH) = gxH$. The map $\xi : x \mapsto xH$ is a homomorphism of right G -set.

Let X be a set and I an ideal. By $FM_X(I)$ we will denote the set of all square X -matrices over I having almost all entries equal to zero.

Theorem 3.8. *Let H and G be two groups, X an G - H -set and R a G -graded ring. Let $P = \bigoplus_{x \in G} R(x)$ and $U = \bigoplus_{C \in X/H} R(C)$. Let $A = \overline{\text{END}}(P_R)_H$, $S = \text{End}_{gr-(R, X/H, G)}(U)$, $J = \{\alpha \in A \mid \alpha(R(x)) = 0 \text{ for almost all } x \in X\}$ and $I = \{\alpha \in S \mid \alpha(R(C)) = 0 \text{ for almost all } C \in X/H\}$.*

There exists an isomorphism $\Phi : A \rightarrow FCM_H(S)$ such that $\Phi(J) = FM_H(I)$.

Proof: Let $\xi : X \rightarrow X/H$ be the map which associates to $x \in X$ the class xH . Set $Q = T_\xi(P)$.

Plainly $Q = \bigoplus_{x \in X} R(xH) = U^{(H)}$. Therefore there is a canonical isomorphism $\Phi : \text{End}_{(G, X/H, R)-gr}(Q) \rightarrow FCM_H(\text{End}_{(G, X/H, R)-gr}(U))$ such that $\Phi(K) = FM(I)$ being $K = \{\alpha \in \text{End}_{(G, X/H, R)-gr}(Q) \mid u_x \alpha = 0 \text{ for almost all } x \in X\}$. The theorem follows by realizing that $\text{End}_{(G, X/H, R)-gr}(Q) = \overline{\text{END}}(P)_H$ and hence $J = K$. ■

Remark. Last theorem implies [CM, Theorem 3.5], [B2, Theorem 2.2] and [NRV, Corollary 2.2].

Lemma 3.9. *Let A be a ring and X a set. Let $B = M_X(A)$ be the finite row and column matrix ring over R indexed by X . Let $x \in X$ and P_x the $X \times X$ matrix having 1 at the (x, x) -th entry and zeros elsewhere. The functor $\text{Hom}({}_B Bp_x, -)$ defines an equivalence of categories between $\text{Gen}({}_B Bp_x)$ and $A\text{-mod}$.*

Proof: Let $P = Bp_x$. ${}_B P$ is finitely generated and projective and $\text{End}({}_B P)$ is isomorphic to A . It only remains to prove that ${}_B P$ generates each of its submodules and apply [F, Theorem 2.6 and Lemma 2.2].

Let M be a submodule of ${}_B P$. For any $y \in X$ let $M_y = \{\alpha(y, x) \mid \alpha \in M\}$. If $y, z \in X$ and $\alpha \in M$, then $(e_{z,y} \alpha)(z, x) = \alpha(y, x)$, $e_{z,y}$ being the $X \times X$ -matrix having 1 at the (z, y) -entry and zeros elsewhere. Therefore, $M_y = M_z$, that is, $N = M_y$ does not depend on $y \in X$.

Let $M' = \{\alpha \in P \mid \alpha(y, x) \in N \text{ for all } y \in X\}$. We claim that $M = M'$. The inclusion $M \subseteq M'$ is clear. Let $\alpha \in M'$. For every $y \in X$ there exists $\beta_y \in M$ such that $\alpha(y, x) = \beta_y(y, x)$. Then $\alpha = \sum_{\alpha(y,x) \neq 0} p_y \beta_y \in M$.

For every $a \in N$ let \hat{a} be the $X \times X$ matrix having a at the (x, x) -th entry and zeros elsewhere. Let $f_a : P \rightarrow M$ be the right multiplication

by \hat{a} map. Then, for every $\alpha \in M$, $\alpha = \sum_{\alpha(y,x) \neq 0} f_{\alpha(y,x)}(e_{(y,x)})$. Therefore M is generated by ${}_B P$. ■

Corollary 3.10. *With the same hypothesis and notation than Theorem 2.8, the categories $\text{Gen}({}_A J)$ and $\text{Gen}({}_S I)$ are equivalent.*

Proof: Let $B = FCM_H(S)$ and $C = M_H(S)$. Let $P = Cp_e$ being p_e the $H \times H$ -matrix having 1 in S at the (e, e) -position and zeros elsewhere.

By Lemma 3.9, $F = \text{Hom}({}_C P, -) : \text{Gen}({}_C P) \rightarrow S\text{-mod}$ is an equivalence of categories. Let $Q = FM_X(J)p_e$. Then $F(Q) \cong I$ and hence F induces an equivalence of categories between $\text{Gen}({}_C Q)$ and $\text{Gen}({}_S I)$.

Let $K = FM_H(I)$. K is isomorphic to $Q^{(H)}$ as a left C -module. Therefore $\text{Gen}({}_C K) = \text{Gen}({}_C Q)$.

By Lemma 3.5 $\text{Gen}({}_C K)$ is equivalent to $\text{Gen}({}_B K)$.

Now, by using the isomorphism Φ of Theorem 3.8, we have that $\text{Gen}({}_B K)$ is equivalent to $\text{Gen}({}_A J)$.

We conclude that $\text{Gen}({}_B J)$ is equivalent to $\text{Gen}({}_S I)$. ■

We finish this section with a version of Cohen-Montgomery Duality Theorem for actions.

Theorem 3.11. *Let $R * G$ be a crossed product and H a subgroup of G . Let $P = \bigoplus_{C \in G/H} R(C)$. Then $\text{End}_{gr-(R,G/H)}(P)$ is isomorphic to $FCM_{G/H}(R * H)$.*

Proof: For every $g \in G$ there is a unit \bar{g} in $R * G$ of degree g . Then, for every $g \in G$ and every $C \in G/H$, left multiplication by \bar{g} is a graded isomorphism between $R(C)$ and $R(gC)$. Therefore $P \cong R(H)^{(G/H)}$ in $(G/H, R) - gr$ and hence $\text{End}_{gr-(R,G/H)}(P)$ is isomorphic to $FCM_{G/H}(\text{End}_{gr-(R,G/H)}(R(H)))$. Finally, $\text{End}_{gr-(R,G/H)}(R(H)) \cong R * H$, by Lemma 1.5. ■

4. Separable functors

Separable functors have been introduced in [NVV] and studies in [NVV] and [Ra]. Let us recall the definition.

Definition. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor between two arbitrary categories \mathcal{C} and \mathcal{D} . F is said to be separable if for any two objects $C, C' \in \mathcal{C}$, there is a map $\varphi = \varphi_{C,C'} : \text{Hom}_{\mathcal{D}}(FC, FC') \rightarrow \text{Hom}_{\mathcal{C}}(C, C')$ such that:

SF1. For all $f \in \text{Hom}_{\mathcal{C}}(C, C')$, $\varphi(Ff) = f$.

SF2. For every commutative diagram in \mathcal{D} of the type

$$\begin{array}{ccc} FC & \xrightarrow{h} & FC_2 \\ F(f) \downarrow & & \downarrow F(g) \\ FC'_1 & \xrightarrow{k} & FC'_2 \end{array}$$

the following diagram in \mathcal{C} is also commutative

$$\begin{array}{ccc} C_1 & \xrightarrow{\varphi(h)} & C_2 \\ f \downarrow & & \downarrow g \\ C'_1 & \xrightarrow{\varphi(k)} & C'_2 \end{array}$$

In this section we first recall a fundamental theorem of Rafael [Ra] on adjoint separable functors. Then we study conditions for the functors T^* and T_* , defined in Section 1, to be separable. When $R = R'$ and ρ is the identity map on R , then T^* is always separable while the separability test for T_* can be simplified.

In [Ra] the authors give a complete characterization of separable adjoint functors in terms of the unit and counit of the adjunction. Explicitly they obtain the following result:

Theorem 4.1 [Ra]. *Let $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ be functors such that F is left adjoint of G . Let $u : 1_{\mathcal{C}} \rightarrow GF$ be the unit and $c : FG \rightarrow 1_{\mathcal{D}}$ the counit of the adjunction.*

- (a) *F is separable if and only if u splits (i.e. There exists a natural transformation $\varepsilon : GF \rightarrow 1_{\mathcal{C}}$ such that $\varepsilon_C \circ u_C = 1_C$ for every $C \in \mathcal{C}$).*
- (b) *G is separable if and only if c cosplits (i.e. There exists a natural transformation $\mu : 1_{\mathcal{D}} \rightarrow FG$ such that $c_D \circ \mu_D = 1_D$ for every $D \in \mathcal{D}$).*

For the sake of completeness we give a proof different from the one given in [Ra].

Proof: By duality it is enough to prove (a).

Assume that F is separable. For every $C, C' \in \mathcal{C}$ let

$$\varphi = \varphi_{C,C'} : \text{Hom}_{\mathcal{D}}(FC, FC') \longrightarrow \text{Hom}_{\mathcal{C}}(C, C')$$

be a map satisfying SF1 and SF2.

For every $C \in \mathcal{C}$, set $\epsilon_C = \varphi(c_{FC}) \in \text{Hom}_{\mathcal{C}}(GF(C), C)$.

First we prove that $\epsilon : GF \rightarrow 1_{\mathcal{C}}$ is a natural transformation. Indeed, let $f \in \text{Hom}_{\mathcal{C}}(C, C')$, then the diagram

$$\begin{array}{ccc} FGFC & \xrightarrow{FGF(f)} & FGFC' \\ c_{FC} \downarrow & & \downarrow c_{FC'} \\ FC & \xrightarrow{F(f)} & FC' \end{array}$$

is commutative, because c is a natural transformation. Therefore the following diagram is also commutative:

$$\begin{array}{ccc} GFC & \xrightarrow{GF(f)} & GFC' \\ \epsilon_C \downarrow & & \downarrow \epsilon_{C'} \\ C & \xrightarrow{f} & C' \end{array}$$

Now we prove that $\epsilon_C u_C = 1_C$ for every $C \in \mathcal{C}$. Indeed, the well known formula $c_{FC} \circ F(u_C) = 1_{FC}$ implies the commutativity of the following diagram

$$\begin{array}{ccc} FC & \xrightarrow{F(u_C)} & FGFC \\ 1_{FC} \downarrow & & \downarrow c_{FC} \\ FC & \xrightarrow{F(1_C)} & FC \end{array}$$

and SF2 implies the commutativity of the following diagram:

$$\begin{array}{ccc} C & \xrightarrow{u_C} & GFC \\ 1_C \downarrow & & \downarrow \epsilon_C \\ C & \xrightarrow{1_C} & C \end{array}$$

Conversely, let $\epsilon : GF \rightarrow 1_{\mathcal{C}}$ be a natural transformation satisfying $\epsilon \circ u = 1$. For any $C, C' \in \mathcal{C}$ let $\varphi : \text{Hom}_{\mathcal{D}}(FC, FC') \rightarrow \text{Hom}_{\mathcal{C}}(C, C')$ given by $\varphi(f) = \epsilon_{C'} \circ G(f) \circ u_C$.

If $f \in \text{Hom}_{\mathcal{C}}(C, C')$, then $\varphi(Ff) = \epsilon_{C'} GF(f) u_C = \epsilon_{C'} u_C f = f$.

On the other hand, if the following diagram in \mathcal{D} is commutative

$$\begin{array}{ccc} FC_1 & \xrightarrow{h} & FC_2 \\ F(f) \downarrow & & \downarrow F(g) \\ FC'_1 & \xrightarrow{k} & FC'_2 \end{array}$$

then the diagram

$$\begin{array}{ccc} C_1 & \xrightarrow{\varphi(h)} & C_2 \\ f \downarrow & & \downarrow g \\ C'_1 & \xrightarrow{\varphi(k)} & C'_2 \end{array}$$

is also commutative because $g \circ \varphi(h) = g \circ \varepsilon_{C_2} \circ G(h) \circ u_{C_1} = \varepsilon_{C'_2} \circ GF(g) \circ G(h) \circ u_{C_1} = \varepsilon_{C'_2} \circ G(F(g) \circ h) \circ u_{C_1} = \varepsilon_{C'_2} \circ G(k) \circ GF(f) \circ u_{C_1} = \varepsilon_{C'_2} \circ G(k) \circ u_{C'_1} \circ f = \varphi(k) \circ f$. ■

Theorem 4.1 can be used to give a simple criterion on when a right adjoint functor of the form $G : gr - (R, X, G) \rightarrow \mathcal{D}$ is separable, for \mathcal{D} an additive category. Next proposition was inspired from Proposition 2.2 in [Ra].

Proposition 4.2. *Let \mathcal{C} be a full subcategory of $gr - (R, X, G)$ such that $R(x) \in \mathcal{C}$ for every $x \in X$. Let $G : \mathcal{C} \rightarrow \mathcal{D}$ be a right adjoint functor, F a left adjoint of G and $c : FG \rightarrow 1_{\mathcal{C}}$ the counit of the adjunction. The following conditions are equivalent:*

- (a) G is separable.
- (b) For any $x \in X$ there exists $m_x \in FG(R(x))_x$ satisfying the following two conditions:
 - (1) For any $x \in X$, $c_{R(x)}(m_x) = 1$.
 - (2) For any $x \in X$, $g \in G$ and $r \in R_g$, $FG(\lambda_r)(m_{xg}) = m_x r$ (where $\lambda_r \in \text{Hom}_{gr-(R, X, G)}(R(xg), R(x))$ is the left multiplication-by- r map).

Proof: Assume first that G is separable. By Theorem 4.1, there exists a natural transformation $\mu : 1_{\mathcal{C}} \rightarrow FG$ such that $c \circ \mu = 1$. By Lemma 1.5 for every $x \in X$ there exists $m_x \in FG(R(x))$ such that $\mu_{R(x)} = \lambda_{m_x}$. Then $c_{R(x)}(m_x) = (c_{R(x)} \circ \mu_{R(x)})(1) = 1$. On the other hand, if $r \in R_g$, then for every $x \in X$ the following diagram

$$\begin{array}{ccc} R(xg) & \xrightarrow{\mu_{R(xg)}} & FG(R(xg)) \\ \lambda_r \downarrow & & \downarrow FG(\lambda_r) \\ R(x) & \xrightarrow{\mu_{R(x)}} & FG(R(x)) \end{array}$$

is commutative. Therefore $FG(\lambda_r)(m_{xg}) = [FG(\lambda_r) \circ \mu_{R(xg)}](1) = [\mu_{R(x)} \circ \lambda_r](1) = m_x r$.

Let now $\{m_x \in FG(x)_x | x \in X\}$ be a system of elements satisfying conditions (1) and (2). For every $N \in gr - (R, X, G)$ define $\mu_N : N \rightarrow FG(N)$ by $\mu_N(n) = FG(\lambda_n)(m_x)$ if $n \in N_x$ and extend μ_N to N by linearity.

μ_M is R -linear because if $n \in N_x$ and $r \in R_g$, then $\mu_N(nr) = FG(\lambda_{nr})(m_{xg}) = FG(\lambda_n)FG(\lambda_r)(m_{xg}) = FG(\lambda_n)(m_{xr}) = FG(\lambda_n)(m_x)r = \mu_N(n)r$.

Now we prove that μ is a natural transformation. Let $f \in \text{Hom}_{gr-(R, X, G)}(N, N')$ and $n \in N_x$. Then

$$\begin{aligned} [FG(f) \circ \mu_N](n) &= FG(f \circ \lambda_n)(m_x) = FG(\lambda_{f(n)})(m_x) = \\ &= \mu_{N'}(f(n)) = (\mu_{N'} \circ f)(n). \end{aligned}$$

Finally, if $N \in \mathcal{C}$ and $n \in N_x$, then $(c_N \circ \mu_N)(n) = [c_N \circ FG(\lambda_n)](m_x) = [\lambda_n \circ c_{R(x)}](m_x) = \lambda_n(1) = n$.

Thus, applying Theorem 4.1 we conclude that G is separable. ■

For the rest of this section we consider the following situation: Let $T = (\rho, \xi, \gamma)$ where $\rho : R \rightarrow R'$ is a ring homomorphism, $\xi : X \rightarrow X'$ is a map, $\gamma : G \rightarrow G'$ is a group homomorphism, R is a G -graded ring, R' is a G' -graded ring, X is a right G -set, X' is a right G' -set and the following conditions hold:

$$\rho(R_g) = R'_{\gamma(g)} \quad \xi(xg) = \xi(x)\gamma(g) \quad (g \in G, x \in X).$$

Let T^* and T_* be the functors defined in Section 1. We are going to study when T^* and T_* are separable.

Proposition 4.3. *Let $T = (\rho, \xi, \gamma)$ as above. The following conditions are equivalent:*

- (1) T^* is separable.
- (2) For every $x \in X$ there is $\varepsilon_x \in \text{Hom}_{gr-(R, X, G)}(T_*(R'(\xi(x)), R(x)))$ such that:
 - (a) For every $x \in X$, $\varepsilon_x(1^x) = 1$.
 - (b) For every $x, y \in X$, $g \in G$, $r \in R_g$ and $r' \in R'(\xi(xg))_{\xi(y)}$,

$$\varepsilon_x((\rho(r)r')^y) = r\varepsilon_{xg}((r')^y).$$

Proof: First let us note that the unit u of the adjunction pair (T^*, T_*) is given by:

$$\begin{aligned} u_M : M &\longrightarrow T_*T^*(M) \quad (M \in gr - (R, X, G)) \\ m &\longmapsto \sum_{x \in X} (m_x \otimes 1)^x \quad (m \in M). \end{aligned}$$

On the other hand the canonical isomorphism $R' \rightarrow R \otimes_R R'$ induces an isomorphism $\phi_x : R'(\xi(x)) \rightarrow T^*(R(x))$ for every $x \in X$.

(1) \Rightarrow (2) Assume that T^* is separable. By Theorem 4.1 there exists a natural transformation $\varepsilon : T_*T^* \rightarrow 1_{gr-(R,X,G)}$ such that $\varepsilon_M \circ u_M = 1_M$ for every $M \in gr-(R,X,G)$.

For every $x \in X$ let $\varepsilon_x = \varepsilon_{R(x)} \circ T_*(\phi_x)$. We are going to see that $\{\varepsilon_x | x \in X\}$ satisfies conditions (a) and (b).

$$\varepsilon_x(1^x) = \varepsilon_{R(x)}((1 \otimes 1)^x) = [\varepsilon_{R(x)} \circ u_{R(x)}](1) = 1.$$

Let now $g, h \in G$, $r \in R_g$ and $r' \in R'(\xi(xg))_{\xi(y)}$. Let $\lambda_r : R \rightarrow R$ be the left multiplication by r map and $\lambda_{\rho(r)} : R' \rightarrow R'$ left multiplication by $\rho(r)$. By Lemma 1.3, one has

$$\lambda_r \in \text{Hom}_{gr-(R,X,G)}(R(xg), R(x))$$

and

$$\lambda_{\rho(r)} \in \text{Hom}_{gr-(R',X',G')}(R'(\xi(xg)), R'(\xi(x))).$$

Furthermore the following diagram is commutative

$$\begin{array}{ccccc} T_*(R'(\xi(xg))) & \xrightarrow{T_*(\phi_{xg})} & T_*T^*(R(xg)) & \xrightarrow{\varepsilon_{R(xg)}} & R(xg) \\ T_*(\lambda_{\rho(r)}) \downarrow & & \downarrow T_*T^*(\lambda_r) & & \downarrow \lambda_r \\ T_*(R'(\xi(x))) & \xrightarrow{T_*(\phi_x)} & T_*T^*(R(x)) & \xrightarrow{\varepsilon_{R(x)}} & R(x) \end{array}$$

Therefore, $\varepsilon_x((\rho(r)r')^y) = [\varepsilon_{R(x)} \circ T_*(\phi_x)]((\lambda_{\rho(r)}(r'))^y) = [\varepsilon_{R(x)} \circ T_*(\phi_x)]([T_*(\lambda_{\rho(r)})]((r')^y)) = [\lambda_r \circ \varepsilon_{xg}]((r')^y) = r\varepsilon_{xg}((r')^y)$.

(2) \Rightarrow (1) Assume that for every $x \in X$ there exists $\varepsilon_x \in \text{Hom}_{(G,X,R)-gr}(T_*(R'(\xi(x))), R(x))$ satisfying conditions (a) and (b). For every $M \in gr-(R,X,G)$ let $\varepsilon_M : T_*T^*(M) \rightarrow M$ be the homomorphism given by $\varepsilon_M((m \otimes r')^y) = m\varepsilon_x((r')^y)$ whenever $m \in M_x$ and $r' \in R'(\xi(x))_{\xi(y)}$ and extended by linearity. Note that ε_M is well defined. Indeed, let $m \in M_x$, $r \in R_g$ and $r' \in R'(\xi(xg))_{\xi(y)}$. Then $\varepsilon_M(mr \otimes r') = mr\varepsilon_{xg}((r')^y) = m\varepsilon_x((\rho(r)r')^y) = \varepsilon_M((m \otimes rr')^y)$.

It is an easy exercise to check that $\varepsilon_M \in \text{Hom}_{gr-(R,X,G)}(T_*T^*(M), M)$. Moreover, for every $m \in M_x$, $[\varepsilon_M \circ u_M](m) = \varepsilon_M((m \otimes 1)^x) = m\varepsilon_x(1^x) = m$. It only remains to prove that ε is a natural transformation from T_*T^* to $1_{gr-(R,X,G)}$ and apply Theorem 4.1.

Let $f \in \text{Hom}_{gr-(R,X,G)}(M, N)$, $m \in M_x$ and $r' \in R'(\xi(x))_{\xi(y)}$. Then $[\varepsilon_N \circ T_*T^*(f)]((m \otimes r')^y) = \varepsilon_N((f(m) \otimes r')^y) = f(m)\varepsilon_x((r')^y) = f(m\varepsilon_x((r')^y)) = f(\varepsilon_M((m \otimes r')^y)) = [f \circ \varepsilon_M]((m \otimes r')^y)$. ■

Proposition 4.4. *Let $T = (\rho, \xi, \gamma)$ as above. The following conditions are equivalent:*

- (a) T_* is separable.
 (2) There exists a system

$$\sum = \{ (a'_{(x',i)}, b'_{(x',i)}, x_{(x',i)}) \in R'_{g'_{(x',i)}} \times R'_{(g'_{(x',i)})^{-1}} \times X \mid x' \in X', \\ i = 1, 2, \dots, m_{x'} \}$$

satisfying the following conditions.

- (1) For every $x' \in X'$ and $i = 1, 2, \dots, m_{x'}$, $x'g'_{(x',i)} = \xi(x_{(x',i)})$.
 (2) For every $x' \in X'$, $\sum_{i=1}^{m_x} a'_{(x',i)} b'_{(x',i)} = 1$.
 (3) For every $r' \in R'_{g'}$, and every $x' \in X'$, the following equality holds in $T_*(R'(x')) \otimes_R R'$.

$$\sum_{i=1}^{m_{x'}} (a'_{(x',i)})^{x_{(x',i)}} \otimes (b'_{(x',i)}) r' = \sum_{i=1}^{m_{x'g'}} (r' a'_{(x'g',i)})^{x_{(x'g',i)}} \otimes (b'_{(g'x',i)}).$$

Proof: First let us remark that the counit $c : T^*T_* \rightarrow 1$ of the adjunction pair (T^*, T_*) is given by:

$$c_N : T_*(N) \otimes_R R' \longrightarrow N \quad (N \in gr - (R', X', G')) \\ c_N(n^x \otimes r') = nr' \quad (r' \in R', n \in N_{\xi(x)}).$$

(a) \Rightarrow (b) Assume that T_* is separable. By Proposition 4.2, for every $x' \in X'$ there exists $K_{x'} \in (T_*(R'(x')) \otimes_R R')_{x'}$ such that

- (1) $c_{R'(x')}(K_{x'}) = 1$.
 (2) For any $x' \in X'$, $g' \in G'$ and $r' \in R'_{g'}$, $T^*T_*(\lambda_{r'})(K_{x'g'}) = K_{x'r'}$.

Each $K_{x'}$ can be expressed as $\sum_{i=1}^{n_x} (\alpha'_{(x',i)})^{x_{(x',i)}} \otimes \beta'_{(x',i)}$ where $\alpha'_{(x',i)}$ and $\beta'_{(x',i)}$ are homogeneous for every possible x' and i . For every possible x' and i , let $g'_{(x',i)}$ be the degree of $\alpha'_{(x',i)}$ and $h'_{(x',i)}$ the degree of $\beta'_{(x',i)}$. Then $x'g'_{(x',i)} = \xi(x_{(x',i)})$ and $\xi(x_{(x',i)})h'_{(x',i)} = x'$.

Let $\sum = \{ (a'_{(x',i)}, b'_{(x',i)}, x_{(x',i)}) \mid x' \in X', i = 1, 2, \dots, m_x \}$ be the system which results by eliminating in the system $\{ (\alpha'_{(x',i)}, \beta'_{(x',i)}, x_{(x',i)}) \mid x' \in X', i = 1, 2, \dots, n_{x'} \}$ any upla $\{ \alpha'_{(x',i)}, \beta'_{(x',i)}, x_{(x',i)} \}$ such that $g'_{(x',i)}h'_{(x',i)} \neq e$. Plainly \sum satisfies (1).

Since $1 = c_{R'(x')}(K_{x'}) = \sum_{i=1}^{n_{x'}} \alpha'_{(x',i)} \beta'_{(x',i)}$, then the homogeneous component of degree e of this element is equal to 1. But this homogeneous component is just $\sum_{i=1}^{m_{x'}} a'_{(x',i)} b'_{(x',i)}$. Thus \sum satisfies (2).

Consider G' as a right G -set by: $g'g = g'\gamma(g)$ ($g \in G, g' \in G'$). Consider in $M = T_*(R'(x'))$ the following G' -grading: A typical element r'^x in $T_*(R'(x'))_x$ has degree $g' \in G'$ if $r' \in R'_{g'}$. Plainly $M \in gr-(R, G', G)$. Let $S = (\rho, 1_{G'}, \gamma)$. Then $S^*(M) = M \otimes_R R' \in gr-(R', G', G')$ where G' is considered as a right G' -set by regular action.

Given $r' \in R'_{g'}$ and $x' \in X'$, then $A = \sum_{i=1}^{n_{x'}} (\alpha'_{(x',i)})^{x_{(x',i)}} \otimes \beta'_{(x',i)} r' = K_{x'} r' = T^* T_*(r') K_{x'g'} = \sum_{i=1}^{n_{x'g'}} (r' \alpha'_{(x'g',i)})^{x_{(x'g',i)}} \otimes \beta'_{(x'g',i)}$. Consider A as an element of $S_*(M)$. For every $i = 1, 2, \dots, n_{x'}$

$$(r' \alpha'_{(x',i)})^{x_{(x',i)}} \otimes \beta'_{(x',i)} \in S^*(M)_{g'g_{(x',i)} h'_{(x',i)}}$$

and

$$(\alpha'_{(x'g',i)})^{x_{(x'g',i)}} \otimes \beta'_{(x'g',i)} r' \in S^*(M)_{g'_{(x',i)} h'_{(x',i)} g'}.$$

Therefore,

$$\sum_{i=1}^{m_{x'}} (\alpha'_{(x',i)})^{x_{(x',i)}} \otimes b'_{(x',i)} r' = A_{g'} = \sum_{i=1}^{m_{x'g'}} (r' \alpha'_{(x'g',i)})^{x_{(x'g',i)}} \otimes a'_{(x'g',i)}.$$

This proves (3) for \sum .

(b) \Rightarrow (a) Let

$$\sum = \{ \{ (a'_{(x',i)}, b'_{(x',i)}, x_{(x',i)}) \in R'_{g'_{(x',i)}} \times R'_{(g'_{(x',i)})^{-1}} \times X | x' \in X', \\ i = 1, 2, \dots, n_{x'} \}$$

be a system satisfying (1), (2) and (3).

For all $x' \in X'$, let $K_{x'} = \sum_{i=1}^{m_{x'}} (a'_{(x',i)})^{x_{(x',i)}} \otimes b'_{(x',i)}$. Then $c_{R'(x')}(K_{x'}) = \sum_{i=1}^{m_{x'}} a'_{(x',i)} b'_{(x',i)} = 1$ and for every $r' \in R'_{g'}$, $T^* T_*(\lambda_{r'})(K_{x'g'}) = \sum_{i=1}^{m_{x'g'}} (r' \alpha'_{(x'g',i)})^{x_{(x'g',i)}} \otimes a'_{(x'g',i)} = \sum_{i=1}^{m_{x'}} (a'_{(x',i)})^{x_{(x',i)}} \otimes a'_{(x',i)} r' = K_{x'} r'$.

By Proposition 4.2, we concluded that T_* is separable. ■

In the proof of last Proposition, we have considered G' as a right G -set by $g'g = g'\gamma(g)$ ($g \in G, G' \in G'$). Actually we can consider G' as a G - G -set by putting $gg' = \gamma(g)g'$.

For every $x' \in X'$, the right R -module $T_*(R'(x'))$ becomes an object of $gr - (R, G', G)$ by assigning degree g' to r'^x whenever $r' \in R'_{g'}$, with $x'g' = \xi(x)$.

We can consider X as a left G -set by $gx = xg^{-1}$ ($g \in G, x \in X$). Similarly X' can be considered as a left G' -set.

Moreover $\gamma(g)\xi(x) = \xi(gx)$ for every $g \in G$ and $x \in X$. Therefore, we can define two functors $T^0 = R' \otimes_R - : (G, X, R) - gr \rightarrow (G', X', R') - gr$ and $T_0 : (G', X', R') - gr \rightarrow (G, X, R) - gr$ which are the left hand versions of T^* and T_* . By symmetry, for every $x' \in X'$, $T_0((x')R')$ is an object of $(G, G', R) - gr$ by assigning degree g' to r'^x whenever $r' \in R'_{g'}$, with $g'x' = \xi(x)$.

Set $S = (\rho, 1_{G'}, \gamma)$. For every $x' \in X'$, $S^*(T_*(R'(x'))) = T_*(R'(x')) \otimes_R R' \in gr - (R', G', G') = gr - R'$ and $S^0(T_0((x')R')) \in (G', G', R') - gr = R - gr$.

Lemma 4.5. *For every $x' \in X'$ and every $g' \in G'$, $S^*(T_*(R'(x')))$ is isomorphic to ${}_g S^0(T_0((g'x')R'))$.*

Proof: Let $\Phi : S^*(T_*(R'(x')))_g \rightarrow {}_g S^0((g'x')(R'))$ the homomorphism given by $\Phi(a'^x \otimes b') = a' \otimes b'^x$ ($x \in X, a' \in R'_{g'}$, $b' \in R'_{g'}$, with $x'\sigma' = \xi(x)$ and $\sigma'\tau' = g'$) and extended by linearity.

Φ is well defined. Indeed, let $a' \in R'_{g'}$, $b' \in R'_{g'}$, $r \in R_g$ and assume that $\sigma'\gamma(g)\tau' = g'$ and $x'\sigma' = \xi(x)$. Then $\Phi((a')^x r \otimes b') = \Phi((a'\rho(r))^{xg} \otimes b') = a'\rho(r) \otimes (b')^{xg} = a' \otimes r(b')^{g^{-1}x} = a' \otimes (\rho(r)b')^x = \Phi(a'^x \otimes \rho(r)b)$.

And Φ is obviously an isomorphism. ■

Proposition 4.6. *Let $T = (\rho, \xi, \gamma)$ as above. T_* is separable if and only if T_0 is separable.*

Proof: By left-right symmetry, T_0 is separable if and only if there exists a system

$$\sum = \{(a'_{(x',i)}, b'_{(x',i)}, x_{(x',i)}) \in R'_{(h'_{(x',i)})^{-1}} \times R_{h'_{(x',i)}} \times X | x' \in X', \\ i = 1, 2, \dots, m_{x'}\}$$

satisfying the following conditions.

(1') For every $x' \in X'$ and $i = 1, 2, \dots, m_{x'}$, $h'_{(x',i)}x' = \xi(x_{(x',i)})$.

(2') For every $x' \in X'$, $\sum_{i=1}^{m_{x'}} a'_{(x',i)} b'_{(x',i)} = 1$.

(3') For every $r' \in R'_h$, and every $y' \in X'$, the following equality holds in $R' \otimes_R T_0((x')R')$

$$\sum_{i=1}^{m_{h'y'}} a'_{(h'y',i)} \otimes (b'_{(h'y',i)} r')^{x_{(h'y',i)}} = \sum_{i=1}^{m_{y'}} r' a'_{(y',i)} \otimes (b'_{(y',i)})^{x_{y',i}}.$$

Assume that T_* is separable. Then there exists a system

$$\sum = \{ (a'_{(x',i)}, b'_{(x',i)}, x_{(x',i)}) \in R'_{g'_{(x',i)}} \times R'_{(g'_{(x',i)})^{-1}} \times X | x' \in X', \\ i = 1, 2, \dots, m_{x'} \}$$

satisfying conditions (1), (2) and (3) of Proposition 4.4. For every $x' \in X'$ and every $i = 1, 2, \dots, m_{x'}$, let $h'_{(x',i)} = (g'_{(x',i)})^{-1}$. Then

$$\sum = \{ (a'_{(x',i)}, b'_{(x',i)}, x_{(x',i)}) \in R'_{(h'_{(x',i)})^{-1}} \times R_{h'_{(x',i)}} \times X | x' \in X', \\ i = 1, 2, \dots, m_{x'} \}$$

satisfies conditions (1') and (2'). On the other hand, let $r' \in R'_{h'}$ and $y' \in X$. Set $g' = h'^{-1}$ and $x' = h'y'$. By assumption the following equality holds in $T_*(R(x')) \otimes_R R'$.

$$\sum_{i=1}^{m_{x'}} (a'_{(x',i)})^{x_{(x',i)}} \otimes b'_{(x',i)} r' = \sum_{i=1}^{m_{x'h'}} (r' a'_{(x'h',i)})^{x_{(x'h',i)}} \otimes (b'_{(x'h',i)})$$

and this element has degree h' when we view $T_*(R(x')) \otimes_R R'$ as $S^*(T_*(R(x')))$. Therefore, by using the isomorphism of Lemma 4.5, the following equality holds in $R' \otimes_R T_0((y')R')$

$$\sum_{i=1}^{m_{h'y'}} a'_{(h'y',i)} \otimes (b'_{(h'y',i)} r')^{x_{(h'y',i)}} = \sum_{i=1}^{m_{y'}} r' a'_{(y',i)} \otimes (b'_{(y',i)})^{x_{y',i}}.$$

We conclude that T_0 is separable. ■

Let us consider now the particular case when $R = R'$ and ρ is the identity map of R . By Lemma 1.8, in order to study T^* and T_* we can assume that $G = G'$ and γ is the identity map of G . In that case $T^* = T_\xi$ and $T_* = S^\xi$.

Corollary 4.7. *Let R be a G -graded ring and $\xi : X \rightarrow X'$ a morphism of right G -sets. The functor T_ξ is separable.*

Proof: For every $x \in X$ let $\varepsilon_x : S^\xi(R(\xi(x))) \rightarrow R(x)$ given by $\varepsilon_x(r^y) = r_y$ (where the y -th component of r is computed in $R(x)$). Plainly ε_x satisfies condition (a) of Proposition 4.3. On the other hand, if $x, y \in X, g \in G, r \in R_g$ and $r' \in R(\xi(x))_{\xi(y)}$, then $\varepsilon_x((rr')^y) = (rr')_y = \sum_{xh=y} (rr')_h = \sum_{xh=y} r(r')_{g^{-1}h} = r \sum_{xgh=y} r'_h = r\varepsilon_{xg}(r'^y)$. Now Proposition 4.3 applies. ■

Corollary 4.8. *Let $T = (\rho, \xi, \gamma)$ as above. If R is a direct summand of R' as R - R -bimodules, then T^* is separable.*

Proof: Consider the following commutative diagram of functors

$$\begin{array}{ccc} gr - (R, X, G) & \xrightarrow{T^*} & gr - (R', X', G') \\ U_R \downarrow & & \downarrow U_{R'} \\ \text{mod} - R & \xrightarrow{-\otimes_R R'} & \text{mod} - R' \end{array}$$

where U_R and $U_{R'}$ are the functor which forget the grading. Let Y be a singleton and consider G acting on Y . Let $\varphi : X \rightarrow Y$ be the only possible map. Then $\text{mod} - R = gr - (R, Y, G)$ and $U_R = T_\varphi$. Therefore U_R is separable.

By Proposition 1.3 of [NVV], $-\otimes_R R'$ is separable. Thus $U_{R'} \circ T^* = (-\otimes_R R') \circ U_R$ is separable and hence T^* is separable (see [NVV, Lemma 1.1]). ■

Corollary 4.9. *Let R be a G -graded ring and $\xi : X \rightarrow X'$ a morphism of right G -sets. The following conditions are equivalent:*

- (a) S^ξ is separable.
- (b) For every $x \in X$ there exists $a^x \in R_e$ such that:
 - (SS1) For every $x' \in X'$, the set $\{x \in \xi^{-1}(x') | a^x \neq 0\}$ is finite and

$$\sum_{\xi(x)=x'} a^x = 1.$$

- (SS2) For every $r \in R_g$ and $\chi \in X, a^{\chi}r = ra^{\chi g}$.

Proof: (a) \Rightarrow (b) Assume that S^ξ is separable. Let

$$\sum = \{ (a_{(x',i)}, b_{(x',i)}, x_{(x',i)}) \in R_{g_{(x',i)}} \times R_{(g_{(x',i)})^{-1}} \times X | x' \in X', i = 1, 2, \dots, m_x \}$$

be a system satisfying conditions (1), (2) and (3) of Proposition 4.4(b). For any $x \in X$ let

$$a^x = \sum \{a_{(\xi(x),i)} b_{(\xi(x),i)} | i = 1, 2, \dots, m_{\xi(x)}, x_{(\xi(x),i)} = xg_{(x',i)}\}.$$

If we fix an $x' \in X'$, then for every $x \in \xi^{-1}(x')$, $a^x = 0$, unless $x \in \{x_{(x',i)}(g_{(x',i)})^{-1} | i = 1, 2, \dots, m_{x'}\}$. Therefore, $a^x = 0$ for almost all $x \in \xi^{-1}(x')$.

Moreover, for a fixed $x' \in X'$, $\sum_{\xi(x)=x'} a^x = \sum_{i=1}^{m_{x'}} a_{(x',i)} b_{(x',i)} = 1$. Note that this implies that ξ has to be surjective.

On the other hand, if we identify $T_*(R(x')) \otimes_R R$ with $T_*(R(x'))$ by the canonical isomorphism, then for every $r \in R_g$,

$$(*) \quad \sum_{i=1}^{m_{x'g}} (ra_{(x'g,i)})^{x_{(x'g,i)}} b_{(x'g,i)} = \sum_{i=1}^{m_{x'}} (a_{(x',i)})^{x_{(x',i)}} b_{(x',i)} r.$$

The left hand term in the previous expression is

$$\begin{aligned} \mathcal{L} &= \sum_{i=1}^{m_{x'g}} (ra_{(x',i)})^{x_{(x'g,i)}} b_{(x'g,i)} = \\ &= \sum_{i=1}^{m_{x'g}} (ra_{(x'g,i)} b_{(x'g,i)})^{x_{(x'g,i)}} (g_{(x'g,i)})^{-1} = \sum_{\xi(x)=x'g} (ra^x)^x. \end{aligned}$$

Arguing in a similar way we obtain the following expression for the right hand term in (*).

$$\mathcal{R} = \sum_{i=1}^{m_{x'}} (r_{x',i})^{x_{x',i}} s_{x',i} r = \sum_{\xi(x)=x'} (a^x)^x r = \sum_{\xi(x)=x'} (a^x r)^{xg}.$$

Therefore $(ra^{xg})^{xg} = \mathcal{L}_{xg} = \mathcal{R}_{xg} = (a^x r)^{xg}$ and hence $ra^{xg} = a^x r$ for every $x \in X$.

(b) \Rightarrow (a) Let $\{a^x | a \in A\}$ be a system satisfying conditions (SS1) and (SS2) of (b). For every $x' \in X'$ and any $x \in \xi^{-1}(x')$, let $a_{(x',x)} = a^x$ and $b_{(x',x)} = 1$. Then the system $\{(a_{(x',x)}, b_{(x',x)}, x) | x' \in X', x \in \xi^{-1}(x'), a^x \neq 0\}$ satisfies conditions (1), (2) and (3) of Proposition 4.4(b). Indeed, conditions (1) and (2) are obvious. Let now $0 \neq r \in R_g$.

Then

$$\begin{aligned} \sum_{x \in \xi^{-1}(x')} (a_{(x',x)})^x \otimes b_{(x',x)} r &= \sum_{x \in \xi^{-1}(x')} (a^x)^x r = \sum_{x \in \xi^{-1}(x')} (a^x r)^{xg} = \\ &= \sum_{x \in \xi^{-1}(x')} (ra^{xg})^{xg} = \sum_{y \in \xi^{-1}(x'g)} (ra^y)^y = \\ &= \sum_{x \in \xi^{-1}(x'g)} (ra_{(x'g,x)})^x \otimes b_{(x'g,x)}. \end{aligned}$$

Proposition 4.4 applies to finish the proof. ■

Definition. Let $\xi : X \rightarrow X'$ be a morphism of right G -sets. A ξ -separability system in a G -graded ring R is a system $\{a^x | x \in X\}$ of elements in R_e satisfying conditions (SS1) and (SS2) of Corollary 4.9.

Remarks. (1) Note that if $X = G$, X' is a singleton and $\xi : G \rightarrow X'$ is the unique possible map, then R has a ξ -separability system if and only if R has a separability system in the sense of [Ra].

(2) Let $\xi : X \rightarrow X'$ be a morphism of right G -sets and R a G -graded ring.

- (a) If S_ξ is separable (or equivalently, if R has a ξ -separability system), then ξ is epic.
- (b) If ξ is epic and for every $x' \in X'$, $\xi^{-1}(x')$ is finite with order invertible in R , then $a^x = |\xi^{-1}(\xi(x))|^{-1}$ is a ξ -separability system whenever $|\xi^{-1}(x')|$ is constant in each orbit of X' .
- (c) Assume that ξ is epic. For every orbit Y in X let $\xi_Y : Y \rightarrow \xi(Y)$ be the restriction of ξ to Y . R has a ξ -separability system if and only if it has a ξ_Y -separability system for every orbit Y of X . Namely, if for every orbit Y in X , $\{a^y | y \in Y\}$ is a ξ_Y -separability system for R , then $\{a^x | x \in X\}$ is a ξ -separability system for R .

5. Weak dimension

Let G be a group, H a subgroup of G and R a G -graded ring.

Let G/H be the set of left H -cosets and $\varphi_H : G \rightarrow G/H$ the canonical projection. In order to simplify the notation let us denote by T_H and S_H the functors $T_{\varphi_H} : R - gr \rightarrow (R, G/H) - gr$ and $S^{\varphi_H} : (G/H, R) - gr \rightarrow R - gr$ respectively.

For a (graded) ring $(gr.)w.\dim(R)$ will denote the (graded) weak dimension of R . Similarly, for a R -module, $f.d.(M)$ will denote the flat dimension of M . Recall that, for a graded module ${}_R M$, the graded flat dimension coincides with the flat dimension (see e.g. [NV1]).

Next theorem extends Theorem 7 in [Rf1] and gives an approach to answer Question 2 in the same paper.

Theorem 5.1. *Let R be a G -graded ring. Assume that for every finitely generated subgroup H of G the functor S_H is separable. Then $\text{gr.w. dim}(R) = \text{w. dim}(R)$.*

Remark. Note that if S_H is separable, then the H -graded ring $R_{(H)} = \bigoplus_{gh \in H} R_h$ has a separability system (in the sense of [Ra]). Indeed, S_H is separable if and only if R has a φ_H -separability system. But if $\{x^g | g \in G\}$ is a φ_H -separability system, then $\{x^h | h \in H\}$ is a separability system for $R_{(H)}$.

The converse is not true in general as the following example shows.

Example. Let K be a field of characteristic 2. Let $G = Z_2 \times Z_2$ and H the subgroup of G generated by $(1, 1)$. Consider the ring R of the matrices of the form $\begin{pmatrix} a & b & c \\ 0 & a & 0 \\ 0 & 0 & a \end{pmatrix}$ with entries in K . R has a G -grading given by:

$$R_{(0,0)} = \left\{ \begin{pmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{pmatrix} \mid a \in K \right\}, \quad R_{(1,1)} = 0$$

$$R_{(1,0)} = \begin{pmatrix} 0 & K & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad R_{(0,1)} = \begin{pmatrix} 0 & 0 & K \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Then $\{a^{(0,0)} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, a^{(1,1)} = 0\}$ is a separability system for $R_{(H)}$. But R has not a φ_H -separability system. Indeed, if $\{a^g | g \in G\}$ is a φ_H -separability system, then $a^{(1,0)} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} a^{(0,0)}$ and $a^{(0,1)} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} a^{(0,0)}$. Therefore $a^{(1,0)} = a^{(0,0)} = a^{(0,1)}$ and $1 = a^{(0,1)} + a^{(1,0)} = 2a^{(1,0)} = 0$.

Lemma 5.2. *Let ${}_R P$ be a projective left R -module and $M \in \text{Gen}({}_R P)$. Then ${}_R M$ is projective if and only if it is projective in the category $\text{Gen}({}_R P)$.*

Proof: Let $f : K \rightarrow L$ be a homomorphism in the category $\text{Gen}({}_R P)$. If f is surjective, then f is an epimorphism in the category $\text{Gen}({}_R P)$.

Conversely, if f is an epimorphism in the category $\text{Gen}({}_R P)$ and C is the cokernel of f , then the canonical projection $p : L \rightarrow C$ is also a homomorphism in $\text{Gen}({}_R P)$ such that $p \circ f = 0$. Thus $C = 0$, this implies that f is surjective.

Since ${}_R M \in \text{Gen}({}_R P)$, then there exists an epimorphism $f : P^{(I)} \rightarrow M$ which is an epimorphism both in $\text{Gen}({}_R P)$ and $R\text{-mod}$. Thus ${}_R M$ is simultaneously projective in $R\text{-mod}$ and $\text{Gen}({}_R P)$. ■

Let \mathcal{C} be a Grothendieck category. Recall that an object F in \mathcal{C} is said to be flat if for every finitely presented object M in \mathcal{C} and every epimorphism $f : N \rightarrow F$, the correspondent homomorphism of abelian groups $\text{Hom}_{\mathcal{C}}(M, f) : \text{Hom}_{\mathcal{C}}(M, N) \rightarrow \text{Hom}_{\mathcal{C}}(M, F)$ is an epimorphism (see e.g. [Po]).

When I is an idempotent left ideal of a ring R , then $\text{Gen}({}_R I)$ is a Grothendieck category (see e.g. [W]) because it is closed under submodules. Of course, a left ideal with local units is an idempotent ideal.

Proposition 5.3. *Let I be a left ideal of R with local units. Then, an object F in $\text{Gen}({}_R I)$ is flat in this category if and only if it is flat as a left R -module.*

Proof: Since $\text{Gen}({}_R I)$ is closed under subobjects, the finitely generated (resp. finitely presented) objects of $\text{Gen}({}_R I)$ are just the finitely generated (resp. finitely presented) left R -modules generated by ${}_R I$. Moreover, as we have remarked in the proof of the previous lemma the epimorphisms in $\text{Gen}({}_R I)$ are the epimorphisms in $R\text{-mod}$ between objects in $\text{Gen}({}_R I)$. Therefore the necessary condition is obvious.

Let now F be a flat object in $\text{Gen}({}_R I)$. First we will see that for every idempotent $e \in I$, eF is flat as a left eRe -module. Consider Re as an $R - eRe$ -bimodule and the functors $H = \text{Hom}({}_R Re, -) : R\text{-mod} \rightarrow eRe\text{-mod}$ and $T = eR \otimes_{eRe} - : eRe\text{-mod} \rightarrow R\text{-Mod}$. Let $\mu : 1_{eRe\text{-mod}} \rightarrow HT$ and $\varepsilon : TH \rightarrow 1_{R\text{-mod}}$ be the unit and the counit of the adjoint pair (T, H) .

Let M be a finitely presented left eRe -module and $f : M \rightarrow eF$ a homomorphism of left eRe -modules, then $T(M)$ is a finitely presented left R -module in $\text{Gen}({}_R I)$. Let $p : I^{(X)} \rightarrow F$ be an epimorphism. By flatness of F in $\text{Gen}({}_R I)$ there exists $g : T(M) \rightarrow I^{(X)}$ such that $p \circ g = \varepsilon_F \circ T(f)$. Since $Re \otimes_{eRe} M$ is finitely generated, there exists $Y \subseteq X$ finite such that $\text{Im}(g) \subseteq I^{(Y)}$ and hence there exist $g' : T(M) \rightarrow I^{(Y)}$ and $p' : I^{(Y)} \rightarrow F$ such that $p' \circ g' = \varepsilon_F \circ T(f)$. Applying the functor

H , one has a commutative diagram

$$\begin{array}{ccccc}
 M & \xrightarrow{\mu_M} & HT(M) & \xrightarrow{H(g')} & H(I^{(Y)}) \simeq (eR)^{(Y)} \\
 f \downarrow & & HT(f) \downarrow & & \downarrow H(p') \\
 Fe & \xrightarrow{\mu_{Fe}} & HTH(F) & \xrightarrow{H(\varepsilon_F)} & H(F)
 \end{array}$$

but μ_X is an isomorphism for every $X \in eRe\text{-mod}$. On the other hand $H(\varepsilon_F) \circ \mu_{H(F)} = 1_{H(F)}$. Thus f factors through ${}_{eRe}(Re)^{(Y)}$ which is projective and finitely generated and hence f factors throughout a finite product of copies of eRe . This proves that ${}_{eRe}eF$ is flat.

Now we prove that ${}_R F$ is flat. Let J be a right ideal of R and assume that $\sum_{i=1}^n r_i m_i = 0$ for $m_1, m_2, \dots, m_n \in F$ and $r_1, r_2, \dots, r_n \in J$. Let e be an idempotent in I such that $em_i = m_i$ for every $i = 1, 2, \dots, n$. Then $\sum_{i=1}^n r_i \otimes m_i = \sum_{i=1}^n r_i \otimes em_i = \sum_{i=1}^n r_i e \otimes em_i$. Therefore $\sum_{i=1}^n r_i e \otimes em_i$ belongs to the kernel of the canonical $Je \otimes_{eRe} eF \rightarrow eRe$, which is injective. Therefore it is 0 in $Je \otimes_{eRe} eF$ and hence it is 0 in $J \otimes_R F$. ■

Lemma 5.4. *Let $\xi : X \rightarrow X'$ be a morphism of left G -sets and assume that G acts freely on X . For every $x' \in X'$, $S_\xi((x')R)$ is isomorphic to $\bigoplus_{\xi(x)=x'} (x)R$. Therefore S_ξ preserves projectivity.*

Proof: For every $x \in \xi^{-1}(x')$ set $f_x : (x)R \rightarrow S_\xi((x')R)$ given by: $f_x(r) = r^y$ if $r \in {}_y(x)R$. Note that if $r \in R_g$ and $gx = y$ then $gx' = \xi(y)$ and hence $r \in {}_{\xi(y)}(x')R$. If $r \in R_g$ and $s \in R_h$, then $f_x(rs) = (rs)^{g^h x} = r(s)^{hx} = r f_x(s)$. Thus f_x is a homomorphism of left R -modules and it is obviously graded. Therefore there is an homomorphism $f : \bigoplus_{\xi(x)=x'} (x)R \rightarrow S_\xi((x')R)$ such that f restricted to $(x)R$ is equal to f_x for every $x \in \xi^{-1}(x')$.

Now we show that f is bijective. Let $\sum_{x \in X} r^x \in \text{Ker}(f)$ where $r^x \in (x)R$. Since f is a graded homomorphism, then $\sum_{x \in X} {}_y(r^x) \in \text{Ker}(f)$ for any $y \in X$. If $g \in \text{Supp}({}_y(r^x)) \cap \text{Supp}({}_y(r^{x'}))$, then $gx = y = gx'$, therefore $x = x'$. Thus $0 = f(\sum_{x \in X} {}_y(r^x)) = (\sum_{x \in X} {}_y(r^x))^y$ and hence ${}_y(r^x) = 0$ for every $x, y \in X$. This proves that f is injective. On the other hand, if $r^y \in S_\xi((x')R)_y$, then $r \in {}_{\xi(y)}(x')R$. Therefore, $gx' = \xi(y)$ for every $g \in \text{Supp}(r)$. For every $g \in G$, let $x_g = g^{-1}y$. Then $r_g \in {}_{y(x_g)}R$ and $f_{x_g}(r_g) = (r_g)^y$, thus $f(\sum_{g \in G} r_g^{x_g}) = r^y$.

Now if $P \in (G, X', R) - gr$ is projective, then it is a direct summand of copies of modules of the form $(x')R$ for some $x' \in X'$. Since S^ξ preserves direct sums, $S^\xi(P)$ is isomorphic to a direct summand of a direct sum of modules of the form $(x)R$ for some $x \in X$. Therefore $S^\xi(P)$ is projective. ■

Let R be a G -graded ring. Let $P = \sum_{g \in G} R(g)$. For every $g \in G$ let

$$u_g : R(g) \longrightarrow P$$

the canonical monomorphism. Consider P as an $G \times G$ -graded R - R -bimodule as in Section 2. For every subgroup H of G , we will consider G as an G/H -set. We will use the following notation:

$$A_H = \text{END}(P_R)_H$$

$$J_H = \{\alpha \in A_H \mid \alpha u_x = 0 \text{ for almost all } g \in G\}$$

$$U_H = \bigoplus_{C \in G/H} R(C) \text{ considered as an } G/H \times G/H\text{-graded bimodule as}$$

in Section 2.

$$S_H = \text{End}_{gr-(R, G/H)}(U)$$

$$I_H = \{\alpha \in S_H \mid \alpha u_C = 0 \text{ for almost all } C \in G/H\}.$$

Lemma 5.5. J_H is projective as a left A_H -module.

Proof: Actually $J_H = \bigoplus_{g \in G} A_H P_g$. ■

Proof of Theorem 5.1:

First it is well known that $gr.w. \dim(R) \leq w. \dim(R)$ (see e.g. [NV1]).

Assume that $gr.w. \dim(R) \leq n$. Let H be a finitely generated subgroup of G and $M \in (G/H, R) - gr$. Let

$$\dots \longrightarrow P_2 \xrightarrow{f_2} P_1 \xrightarrow{f_1} P_0 \xrightarrow{f_0} M \longrightarrow 0$$

be a projective resolution in $(G/H, R) - gr$. Then by Lemma 5.4

$$\dots \longrightarrow S_H(P_2) \xrightarrow{S_H(f_2)} P_1 \xrightarrow{S_H(f_1)} P_0 \xrightarrow{S_H(f_0)} M \longrightarrow 0$$

is a projective resolution in $R - gr$. Therefore, $\text{Ker}(S_H(f_{n-1}))$ is flat (being $f_{-1} = 0 : M \rightarrow 0$). But $\text{Ker}(S_H(f_{n-1})) = S_H(\text{Ker}(f_{n-1}))$ and S_H is separable, therefore $\text{Ker}(f_{n-1})$ is flat. Thus for every $M \in (G/H, R) - gr$, $fd(M) \leq n$.

Let $B = \overline{\text{END}}(R_P)_H$. Then $J_H = \{\alpha \in B \mid u_g \alpha = 0 \text{ for almost all } g \in G\}$. By Theorem 3.4 $(G/H, R) - gr$ is equivalent to $\text{Gen}(S_H(I_H))$.

By Corollary 3.10, $\text{Gen}(B(J_H))$ is equivalent to $\text{Gen}(S_H(I_H))$. Finally, by Lemma 3.5 $\text{Gen}(B(J_H))$ is equivalent to $\text{Gen}(A_H(J_H))$. Thus $fd(M) \leq n$ for all $M \in \text{Gen}(A_H(J_H))$.

For every subgroup H of G , I_G is isomorphic to $(I_H)^{(G/H)}$ as a right A_H -module. Thus $\text{Gen}(A_H(I_G)_{A_H}) = \text{Gen}(A_H(I_H))$.

Let $M \in \text{Gen}(A_G(J_G))$ and

$$\dots \longrightarrow P_2 \xrightarrow{f_2} P_1 \xrightarrow{f_1} P_0 \xrightarrow{f_0} M \longrightarrow 0$$

a projective resolution of M such that $P_n \in \text{Gen}(A_G(J_G))$ for every $n \geq 0$. Note that such a projective resolution exists because $\text{Gen}(A_G(J_G))$ is closed under submodules and $A_G(J_G)$ is projective (see Lemma 5.5). Since $A_H(A_G)$ is projective and restriction of scalars sends modules in $\text{Gen}(A_G(J_G))$ to modules in $\text{Gen}(A_H(J_G))$, then ${}_{A_H} \text{Ker}(f_{n-1})$ is flat. But A_G is the direct limit of the A_H 's where H runs on the finitely generated subgroups of G . Therefore ${}_{A_H} \text{Ker}(f_{n-1})$ is flat and hence $fd(A_G M) \leq n$ for every $M \in \text{Gen}(A_G(J_G))$.

Now by using the equivalence $\text{Gen}(A_G(J_G)_{A_G}) \cong \text{Gen}(B_G J)$ of Lemma 3.5 one has that $fd(B_G M) \leq n$ for every $M \in \text{Gen}(B_G(J_G))$. And using the equivalence $\text{Gen}(B_G(J_G)) \cong \text{Gen}(S_G I_G) = R\text{-mod}$ of Corollary 3.10 for $G = X = H$ one has that $w. \dim(R) \leq n$. ■

References

[A] T. ALBU, Pure ideals, quotient categories and infinite group-graded rings, *Comm. in Algebra* **18(3)** (1990), 839–862.
 [AN] T. ALBU AND C. NĂSTĂSESCU, Infinite group-graded rings, rings of endomorphisms and localization, *J. of Pure and Appl. Algebra* **59** (1989), 125–150.
 [AM] P. N. ANH AND L. MÁRKI, Morita equivalence for rings without identity, *Tsukuba J. Math.* **11** (1987), 1–16.
 [AW] P. N. ANH AND R. WIEGANDT, On Grothendieck categories: Morita duality, Preprint.
 [B1] M. BEATTIE, A Generalization of the smash product of a graded ring, *J. of Pure and Appl. Algebra* **52** (1988), 219–226.
 [B2] M. BEATTIE, Duality theorems for rings with actions and coactions, *J. Algebra* **115** (1988), 302–312.
 [CM] M. COHEN AND S. MONTGOMERY, Group graded rings, smash product and group actions, *Trans. Am. Math. Soc.* **284** (1984), 237–258.

- [CF] R. R. COLBY AND K. R. FULLER, Exactness of the double dual and Morita duality for Grothendieck categories, *J. Algebra* **82** (1983), 546–558.
- [D1] E. C. DADE, Clifford Theory for group graded rings, *J. Reine Angew. Math.* **369** (1986), 40–86.
- [D2] E. C. DADE, Clifford Theory for group graded rings II, *J. Reine Angew. Math.* **387** (1988), 148–181.
- [F] K. R. FULLER, Density and equivalences, *J. of Algebra* **29** (1974), 528–550.
- [GG] J. L. GÓMEZ PARDO AND P. A. GUIL ASENSIO, Linear compactness and Morita duality for Grothendieck categories, to appear in *J. Algebra*.
- [GN] J. L. GÓMEZ PARDO AND C. NĂSTĂSESCU, Relative projectivity, graded Clifford theory and applications, *J. Algebra* **141** (1991), 484–504.
- [M] C. MENINI, Functors between categories of graded modules. Applications, Preprint.
- [MN] C. MENINI AND C. NĂSTĂSESCU, When are induced and coinduced functors equivalent?, Preprint.
- [MR] C. MENINI AND A. DEL RÍO, Morita duality and graded rings, *Comm. in Algebra* **19** (1991), 1765–1794.
- [NRV] C. NĂSTĂSESCU, S. RAIANU AND F. VAN OYSTAEYEN, Modules graded by G -sets, *Math. Z.* **203** (1990), 605–627.
- [NVV] C. NĂSTĂSESCU, M. VAN DER BERGH AND F. VAN OYSTAEYEN, Separable functor, applications to graded rings and modules, *J. Algebra* **123** (1989), 397–413.
- [NV1] C. NĂSTĂSESCU AND F. VAN OYSTAEYEN, “Graded ring theory,” North Holland Mathematical Library **28** (North Holland, Amsterdam, 1982).
- [NV2] C. NĂSTĂSESCU AND F. VAN OYSTAEYEN, Clifford Theory for subgroups of graded groups, Preprint.
- [Po] N. POPESCU, “Abelian categories with applications to rings and modules,” LMS Monographs **3**, Acad. Press, 1973.
- [Q] D. QUINN, Group-graded rings and duality, *Trans. A.M.S.* **292**.
- [Ra] M. D. RAFAEL, Separable functors revisited, *Comm. in Algebra* **18** (1990), 1445–1459.
- [R1] A. DEL RÍO, Weak dimension of group-graded rings, *Publicacions Matemàtiques* **34** (1990), 209–216.
- [R2] A. DEL RÍO, Graded rings and equivalences of categories, *Comm. in Algebra* **19** (1991), 997–1012.

- [S] B. STENSTRÖM, "*Ring of quotients*," Springer-Verlag, New York-Heidelberg-Berlin, 1975.
- [W] R. WISBAUER, Localization of modules and the central closure of rings, *Comm. in Algebra* **9** (1981), 1455–1493.

Departamento de Matemáticas
Universidad de Murcia
30071 Murcia
SPAIN

Rebut el 7 de Gener de 1992