

A NUMERICAL ALGORITHM FOR FINDING SOLUTIONS OF A GENERALIZED NASH EQUILIBRIUM PROBLEM

LUIZ CARLOS MATIOLI, WILFREDO SOSA, AND JINYUN YUAN

ABSTRACT. A family of nonempty closed convex sets is built by using the data of the Generalized Nash equilibrium problem (GNEP). The sets are selected iteratively such that the intersection of the selected sets contains solutions of the GNEP. The algorithm introduced by Iusem-Sosa (2003) is adapted to obtain solutions of the GNEP. Finally some numerical experiments are given to illustrate the numerical behavior of the algorithm.

1. INTRODUCTION

The standard definition of a non-cooperative game usually requires that each player in the game has a feasible set (or strategies set) independently on the rival's strategies. In a game with N players, for player i -th, we denote by $K_i \subset \mathbb{R}^{n_i}$ its feasible set and the function $h_i: \prod_i^N K_i \rightarrow \mathbb{R}$ will be called the loss function.

So, in this game, the feasible set of the game is $\prod_i^N K_i$. The goal of this game is

find \bar{x} in the feasible set of the game such that $h_i(y) \geq h_i(\bar{x}) \forall y \in \prod_i^N K_i$ and $\forall i \in \{1, \dots, N\}$. It was well understood from the early developments in the field (see [1, 8, 10]) that in many cases the interaction among the players can also take place at the feasible set of each player, in this case the feasible set for each player depend on the strategy of the other players, so the feasible set of the game can be different to the product of the feasible set of each player. We speak of Generalized Nash game, when the feasible set of each player depend on the strategies of the other players. Next we shall introduce some notations.

The index set of players is denoted by $I = \{1, 2, \dots, N\}$. For each $x \in \mathbb{R}^n$ and each $i \in I$, $x = [x_i]_{i \in I}$, where $x_i \in \mathbb{R}^{n_i}$ and $n = \sum_{i \in I} n_i$. Now, taking $\Lambda = \prod_{j \in I, j \neq i} \mathbb{R}^{n_j}$, we define $x_{-i} = P_\Lambda(x)$ where $P_C(x)$ denotes the orthogonal projection of x on the set C . Note that $x_i = P_{\mathbb{R}^{n_i}}(x)$. Now, for each $x \in \mathbb{R}^n$, each $i \in I$ and each

Key words and phrases. Convex feasibility Problem, Generalized Nash Equilibrium Problem, Projection methods.

The work was partially supported by the CNPq, CAPES, and Fundação Araucária, Brazil, CONCYTEC (projects STIC-AMSUD), Peru and Fundación Carolina and CRM, Spain.

$\rho \in \mathbb{R}^{n_i}$, we define $x(i, \rho) \in \mathbb{R}^n$ as $(x(i, \rho))_i = \rho$ and $(x(i, \rho))_{-i} = x_{-i}$. Note, that this notation is different to the classical notation, but we prefer it because we feel it more flexible than the classical one, from the mathematical view point.

For the Generalized Nash game considered here, we denote its feasible set by K and we consider K as a convex closed set. For each $x \in K$, and each $i \in I$, the set $K_i(x) = \{\rho \in \mathbb{R}^{n_i} : x(i, \rho) \in K\}$ will be the feasible set of the i -th player when the other players choose strategies x_{-i} . It is easy to see that

$$(1) \quad K = \{x \in \mathbb{R}^n : x_i \in K_i(x), i = 1, \dots, N\}.$$

For each player $i \in I$, we consider its loss function defined by $h_i: \Omega \rightarrow \mathbb{R}$, where Ω is an open set of \mathbb{R}^n and $K \subset \Omega$. Assume that h_i is continuously differentiable on Ω and $h_i(x(i, \cdot)): \mathbb{R}^{n_i} \rightarrow \mathbb{R} \cup \{+\infty\}$ pseudo-convex (in the sense of Karamardian [6]) for each $x \in K$ and each $i \in I$.

Using the above notation, we state the formal definition of the Generalized Nash Equilibrium Problems (GNEP in the sequel) as follows:

The GNEP is a Nash game, in which the feasible set of each player depends on the other player's strategies, which consists of finding a point in the feasible set of the game $\bar{x} \in K$ such that, for each $i \in I$, \bar{x}_i solves the minimization problem defined by:

$$(2) \quad \min h_i(\bar{x}(i, \rho)) \text{ subject to } \rho \in K_i(\bar{x}).$$

For recent study of the GNEP, Facchinei and Kanzow have given an excellent survey in [2].

The paper is organized as follows. Some basic results are given In Section 2. A numerical algorithm based on successive projection for solving the GNEP is introduced in Section 3, as a specialization of the other one introduced by Iusem and Sosa ([4]). The convergence of the algorithm is studied as well in the section 3. Compared with other methods, our method is simpler and more efficient. Our method does not require extras variables. Some numerical experiments and comments are given in the last section. In terms of numerical tests, our method can be applied to solve large scale problems.

2. PREVIOUS RESULTS

Given the GNEP, we define:

$$(3) \quad F: \Omega \rightarrow \mathbb{R}^n \text{ by } F(x) = [\nabla_i h_i(x(i, \cdot))(x_i)]_{i \in I}$$

$$(4) \quad f: \Omega \times \mathbb{R}^n \rightarrow \mathbb{R} \text{ by } f(x, y) = \langle F(x), x - y \rangle$$

for each $x \in K$:

$$(5) \quad L_f(x) = \{y \in K : f(x, y) \geq 0\}$$

Remark 2.1. We point out that $f(x, y) = \sum_{i \in I} \langle \nabla_i h_i(x(i, \cdot))(x_i), x_i - y_i \rangle \forall x, y \in K$. So, when $y = x(i, \rho)$ for $i \in I$ and $\rho \in K_i$, then $f(x, y) = \langle \nabla_i h_i(x(i, \cdot))(x_i), x_i - \rho \rangle$, because for each $j \neq i$, $y_j = x_j$ and so $x_j - y_j = 0$.

The next lemma follows directly from the previous definitions without proof.

Lemma 2.2. *The following statements hold.*

- (1) *The function f is continuous on $\Omega \times \mathbb{R}^n$.*
- (2) *For each $x \in \Omega$, $f(x, x) = 0$.*
- (3) *For each $x \in \Omega$, $f(x, \cdot): \mathbb{R}^n \rightarrow \mathbb{R}$ is linear affine (in particular is convex and concave).*
- (4) *For each $y \in \Omega$, $L_f(y)$ is nonempty closed and convex (in particular if K is a polyhedron, then $L_f(y)$ is also a polyhedron).*

Lemma 2.3. *If $x^* \in K$ is a local minimal point of the function $f(\cdot, x^*): K \rightarrow \mathbb{R}$, then x^* is a solution of GNEP.*

Proof. If $x^* \in K$ is local minimal point of the function $f(\cdot, x^*) : K \rightarrow \mathbb{R}$, then there exists a nonempty open convex set U such that $x^* \in U \cap K$ and $0 = f(x^*, x^*) \leq f(y, x^*)$ for each $y \in U \cap K$. Now, $\forall y \in K$ and $y \neq x^*$, there exists $\bar{t} \in]0, 1[$ such that $x_t = ty + (1 - t)x^* \in U \cap K$ for each $t \in]0, \bar{t}[$. But, $0 = f(x_t, x_t) = tf(x_t, y) + (1 - t)f(x_t, x^*)$ for each $t \in]0, \bar{t}[$. Thus, $f(x_t, y) \leq 0$ for each $t \in]0, \bar{t}[$. In the limit, we have that $f(x^*, y) \leq 0$. Now, for each $i \in I$, $\rho \in K_i(x^*)$ and $y = x^*(i, \rho)$, we have, from the previous remark, that $\langle \nabla_i h_i(x^*), \rho - x_i^* \rangle = -f(x^*, y) \geq 0$. Therefore, the statement follows from the pseudo-convexity of the functions h_i for each $i \in I$. \square

Lemma 2.4. *If there exists $x^* \in K$ such that $F(x^*) = 0$, then x^* is a solution of GNEP.*

Proof. Follows from the pseudo convexity of the functions h_i for each $i \in I$. \square

3. THE ALGORITHM

In this section, we introduce an algorithm, based on a successive projection scheme. Each iteration basically consists of two steps: an inexact local minimization of a continuous function over a compact set, and an orthogonal projection onto a set of the form $L_f(y)$ for some $y \in K$. The algorithm requires a constant $\alpha \in (0, 1)$ and three parameter sequences: relaxation parameters $\{\lambda_m\} \subset [\alpha, 2 - \alpha]$, precision parameters $\{\varepsilon_m > 0\} \downarrow 0$ for the inexact maximization, and $\{\delta_m > \alpha\} \uparrow \delta$ for the local minimization respectively. The algorithm generates two sequences $\{x^m\}, \{y^m\} \subset K$ in the following way.

Algorithm 3.1.

Initial step: Choose $x^0 \in K$.

Iterative step: Given $x^m \in K$,

a) Find $y^m \in K \cap \bar{B}(x^m, \delta_m)$ satisfying

$$(6) \quad f(y^m, x^m) \leq \min\{f(y, x^m) : y \in K \cap \bar{B}(x^m, \delta_m)\} + \varepsilon_m.$$

b) Compute $x^{m+1} \in K$ as

$$(7) \quad x^{m+1} = x^m + \lambda_m(P_{L_f(y^m)}(x^m) - x^m).$$

Next we shall study the properties of the sequences generated by Algorithm 3.1.

Lemma 3.1. *Given a convergent sequence $\{x^m\} \subset \mathbb{R}^n$ and $\{\delta_m > \alpha\} \uparrow \delta$. If x^* is the limit of the sequence $\{x^m\}$, then for each $\epsilon > 0$, there exists $M > 0$ such that for all $m \geq M$, the following statements hold.*

- (1) $\|x^m - x^*\| < \frac{\epsilon}{2}$.
- (2) $\delta_m > \delta - \frac{\epsilon}{2}$.
- (3) $B(x^*, \delta - \epsilon) \subset B(x^m, \delta_m) \subset B(x^*, \delta + \frac{\epsilon}{2})$.
- (4) For all $y \in B(x^*, \delta)$, there exists $i \geq m$ such that $\delta_i \geq \|y - x^i\|$

Proof. Since $\{x^m\}$ converges to x^* , the sequence $\{\|x^m - x^*\|\}$ converges to zero. Hence, there exists $M_1 > 0$ such that $\|x^m - x^*\| < \frac{\epsilon}{2}$ for each $m \geq M_1$. Also since $\{\delta_m\} \uparrow \delta = \sup\{\delta_m : m \in \mathbb{N}\}$, there exists $M_2 > 0$ such that $\delta_m > \delta - \frac{\epsilon}{2}$. Then, items (1) and (2) follows from taking $M = \max\{M_1, M_2\}$. Now, for $y \in B(x^*, \delta - \epsilon)$, $\|y - x^m\| \leq \|y - x^*\| + \|x^* - x^m\| \leq \delta - \epsilon + \frac{\epsilon}{2} = \delta - \frac{\epsilon}{2} < \delta_m$. The other inclusions are analogous. Thus item (3) follows. For item (4), suppose that there exists $m \geq M$ and $y \in B(x^*, \delta)$ such that $\delta_i < \|y - x^i\|$ for all $i \geq m$. Taking the limit, we obtain $\delta \leq \|y - x^*\|$ which is a contradiction because $y \in B(x^*, \delta)$. The proof is complete. \square

Now, we start the convergence analysis of the algorithm

Proposition 3.2. *Algorithm 3.1 is well defined.*

Proof. Since $x^m \in K$ for all m , all intersections $K \cap \bar{B}(x^m, \delta_m)$ are nonempty and trivially compact. It follows from the continuity of f that $f(\cdot, x^m)$ attains its minimum over $K \cap \bar{B}(x^m, \delta_m)$. Hence, there exists y^m satisfying (6). Since, $L_f(y^m)$ is closed and convex, x^{m+1} is uniquely defined by (7). \square

Lemma 3.3. *Let $\{x^m\}$ and $\{y^m\}$ be the sequences generated by Algorithm 3.1. If there exists $M > 0$ such that $\bigcap_{m \geq M} L_f(y^m)$ is nonempty, then*

- (1) The sequence $\{\|x^m - \bar{x}\|\}$ is convergent, for each $\bar{x} \in \bigcap_{m \geq M} L_f(y^m)$.
- (2) The sequence $\{x^m - P_{L_f(y^m)}(x^m)\}$ converges to zero.
- (3) The sequences $\{x^m\}$ and $\{y^m\}$ are bounded.

Proof. (1) Set $\bar{x} \in \bigcap_{m \geq M} L_f(y^m)$, by (7):

$$\begin{aligned} \|x^{m+1} - \bar{x}\|^2 &= \|x^m - \bar{x}\|^2 + \lambda_m^2 \|x^m - P_{L_f(y^m)}(x^m)\|^2 \\ &\quad + 2\lambda_m \langle x^m - \bar{x}, P_{L_f(y^m)}(x^m) - x^m \rangle \\ &= \|x^m - \bar{x}\|^2 + \lambda_m^2 \|x^m - P_{L_f(y^m)}(x^m)\|^2 - 2\lambda_m \|x^m - P_{L_f(y^m)}(x^m)\|^2 \\ &\quad + 2\lambda_m \langle P_{L_f(y^m)}(x^m) - \bar{x}, P_{L_f(y^m)}(x^m) - x^m \rangle. \end{aligned}$$

Since \bar{x} belongs to $L_f(y^m)$ for each $m \geq M$, the last term in the previous expression is nonnegative, because of the following well known property of orthogonal projections: $\langle P_C(x) - y, P_C(x) - x \rangle \leq 0$ for any closed and convex set C , any $x \in \mathbb{R}^n$ and any $y \in C$. Thus, it follows that, for all $m \geq M$,

$$(8) \quad \|x^{m+1} - \bar{x}\|^2 \leq \|x^m - \bar{x}\|^2 - \lambda_m(2 - \lambda_m)\|x^m - P_{L_f(y^m)}\|^2 \leq \|x^m - \bar{x}\|^2,$$

by the fact that $\lambda_m(2 - \lambda_m) > 0$. We conclude from (8) that the sequence $\{\|x^m - \bar{x}\|\}_{m \geq M}$ is non-increasing, and henceforth, being nonnegative, so $\{\|x^m - \bar{x}\|\}_{m=0}^\infty$ is convergent.

(2) Rewriting (8) and using the conditions on $\{\lambda_m\}$ we get

$$\begin{aligned} \alpha(2 - \alpha)\|x^m - P_{L_f(y^m)}(x^m)\|^2 &\leq \lambda_m(2 - \lambda_m)\|x^m - P_{L_f(y^m)}(x^m)\|^2 \\ &\leq \|x^m - \bar{x}\|^2 - \|x^{m+1} - \bar{x}\|^2. \end{aligned}$$

using the convergence of $\{\|x^m - \bar{x}\|\}$ we obtain, since $\alpha(2 - \alpha) > 0$,

$$(9) \quad \lim_{m \rightarrow \infty} (x^m - P_{L_f(y^m)}(x^m)) = 0.$$

(3) The convergence of $\{\|x^m - \bar{x}\|\}$ also implies that the sequence $\{x^m\}$ is bounded. For each m , since $y^m \in K \cap \bar{B}(x^m, \delta_m)$, $\|y^m\| - \|x^m\| \leq \|y^m - x^m\| \leq \delta_m$. Thus, $\|y^m\| \leq \|x^m\| + \delta_m$. The statement follows from the facts that $\{\|x^m\|\}$ and $\{\delta_m\}$ are bounded. \square

Let $\{x^m\}$ be sequence generated by Algorithm 3.1. Remember that x^* is a cluster point of $\{x^m\}$ if there is a subsequence $\{x^{m_i}\}$ of $\{x^m\}$ such that $\{x^{m_i}\}$ converges to x^* .

Theorem 3.4. *Let $\{x^m\}$ and $\{y^m\}$ be the sequences generated by Algorithm 3.1 and let $M > 0$.*

- i) *If $\bigcap_{m > M} L_f(y^m)$ is nonempty, then any cluster point of $\{x^m\}$ is a local minimizer of $f(\cdot, x^*)$ over K .*
- ii) *If $\{x^m\}$ converges to $x^* \in K$, then x^* is local minimizer of $f(\cdot, x^*)$ over K .*
- iii) *If GNEP lacks solutions, then $\{x^m\}$ is not convergent (though it might be bounded).*

Proof. i) From Lemma 3.3, we know that $\{x^m\}$ and $\{y^m\}$ are bounded. Let x^* be a cluster point of $\{x^m\} \subset K$. Since K is closed, x^* belongs to K . Thus, we can select a subsequence $\{x^{k_m}\}$ of $\{x^m\}$ such that $\lim_{m \rightarrow \infty} x^{k_m} = x^*$ and $\{y^{k_m}\}$ converges to y^* . By (9), $\lim_{m \rightarrow \infty} P_{L_f(y^{k_m})}(x^{k_m}) = x^*$. It follows from the continuity of f that

$$(10) \quad f(y^*, x^*) = \lim_{m \rightarrow \infty} (f(y^{k_m}, P_{L_f(y^{k_m})}(x^{k_m}))) \geq 0,$$

where the inequality follows from the definition of $L_f(y^{k_m})$ and the fact that

$$P_{L_f(y^{k_m})}(x^{k_m}) \in L_f(y^{k_m}).$$

From Item (3) of Lemma 3.1, for any fixed $\epsilon > 0$, there exists $M_\epsilon > 0$ such that $B(x^*, \delta - \epsilon) \subset B(x^{k_m}, \delta_{k_m})$ for all $k_m > M_\epsilon$. Taking $\bar{M} = \max\{M, M_\epsilon\}$, for each $y \in B(x^*, \delta - \epsilon)$ and each $k_m > \bar{M}$, we obtain from equation (6) of Algorithm 3.1 that $f(y^{k_m}, x^{k_m}) \leq f(y, x^{k_m}) + \epsilon_{k_m}$. Hence, in the limit, $f(x^*, x^*) = 0 \leq f(y^*, x^*) \leq f(y, x^*)$ which implies that x^* is a local minimum of $f(\cdot, x^*) : K \rightarrow \mathbb{R}$.

- ii) If $\{x^m\}$ converges to x^* , then $\{x^m\}$ and $\{y^m\}$ are bounded and $\lim_{m \rightarrow \infty} P_{L_f(y^{k_m})}(x^{k_m}) = x^*$. Thus, the statement follows from the same argument as in the previous item.
- iii) Suppose that $\{x^m\}$ converges to x^* . From the previous item, x^* is a local minimum of $f(\cdot, x^*)$. From Lemma 2.3 x^* is a solution of the GNEP which contradicts to the hypothesis of this item. \square

Next we shall consider the Convex Feasibility Problem (CFP in short)

$$\bigcap_{y \in K} L_f(y),$$

which is associated to GNEP.

Corollary 3.5. *If the CFP has solutions, then every cluster point of the sequence $\{x^m\}$ generated by Algorithm 3.1 is a solution of the GNEP.*

Proof. If the CFP has solutions, then $\emptyset \neq \bigcap_{y \in K} L_f(y) \subset \bigcap_{m=1}^{\infty} L_f(y^m)$. The result follows from Theorem 3.4(i) immediately. \square

In the next result we consider the notion of pseudomonotonicity (for details see for instance [6], [5]) for the operator F defined in 3, i.e. F is pseudomonotone if for $x, y \in \text{dom}(T)$: $\langle F(x), y - x \rangle \geq 0$ implies $\langle F(y), x - y \rangle \leq 0$. Note that the previous implication is equivalent to say that $\langle F(x), y - x \rangle > 0$ implies $\langle F(y), x - y \rangle < 0$.

Lemma 3.6. *If F is pseudomonotone, then $\{L_f(y)\}_{y \in K}$ is such that for each $\{x^1, \dots, x^p\} \subset K$ we have that $\text{co}\{x^1, \dots, x^p\} \subset \bigcup_{i=1}^p L_f(x^i)$*

Proof. Suppose that $\exists \bar{x} \in \text{co}\{x^1, \dots, x^p\}$ such that $\bar{x} \notin \bigcup_{i=1}^p L_f(x^i)$, then $\bar{x} = \sum_{i=1}^p \lambda_i x^i$ with $\lambda_i \geq 0$ and $\sum_{i=1}^p \lambda_i = 1$ and $\langle F(x^i), x^i - \bar{x} \rangle < 0$, $\forall i$, then $\langle F(x^i), \bar{x} - x^i \rangle > 0$, $\forall i$. So, from the pseudomonotonicity of F , we have that $\langle F(\bar{x}), x^i - \bar{x} \rangle < 0$, $\forall i$. So, $0 = \langle F(\bar{x}), \bar{x} - \bar{x} \rangle = \sum_{i=1}^p \lambda_i \langle F(\bar{x}), x^i - \bar{x} \rangle < 0$. The statement follows from this contradiction. \square

Corollary 3.7. *Let $\{x^m\}$ and $\{y^m\}$ be the sequences generated by Algorithm 3.1. If F is pseudo-monotone and either $\{x^m\}$ or $\{y^m\}$ is bounded, then every cluster point of the sequence $\{x^m\}$ is a solution of the GNEP.*

Proof. If $\{x^m\}$ is bounded, then $\{y^m\}$ is bounded by the same argument as in the proof of (i) of Theorem 3.4. Thus we may assume that $\{y^m\}$ is bounded. By the previous Lemma and Lemma 2.3 in [4] $\cap_{m=1}^{\infty} L_f(y^m) \neq \emptyset$. The statement follows from (i) of Theorem 3.4. \square

4. NUMERICAL EXPERIMENTS

First of all we set $K = \{x \in \mathbb{R}^n : \langle a^i, x \rangle \leq b_i, x_j \geq 0, i = 1, \dots, m, j = 1, \dots, n\}$ as a polyhedron

In (6) we minimize a linear approximation of f in a compact set, that is, we solve the following subproblem: (given x^m in K and δ^m)

$$(11) \quad \begin{aligned} & \text{minimize} && \langle F(x^m), y - x^m \rangle \\ & \text{s.t.} && y \in \bar{B}(x^m, \delta^m) \cap K \end{aligned}$$

to find y^m . Note that in our implementation the region $\bar{B}(x^m, \delta^m)$ is a box.

In step (b) of Algorithm 3.1, we determine x^{m+1} by an orthogonal projection of x^m onto $L_f(y^m)$, which is equivalent to solving the following subproblem:

$$(12) \quad \begin{aligned} & \text{minimize} && \frac{1}{2} \|z - x^k\|^2 \\ & \text{s.t.} && z \in K \cap \{y : \langle F(x^m), y \rangle \leq \langle F(x^m), x^m \rangle\} \end{aligned}$$

The subproblems (11) and (12) are solved, respectively, by routines *linprog* and *lsqlin* from optimization toolbox of Matlab 6.0.0.88 (R12). The stop criteria is $\frac{\|x^{m+1} - x^m\|}{\|x^{m+1}\|} < 10^{-10}$.

The initial point in K is determined by solving

$$\min_{s,x} \sum_{i=1}^m s_i$$

subject to $Ax + s = b, s, x \geq 0$.

We consider only nonnegative x because negative variables have no real application meaning for most practical problems. In the implementation of Algorithm 3.1 we initialize $\alpha = 0.2$ in Problem 2, and $\alpha = 0.5$ in all other problems. The parameter δ_m controls the radio of box $\bar{B}(x^m, \delta^m)$ and is increased till δ_{max} . The updating form of this parameter is given by $\delta_0 = (\alpha + \delta_{max})/2$ and $\delta_{m+1} = (\delta_m + \delta_{max})/2$. All computations are executed by Matlab 6.0.0.88 on Laptop TOSHIBA Satellite M115-S3144 with intel Core 2 with 2 GB de RAM Memory.

Problem 1: $F(x_1, x_2) = (3x_1^2x_2^2, 3x_1^2x_2^2)$ and $K = \{(x_1, x_2) : x_1 + x_2 \leq 3, x_1 \geq 1; x_2 \geq 1\}$. The exact solution is $x = [1 \ 1]^T$. The numerical results are showed in Table 1, whose first column is the number of iterations, the second and third columns are the decision of players.

Problem 2: This problem is taken from [3] where authors transform the Nash equilibrium problem to Variational Inequality. In sense of Variational Inequality the solution is unique $\bar{x} = (\frac{3}{4}, \frac{1}{4})$ which belongs to solution set of the Nash equilibrium problem. Here the problem is given by: $F(x_1, x_2) = (2x_1 - 2, 2x_2 - 1)$ and $K = \{(x_1, x_2) : x_1 + x_2 \leq 1, x_1 \geq 0, x_2 \geq 0\}$.

m	x_1^m	x_2^m
0	1.3618	1.3618
1	1.0618	1.0618
2	1.0000	1.0000

TABLE 1. Numerical results for Problem 1

Problem 3: We consider the River Basin Pollution problem [7] with three players $j = 1, 2, 3$. The problem is given by $h_j(x) = p_j x_j^2 + 0.01 x_j (x_1 + x_2 + x_3) - q_j x_j$. Now using $F(x) = [\nabla_j h_j(x(j, \cdot))(x_j)]$, $j = 1, 2, 3$, we have

$$F(x) = \begin{bmatrix} 2p_1 x_1 + 0.01(2x_1 + x_2 + x_3) - q_1 \\ 2p_2 x_2 + 0.01(x_1 + 2x_2 + x_3) - q_2 \\ 2p_3 x_3 + 0.01(x_1 + x_2 + 2x_3) - q_3 \end{bmatrix}$$

where $p = [0.01; 0.05; 0.01]^T$ and $q = [2.9; 2.88; 2.85]^T$. The polyhedron K is given by constraints: $g_1(x) = 3.25x_1 + 1.25x_2 + 4.125x_3 \leq 100$, $g_2(x) = 2.2915x_1 + 1.5625x_2 + 2.8125x_3 \leq 100$, and $x_1, x_2, x_3 \geq 0$.

Problema 4[9]: The authors solve the problem by means of Linear Complementary Problem whose solution depends on multipliers. The problem is defined by $F(x) = (F_1(x), F_2(x))^T$ where $F_1(x) = -1 + x_1 + 0.5x_2$ and $F_2(x) = -2 + 0.5x_1 + x_2$. Here the polyhedron is given by $x_1 + x_2 \leq 1$, $x_1, x_2 \geq 0$. The solution founded by our algorithm is $\bar{x} = (x_1, x_2) = (0, 1)$.

Problem 5 [7]: The problem is give by

$$F(x) = \left[\frac{2(x_1 + x_2)}{4} + \frac{2(x_1 - x_2)}{9}; \frac{2(x_1 + x_2)}{4} - \frac{2(x_1 - x_2)}{9} \right].$$

Here $K = \{0 \leq x_1, x_2 \leq 10\}$.

Problems 6–10. We define the problems by

$$F_j(x) = 2p_j x_j + \alpha x_j + \alpha \sum_{k=1}^n x_k - q_j$$

where $F_j(x)$ is the j -th component of $F(x)$, the constants p , α as well as the m constraints are generated by function *rand* of Matlab with corresponding dimensions given in Table 2. The purpose of these problems is to test the numerical behavior of Algorithm 3.1 for large scale problems.

The numerical results and comparisons are given in Table 2 where the first column is the problem, second column is the number of variable, third column is the number of constraints, fourth column is the parameter α , fifth column is the δ_{max} , sixth column is number of iteration, seventh column is the number of the iterations of the method given in the cited references. Here – means that there is no number of iteration for other methods.

Problem	n	m	α	δ_{max}	nit	IT of other methods
Prob2	2	1	0.2	0.3	2	–[3]
Prob3	2	2	0.5	1.3	9	7[7]
Prob4	2	1	0.5	1.3	2	–[9]
Prob5	3	2	0.5	1.5	15	19[7]
Prob6	50	30	0.5	1	3	–
Prob7	100	80	0.5	1	4	–
Prob8	200	120	0.5	1	3	–
Prob9	500	50	0.5	1	5	–
Prob10	1000	30	0.5	1	2	–

TABLE 2. Numerical results and comparisons

The main advantages of Algorithm 3.1 compared with other methods are as follows.

- (1) Algorithm 3.1 doesn't require extra variables unlike other methods given in [7, 9].
- (2) By our numerical results, it seems that the convergence of Algorithm 3.1 is independent on number of variables which will be studied in the future.
- (3) The convergence rate is very competitive from our numerical experiments.
- (4) So far Algorithm 3.1 is the best compared with all existing methods for the problem.

We shall give further study and modifications of Algorithm 3.1 in the next paper.

ACKNOWLEDGMENT

This work was started during the visit of the second author at Department of Mathematics, Federal University of Paraná in 2008 and was finished during the

visit of the same author at Centre de Recerca Matemàtica (CRM) in 2009. The second author likes to thank Prof. Carlos Henrique dos Santos, Prof. Jinyun Yuan and Prof. Juan Enrique Martinez Legaz for their kindest hospitality during his visiting at: Department of Mathematics, Federal University of Paraná, Curitiba, Brazil and the CRM of Barcelona, Spain; respectively.

REFERENCES

- [1] K.J. Arrow, G. Debreu. Existence of an equilibrium for a competitive economy, *Econometrica* **22** (1954), 265–290.
- [2] F. Facchinei, C. Kanzow. Generalized Nash equilibrium problems *4OR* **5** (2007), 173–210.
- [3] F. Facchinei, A. Fischer, V. Piccialli. On generalized Nash games and variational inequalities *Operations Research Letters* **35** (2007), 159–164.
- [4] A.N. Iusem, W. Sosa. Iterative algorithms for equilibrium problems, *Optimization* **52** (2003,) 301–316.
- [5] A.N. Iusem, W. Sosa. New existence results for equilibrium problems, *Nonlinear Analysis* **52** (2003), 621–635.
- [6] S. Karamardian, Complementarity problems over cones with monotone and pseudomonotone maps, *J. Optimization Theory Appl.*, **18(4)** (1976), 445–454.
- [7] J.B. Krawczyk, S. Uryasev. relaxations algorithms to find Nash equilibria with economic applications, *Environmental Modeling and Assessment* **5** (2000), 63–73.
- [8] L.W. McKenzie. On the existence of a general equilibrium for a competitive market, *Econometrica* **27** (1959), 54–71.
- [9] J.S Pang, B.F. Hobbs. Nash-Cournot equilibria in electric power markets with piecewise linear demand functions and joint constraints *Operations Research* **55**, issue 1 (january 2007) 113-127.
- [10] J.B. Rosen. Existence and uniqueness of equilibrium points for concave n -person games, *Econometrica* **33** (1965), 520–534.

LUIZ CARLOS MATIOLI
 DEPARTAMENTO DE MATEMÁTICA - UFPR CENTRO POLITÉNICO
 CP: 19.081, CURITIBA
 PR 81531-980, BRAZIL
E-mail address: `matioli@mat.ufpr.br`

WILFREDO SOSA
 INSTITUTO DE MATEMÁTICA Y CIENCIAS AFINES
 UNIVERSIDAD NACIONAL DE INGENIERIA
 CALLE LOS BIOLOGOS 245, LIMA 12
 LIMA PERÚ
E-mail address: `sosa@uni.edu.pe`

JINYUN YUAN
 DEPARTAMENTO DE MATEMÁTICA - UFPR, CENTRO POLITÉNICO
 CP: 19.081, CURITIBA
 PR, 81531-980, BRAZIL
E-mail address: `jin@mat.ufpr.br`