

## SPACES WITH NOETHERIAN COHOMOLOGY

NATÀLIA CASTELLANA\*, VINCENT FRANJOU\*\*, ALAIN JEANNERET,  
AND JÉRÔME SCHERER\*\*\*

ABSTRACT. Is the cohomology of the classifying space of a  $p$ -compact group, with Noetherian twisted coefficients, a Noetherian module? This note provides, over the ring of  $p$ -adic integers, such a generalization to  $p$ -compact groups of the Evens-Venkov Theorem. We consider the cohomology of a space with coefficients in a module, and we compare Noetherianity over the field with  $p$  elements, with Noetherianity over the  $p$ -adic integers, in the case when the fundamental group is a finite  $p$ -group.

### INTRODUCTION

The main theorem of Dwyer and Wilkerson in [12] states that the mod  $p$  cohomology of the classifying space of a  $p$ -compact group is a finitely generated algebra. This generalizes to  $p$ -compact groups the Evens-Venkov Theorem [13] on the cohomology of a finite group  $G$ . There are however two main differences between these two results. Evens' statements allow a general base ring — any Noetherian ring is allowed, and they include the case of general twisted coefficients (contrary to the early work by Golod, [15], or Venkov, [24]) as follows: if  $M$  is Noetherian over a ring  $R$ , then so is  $H^*(G; M)$  over  $H^*(G; R)$ . Beautiful finite

---

2000 *Mathematics Subject Classification*. Primary 55U20; Secondary 13E05, 18G15, 55N25, 55T10, 55R12, 55R35.

\*The first author has been partially supported by the FEDER/MEC grant MTM2010-20692.

\*\*The second author has been partially supported by the LMJL - Laboratoire de Mathématiques Jean-Leray UMR 6629 CNRS/Université de Nantes, by the ANR grant HGRT BLAN08-2-338236, and by CRM Barcelona.

\*\*\*The fourth author has been partially supported by FEDER/MEC grant MTM2010-20692, the IHES, and by the University of Bern.

generation statements on cohomology have since been proved in numerous situations. For statements as general as Evens' however, proofs have been surprisingly elusive.

This note is concerned with these generalizations for  $p$ -compact groups and  $p$ -local finite groups, as defined by Broto, Levi, and Oliver, [8]. We ask more generally when Noetherianity of the mod  $p$  cohomology algebra  $H^*(Y; \mathbb{F}_p)$  of a space  $Y$  implies that the cohomology with coefficients in a  $R[\pi_1(Y)]$ -module  $M$ ,  $H^*(Y; M)$ , is a Noetherian module over the algebra  $H^*(Y; R)$ . Because the classifying space  $BX$  of a  $p$ -compact group is  $p$ -complete by definition, we work over  $p$ -complete rings (for example  $H^*((BS^3)_p^\wedge; \mathbb{Z})$  is not Noetherian).

**Theorem 2.4.** *Let  $Y$  be a connected space. Then, the graded  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(Y; \mathbb{Z}_p^\wedge)$  is Noetherian if, and only if, the graded  $\mathbb{F}_p$ -algebra  $H^*(Y; \mathbb{F}_p)$  is Noetherian and the torsion in  $H^*(Y; \mathbb{Z}_p^\wedge)$  is bounded.*

**Theorem 3.6.** *Let  $Y$  be a connected space such that  $\pi_1 Y$  is a finite  $p$ -group. Let  $M$  be a  $\mathbb{Z}_p^\wedge[\pi_1 Y]$ -module, which is finitely generated over  $\mathbb{Z}_p^\wedge$ . If the graded  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(Y; \mathbb{Z}_p^\wedge)$  is Noetherian, then  $H^*(Y; M)$  is Noetherian as a module over  $H^*(Y; \mathbb{Z}_p^\wedge)$ .*

This applies to  $p$ -compact group and to  $p$ -local finite groups to show that their  $p$ -adic cohomology algebra is Noetherian, see Theorem 4.2 and 4.5. Note that our proof makes no use of the recent classification of  $p$ -compact groups by Andersen, Grodal, Møller, and Viruel, [2], [3], [21]. Even in the case of a compact Lie group  $G$ , our theorem provides a general finiteness theorem for the cohomology of  $BG$  with twisted coefficients. One of the few explicit computations available in the literature is the case of  $O(n)$ , due to Čadek [10], (see also Greenblatt's [16]).

**Acknowledgements.** This work started when the second author visited the CRM in Barcelona during the emphasis semester on higher categories in 2008 and was continued while the fourth author was visiting the IHES and the University of Bern in 2009. We would like to thank these institutions for their generous hospitality. Our warmest thanks go to the anonymous referee for his careful

reading, improvements, and corrections, which led us to the present version of Theorem 2.4.

## 1. THE COHOMOLOGY AS A GRADED MODULE

Before considering the mod  $p$  or  $p$ -adic cohomology as an algebra, we first make explicit the relationship between two standard milder finiteness assumptions. When the graded vector space  $H^*(Y; \mathbb{F}_p)$  is of finite type, i.e.  $H^n(Y; \mathbb{F}_p)$  is a finite dimensional vector space in each degree  $n$ , is  $H^n(Y; \mathbb{Z}_p^\wedge)$  a finitely generated  $\mathbb{Z}_p^\wedge$ -module in each degree  $n$  as well? This is clearly a necessary condition for the cohomology algebra to be finitely generated. We show that it holds, even for non-simply-connected spaces.

The main tool to relate the mod  $p$  and the  $p$ -adic cohomology is the universal coefficient exact sequence — see for example [23, Theorem 5.5.10]:

$$(1) \quad 0 \rightarrow H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p \xrightarrow{\rho} H^*(Y; \mathbb{F}_p) \xrightarrow{\partial} \mathrm{Tor}(H^{*+1}(Y; \mathbb{Z}_p^\wedge); \mathbb{Z}/p) \rightarrow 0$$

which applies, since  $\mathbb{Z}_p^\wedge$  is a PID and  $\mathbb{Z}/p$  is a finitely generated  $\mathbb{Z}_p^\wedge$ -module.

**Remark 1.1.** The morphism  $\rho$  in (1) is a ring homomorphism which makes the middle term  $H^*(Y; \mathbb{F}_p)$  an  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ -module. Evens observed in [14, p. 272] that  $\partial$  is also a homomorphism of  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ -modules, where  $\mathrm{Tor}(H^*(Y; \mathbb{Z}_p^\wedge); \mathbb{Z}/p)$  has the natural module structure he introduced in [14, Lemma 2].

**Proposition 1.2.** *Let  $Y$  be a connected space with finite fundamental group. The group  $H^n(Y; \mathbb{F}_p)$  is finite for every positive integer  $n$  if and only if the  $\mathbb{Z}_p^\wedge$ -module  $H^n(Y; \mathbb{Z}_p^\wedge)$  is finitely generated for every  $n$ . Under this condition, the  $\mathbb{Z}_p^\wedge$ -module  $H^n(Y; M)$  is finitely generated for any  $n$  and every  $\mathbb{Z}_p^\wedge[\pi_1 Y]$ -module  $M$  which is finitely generated over  $\mathbb{Z}_p^\wedge$ .*

*Proof.* If  $H^n(Y; \mathbb{Z}_p^\wedge)$  is a finitely generated  $\mathbb{Z}_p^\wedge$ -module for any  $n$ , the universal coefficient exact sequence (1) implies that  $H^n(Y; \mathbb{F}_p)$  is finite for any  $n$ .

Conversely, assume that  $H^n(Y; \mathbb{F}_p)$  is finite for every  $n$ . Since the fundamental group of  $Y$  is finite, the space  $Y$  is  $p$ -good by [7, Proposition VII.5.1]

and therefore  $H^n(Y_p^\wedge; \mathbb{F}_p) \cong H^n(Y; \mathbb{F}_p)$ . Likewise, since cohomology with  $p$ -adic coefficients is represented by Eilenberg-MacLane spaces  $K(\mathbb{Z}_p^\wedge, n)$ , which are  $p$ -complete,  $H^n(Y_p^\wedge; \mathbb{Z}_p^\wedge) \cong H^n(Y; \mathbb{Z}_p^\wedge)$ , [7, Proposition II.2.8]. We may therefore assume that  $Y$  is  $p$ -complete and that  $G = \pi_1 Y$  is a finite  $p$ -group, see [12, Proposition 11.14] or [6, Section 5].

If  $Y$  is 1-connected, then [1, Proposition 5.7] applies and  $H^n(Y; \mathbb{Z}_p^\wedge)$  is a finitely generated  $\mathbb{Z}_p^\wedge$ -module for every  $n$ . For the general situation, let us consider the universal cover fibration  $\tilde{Y} \rightarrow Y \rightarrow BG$  where  $G$  is the fundamental group of  $Y$ . We prove by induction that  $H^n(\tilde{Y}; \mathbb{F}_p)$  is finite dimensional for any  $n$ . The induction starts with the trivial case  $n = 0$ . Assume thus that  $H^m(\tilde{Y}; \mathbb{F}_p)$  is finite for all  $m < n$ . Then, in the second page of the Serre spectral sequence in mod  $p$  cohomology, all groups  $E_2^{i,j} = H^i(BG, H^j(\tilde{Y}; \mathbb{F}_p))$  on the lines  $j = 0, \dots, n-1$  are finite. As  $E_\infty^{0,n}$  is finite as well, it follows that  $E_2^{0,n} = H^n(\tilde{Y}; \mathbb{F}_p)^G$  is finite dimensional. Let  $M$  be a non-trivial finite dimensional  $\mathbb{F}_p[G]$ -module. Since  $G$  is a finite  $p$ -group, the fixed point  $M^G$  is non-trivial, and finiteness of  $H^n(\tilde{Y}; \mathbb{F}_p)^G$  implies finiteness of  $H^n(\tilde{Y}; \mathbb{F}_p)$ .

We can now apply the 1-connected case to conclude that  $H^n(\tilde{Y}; \mathbb{Z}_p^\wedge)$  is a finitely generated  $\mathbb{Z}_p^\wedge$ -module for any  $n$ . The Evens-Venkov Theorem [13, Theorem 8.1] now shows that the  $E_2$ -term of the Serre spectral sequence with  $p$ -adic coefficients consists of finitely generated  $\mathbb{Z}_p^\wedge$ -modules. Thus so must be  $H^n(Y; \mathbb{Z}_p^\wedge)$  for any  $n$ .

The second part of the assertion follows by a similar argument. The first part of the proposition and the universal coefficient formula imply that  $H^n(\tilde{Y}; M)$  is a finitely generated  $\mathbb{Z}_p^\wedge$ -module for every  $n$ . We then use the Serre spectral sequence for cohomology with twisted coefficients. The only reference we know is [20, Theorem 3.2] where it is done equivariantly; we need the case of the trivial group action. ▮

## 2. COHOMOLOGY WITH TRIVIAL COEFFICIENTS

We now turn to finite generation of the cohomology algebras  $H^*(Y; \mathbb{Z}_p^\wedge)$  and  $H^*(Y; \mathbb{F}_p)$ , where trivial coefficients are understood. This section should thus

be no more than a warm up, because it seems enough to gain some control on torsion to draw conclusion from the universal coefficient theorem.

Let  $R$  be either the ring  $\mathbb{Z}_p^\wedge$ , or the field  $\mathbb{F}_p$ , and note that both are Noetherian rings. The cohomology  $H^*(Y; R)$  of any connected space is a commutative graded algebra, which is a Noetherian  $R$ -algebra if and only if it is finitely generated as an  $R$ -algebra [18, Theorem 13.1].

**Lemma 2.1.** *Let  $Y$  be a connected space. If the  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(Y; \mathbb{Z}_p^\wedge)$  is Noetherian, then  $H^*(Y; \mathbb{F}_p)$  is a finitely generated module over the algebra  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ .*

*Proof.* The ideal  $\text{Tor}(H^*(Y; \mathbb{Z}_p^\wedge); \mathbb{Z}/p)$  of elements annihilated by  $p$  is a finitely generated ideal of  $H^*(Y; \mathbb{Z}_p^\wedge)$  by assumption. It is therefore also finitely generated as an  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ -module. The conclusion follows from Remark 1.1 on the universal coefficients exact sequence.  $\blacksquare$

To be able to compare Noetherianity of the mod  $p$  and the  $p$ -adic cohomology, we need to analyze the  $p$ -torsion in  $H^*(Y; \mathbb{Z}_p^\wedge)$ . Let us denote by  $T_p H^*(Y; \mathbb{Z}_p^\wedge)$  the graded submodule of  $p$ -torsion elements. The key assumption in the main theorem of the section is that the order of the  $p$ -torsion is bounded. This implies that  $\rho$  is “uniformly power surjective”, a strong form of integrity.

**Lemma 2.2.** *Let  $Y$  be a connected space and let  $d$  be an integer such that  $p^d \cdot T_p H^*(Y; \mathbb{Z}_p^\wedge) = 0$ . Assume that  $H^n(Y; \mathbb{Z}_p^\wedge)$  is a finitely generated  $\mathbb{Z}_p^\wedge$ -module for any  $n$ . If  $u \in H^*(Y; \mathbb{F}_p)$ , then  $u^{p^d}$  belongs to the image of  $\rho : H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p \rightarrow H^*(Y; \mathbb{F}_p)$ .*

*Proof.* Following the elementary proof of [5, Lemma 4.5], we start with the observation that for any element  $x \in H^*(Y; \mathbb{Z}/p^k)$  the  $p$ -th power  $x^p$  lies in the image of the reduction map  $H^*(Y; \mathbb{Z}/p^{k+1}) \rightarrow H^*(Y; \mathbb{Z}/p^k)$ . The argument is as follows: If  $p$  is odd and the degree of  $x$  is odd,  $x^p = 0$  and the conclusion follows. Otherwise,  $\delta(x^p) = p\delta(x) \cdot x^{p-1} = 0$ , because the Bockstein  $\delta$  coming from the short exact sequence  $\mathbb{Z}/p \rightarrow \mathbb{Z}/p^{k+1} \rightarrow \mathbb{Z}/p^k$  is a derivation.

Therefore  $u^{p^d}$  lies in the image of the reduction  $H^*(Y; \mathbb{Z}/p^{d+1}) \rightarrow H^*(Y; \mathbb{F}_p)$ . Consider the two universal coefficients sequences relating the cohomology of  $Y$

with coefficients in  $\mathbb{F}_p$ , respectively in  $\mathbb{Z}/p^{d+1}$  to the cohomology of  $Y$  with coefficients in  $\mathbb{Z}_p^\wedge$  :

$$\begin{array}{ccccccc} 0 \rightarrow & H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{Z}/p^{d+1} & \longrightarrow & H^*(Y; \mathbb{Z}/p^{d+1}) & \longrightarrow & \mathrm{Tor}(H^{*+1}(Y; \mathbb{Z}_p^\wedge); \mathbb{Z}/p^{d+1}) & \rightarrow 0 \\ & \downarrow & & \downarrow & & \downarrow & \\ 0 \rightarrow & H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p & \xrightarrow{\rho} & H^*(Y; \mathbb{F}_p) & \xrightarrow{\partial} & \mathrm{Tor}(H^{*+1}(Y; \mathbb{Z}_p^\wedge); \mathbb{F}_p) & \rightarrow 0 \end{array}$$

where the vertical morphisms are induced by the mod  $p$  reduction  $\mathbb{Z}/p^{d+1} \rightarrow \mathbb{Z}/p$ . Since the  $p$ -adic cohomology groups are finitely generated and have exponent  $p^d$  we see that the morphism between the torsion groups on the right is zero. Thus  $\partial(u^{p^d}) = 0$ , which implies that  $u^{p^d}$  is in  $\mathrm{Im} \rho$ .  $\blacksquare$

**Lemma 2.3.** *Let  $Y$  be a connected space. If the graded  $\mathbb{F}_p$ -algebra  $H^*(Y; \mathbb{F}_p)$  is Noetherian and if  $H^*(Y; \mathbb{Z}_p^\wedge)$  has bounded torsion, then  $H^*(Y; \mathbb{F}_p)$  is a finitely generated module over  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ .*

*Proof.* This is clear since Lemma 2.2 implies that  $H^*(Y; \mathbb{F}_p)$  is integral over  $\mathrm{Im} \rho$ . Explicitly, let us choose homogeneous generators  $w_1, \dots, w_n$  of the graded algebra  $H^*(Y; \mathbb{F}_p)$  and consider the finite set  $W$  of monomials of the form  $w_1^{r_1} \cdots w_n^{r_n}$  with  $0 \leq r_i < p^d$ . We show that the set  $W$  generates  $H^*(Y; \mathbb{F}_p)$  as a module over  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ . For, consider any monomial  $m = w_1^{s_1} \cdots w_n^{s_n}$  in  $H^*(Y; \mathbb{F}_p)$ . Writing the exponents  $s_i = r_i + p^d t_i$  with  $0 \leq r_i < p^d$ , we express  $m = x^{p^d} \cdot w$  for a monomial  $w$  in  $W$  and an homogeneous element  $x$ . By Lemma 2.2,  $x^{p^d}$  lifts to an element  $a$  in  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$  and  $m = \rho(a) \cdot w$ .  $\blacksquare$

**Theorem 2.4.** *Let  $Y$  be a connected space. Then, the graded  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(Y; \mathbb{Z}_p^\wedge)$  is Noetherian if and only if the graded  $\mathbb{F}_p$ -algebra  $H^*(Y; \mathbb{F}_p)$  is Noetherian and the torsion in  $H^*(Y; \mathbb{Z}_p^\wedge)$  is bounded.*

*Proof.* Assume first that  $H^*(Y; \mathbb{Z}_p^\wedge)$  is a Noetherian  $\mathbb{Z}_p^\wedge$ -algebra. By Lemma 2.1,  $H^*(Y; \mathbb{F}_p)$  is a finitely generated module over  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ . Since  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$  is a Noetherian  $\mathbb{F}_p$ -algebra, it follows from [4, Proposition 7.2] that  $H^*(Y; \mathbb{F}_p)$  is also a Noetherian  $\mathbb{F}_p$ -algebra. The torsion part  $T_p H^*(Y; \mathbb{Z}_p^\wedge)$  is an ideal of

the Noetherian algebra  $H^*(Y; \mathbb{Z}_p^\wedge)$ , hence is finitely generated. The order of the torsion is thus bounded by the order of its generators.

Suppose now that  $H^*(Y; \mathbb{F}_p)$  is a Noetherian  $\mathbb{F}_p$ -algebra and that the torsion in  $H^*(Y; \mathbb{Z}_p^\wedge)$  is bounded. Then, by Lemma 2.3,  $H^*(Y; \mathbb{F}_p)$  is a finitely generated module over  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ . As a consequence of the graded version of the so-called Eakin-Nagata Theorem, see Proposition A.1, we infer then that the graded subring  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$  of  $H^*(Y; \mathbb{F}_p)$  is also Noetherian. Thus  $H^*(Y; \mathbb{Z}_p^\wedge)$  is a Noetherian  $\mathbb{Z}_p^\wedge$ -algebra by Corollary A.3.  $\blacksquare$

We end this section with an example which shows that Theorem 2.4 does not hold without the condition on torsion.

**Example 2.5.** Aguadé, Broto, and Notbohm constructed in [1] spaces  $X_k(r)$  for any odd prime  $p$  satisfying :  $H^*(X_k(r); \mathbb{F}_p) \cong \mathbb{F}_p[x_{2r}] \otimes E(\beta^{(k+1)}x_{2r})$  where  $\beta^{(k+1)}$  denotes the Bockstein of order  $k + 1$ . Observe that  $H^*(X_k(r); \mathbb{F}_p)$  is a Noetherian  $\mathbb{F}_p$ -algebra. The torsion of  $H^*(X_k(r); \mathbb{Z}_p^\wedge)$  is unbounded by [1, Remark 5.8]. Theorem 2.4 shows that the algebra  $H^*(X_k(r); \mathbb{Z}_p^\wedge)$  is not Noetherian.

### 3. COHOMOLOGY WITH TWISTED COEFFICIENTS

In this section we work over a ring  $R$  which is either  $\mathbb{Z}_p^\wedge$  or  $\mathbb{F}_p$ . Let  $Y$  be a connected space of which the fundamental group is a finite  $p$ -group. Let  $M$  be a  $R[\pi_1 Y]$ -module which is a finitely generated  $R$ -module. We aim to show that the cohomology with twisted coefficients  $H^*(Y; M)$  is Noetherian as a module over  $H^*(Y; R)$  if  $H^*(Y; R)$  is Noetherian. We shall deal separately with the field of  $p$  elements and with the ring of  $p$ -adic integers.

We start with a standard Noetherianity result.

**Lemma 3.1.** *Let  $R = \mathbb{Z}_p^\wedge$  or  $\mathbb{F}_p$ . Let  $Y$  be a space and let  $0 \rightarrow N \rightarrow M \rightarrow Q \rightarrow 0$  be a short exact sequence of  $R[\pi_1 Y]$ -modules. If both  $H^*(Y; N)$  and  $H^*(Y; Q)$  are Noetherian modules over  $H^*(Y; R)$ , then so is  $H^*(Y; M)$ .*

*Proof.* The long exact sequence in cohomology induced by the short exact sequence of modules is one of  $H^*(Y; R)$ -modules. It exhibits  $H^*(Y; M)$  as an extension of a submodule of  $H^*(Y; Q)$  by a quotient of  $H^*(Y; N)$ .  $\blacksquare$

**3.1. The case of  $\mathbb{F}_p$ -vector spaces.** To prove the next result we follow Minh and Symonds' approach for profinite groups, [19, Lemma 1].

**Theorem 3.2.** *Let  $Y$  be a connected space such that  $\pi_1 Y$  is a finite  $p$ -group and let  $M$  be a finite  $\mathbb{F}_p[\pi_1 Y]$ -module. If the graded  $\mathbb{F}_p$ -algebra  $H^*(Y; \mathbb{F}_p)$  is Noetherian, then  $H^*(Y; M)$  is Noetherian as a module over  $H^*(Y; \mathbb{F}_p)$ .*

*Proof.* We use induction on  $\dim_{\mathbb{F}_p} M$ . Since  $G = \pi_1(Y)$  is a finite  $p$ -group, the fixed points  $M^G$  is not trivial when  $M$  is not trivial. The induction step follows by applying Lemma 3.1 to the short exact sequence  $0 \rightarrow M^G \rightarrow M \rightarrow M/M^G \rightarrow 0$ . **■**

**3.2. The case of  $\mathbb{Z}_p^\wedge$ -modules.** We consider in this section the cohomology with twisted coefficient  $H^*(Y; M)$  of a connected space  $Y$  where  $M$  is a  $\mathbb{Z}_p^\wedge[\pi_1 Y]$ -module which is finitely generated over  $\mathbb{Z}_p^\wedge$ . In a first step, let  $M$  be a  $\mathbb{Z}_p^\wedge[\pi_1 Y]$ -module which is finite (meaning finite as a set).

**Lemma 3.3.** *Let  $Y$  be a connected space such that  $\pi_1 Y$  is a finite  $p$ -group. Let  $M$  be a  $\mathbb{Z}_p^\wedge[\pi_1 Y]$ -module which is finite. If the graded  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(Y; \mathbb{Z}_p^\wedge)$  is Noetherian, then  $H^*(Y; M)$  is Noetherian as a module over  $H^*(Y; \mathbb{Z}_p^\wedge)$ .*

*Proof.* The module  $M$  being finite,  $M$  is a finite abelian  $p$ -group. We perform an induction on the exponent  $e$  of  $M$ . When  $e = 1$ , the module  $M$  has the structure of an  $\mathbb{F}_p$ -vector space. As  $H^*(Y; \mathbb{F}_p)$  is a Noetherian  $\mathbb{F}_p$ -algebra by Theorem 2.4, we know from Theorem 3.2 that  $H^*(Y; M)$  is Noetherian as a module over  $H^*(Y; \mathbb{F}_p)$ . The Noetherian  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(Y; \mathbb{Z}_p^\wedge)$  acts on  $H^*(Y; M)$  through  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ . Therefore  $H^*(Y; M)$  is a Noetherian module over  $H^*(Y; \mathbb{Z}_p^\wedge)$ .

Let us now assume that  $e > 1$  and consider the short exact sequence  $0 \rightarrow M_p \rightarrow M \rightarrow Q \rightarrow 0$  where  $M_p$  is the submodule of  $M$  consisting of elements of order 1 or  $p$ . The induction step follows from Lemma 3.1. **■**

**Remark 3.4.** In the case of trivial coefficient modules our main tool was the universal coefficients exact sequence, but this does not exist in general for twisted coefficients. One basic counter example is given by the free module  $\mathbb{F}_p[G]$  for a finite group  $G$  whose order is divisible by  $p$ . Then  $H^*(BG; M)$  is zero in positive degrees and the universal coefficient formula does not hold.



In a second step we consider, as coefficient of the cohomology, a  $\mathbb{Z}_p^\wedge[\pi_1 Y]$ -module  $M$ , which is free of finite rank over  $\mathbb{Z}_p^\wedge$ .

**Lemma 3.5.** *Let  $Y$  be a connected space such that  $\pi_1 Y$  is a finite  $p$ -group. Let  $M$  be a  $\mathbb{Z}_p^\wedge[\pi_1 Y]$ -module which is free of finite rank over  $\mathbb{Z}_p^\wedge$ . If the graded  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(Y; \mathbb{Z}_p^\wedge)$  is Noetherian, then  $H^*(Y; M)$  is Noetherian as a module over  $H^*(Y; \mathbb{Z}_p^\wedge)$ .*

*Proof.* The short exact sequence  $0 \rightarrow M \xrightarrow{p} M \rightarrow M \otimes \mathbb{F}_p \rightarrow 0$  induces in cohomology a long exact sequence of  $H^*(Y; \mathbb{Z}_p^\wedge)$ -modules. We see that  $H^*(Y; M) \otimes \mathbb{F}_p$  is a sub- $H^*(Y; \mathbb{Z}_p^\wedge)$ -module of  $H^*(Y; M \otimes \mathbb{F}_p)$ . Since the action of  $H^*(Y; \mathbb{Z}_p^\wedge)$  on both  $H^*(Y; M \otimes \mathbb{F}_p)$  and  $H^*(Y; M) \otimes \mathbb{F}_p$  factors through  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ , it follows that  $H^*(Y; M) \otimes \mathbb{F}_p$  is a sub- $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ -module of  $H^*(Y; M \otimes \mathbb{F}_p)$ .

This takes us back to the world of  $\mathbb{F}_p$ -vector spaces and we know by Theorem 3.2 that  $H^*(Y; M \otimes \mathbb{F}_p)$  is a Noetherian module over  $H^*(Y; \mathbb{F}_p)$ . As the latter is a finitely generated module over  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$  by Lemma 2.1, we infer that  $H^*(Y; M \otimes \mathbb{F}_p)$  is a Noetherian module over  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ . Therefore  $H^*(Y; M) \otimes \mathbb{F}_p$  is a Noetherian module over  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$  as well, and since  $H^*(Y; \mathbb{Z}_p^\wedge)$  acts on  $H^*(Y; M) \otimes \mathbb{F}_p$  via  $H^*(Y; \mathbb{Z}_p^\wedge) \otimes \mathbb{F}_p$ , it is a Noetherian module over  $H^*(Y; \mathbb{Z}_p^\wedge)$ .

Set  $A^* = H^*(Y; \mathbb{Z}_p^\wedge)$  and  $N^* = H^*(Y; M)$ . Both are finitely generated  $\mathbb{Z}_p^\wedge$ -modules in each degree by Proposition 1.2, thus also Hausdorff and complete. We then conclude by applying Proposition A.2. ▀

We now prove our main theorem.

**Theorem 3.6.** *Let  $Y$  be a connected space such that  $\pi_1 Y$  is a finite  $p$ -group. Let  $M$  be a  $\mathbb{Z}_p^\wedge[\pi_1 Y]$ -module, which is finitely generated over  $\mathbb{Z}_p^\wedge$ . If the graded  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(Y; \mathbb{Z}_p^\wedge)$  is Noetherian, then  $H^*(Y; M)$  is Noetherian as a module over  $H^*(Y; \mathbb{Z}_p^\wedge)$ .*

*Proof.* Let  $TM$  be the torsion submodule of  $M$  and consider the short exact sequence of  $\mathbb{Z}_p^\wedge[\pi_1 Y]$ -modules  $0 \rightarrow TM \rightarrow M \rightarrow Q \rightarrow 0$ . We know from Lemma 3.3

that  $H^*(Y; TM)$  is a Noetherian  $H^*(Y; \mathbb{Z}_p^\wedge)$ -module and from Lemma 3.5 that so is  $H^*(Y; Q)$ . We conclude by Lemma 3.1.  $\blacksquare$

**Remark 3.7.** Our main theorem makes no assumption, except that the fundamental group be a  $p$ -group. One could try to relax it with transfer arguments, requiring a version of the transfer with twisted coefficients. However, recent work of Levi and Ragnarsson, in the context of  $p$ -local finite group theory, provides [17, Proposition 3.1] an example showing that such a transfer might not have, in general, the properties we need when the fundamental group of the space is not a  $p$ -group.

#### 4. THE CASE OF $p$ -COMPACT GROUPS AND $p$ -LOCAL FINITE GROUPS

We arrive at the promised application to  $p$ -compact groups and  $p$ -local finite groups. By definition, a  $p$ -compact group is a mod  $p$  finite loop space  $X = \Omega BX$ , where the “classifying space”  $BX$  is  $p$ -complete, [12].

**Lemma 4.1.** *Let  $X$  be a  $p$ -compact group. Then the  $p$ -torsion in  $H^*(BX; \mathbb{Z}_p^\wedge)$  is bounded.*

*Proof.* By [12, Proposition 9.9], any  $p$ -compact group admits a maximal  $p$ -compact toral subgroup  $S$  such that  $\iota: BS \rightarrow BX$  is a monomorphism and the Euler characteristic  $\chi$  of the homotopy fibre is prime to  $p$  (see [12, Proof of 2.4, page 430]). Dwyer constructed a transfer map  $\tau: \Sigma^\infty BX \rightarrow \Sigma^\infty BS$  in [11] such that  $\iota \circ \tau$  is multiplication by  $\chi$ . In particular,  $\iota^*: H^*(BX; \mathbb{Z}_p^\wedge) \rightarrow H^*(BS; \mathbb{Z}_p^\wedge)$  is a monomorphism. It is thus enough to show that  $H^*(BS; \mathbb{Z}_p^\wedge)$  has bounded torsion. Now, a  $p$ -toral compact group  $S$  can be constructed, up to  $p$ -completion, as an extension of a finite  $p$ -group  $P$  and a discrete torus  $H = \bigoplus \mathbb{Z}_{p^\infty}$ . The index of  $H$  in  $S$  is thus  $|P|$ , a power of  $p$ . A classical transfer argument shows then that multiplication by  $|P|$  on  $H^*(BS; \mathbb{Z}_p^\wedge)$  factors through the torsion free module  $H^*(BH; \mathbb{Z}_p^\wedge)$ .  $\blacksquare$

**Theorem 4.2.** *Let  $X$  be a  $p$ -compact group, let  $M$  be a finite  $\mathbb{F}_p[\pi_1 BX]$ -module, and let  $N$  be a  $\mathbb{Z}_p^\wedge[\pi_1 BX]$ -module which is finitely generated over  $\mathbb{Z}_p^\wedge$ . Then*

- (1) the  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(BX; \mathbb{Z}_p^\wedge)$  is Noetherian;
- (2) the module  $H^*(BX; M)$  is Noetherian over  $H^*(BX; \mathbb{F}_p)$ ;
- (3) the module  $H^*(BX; N)$  is Noetherian over  $H^*(BX; \mathbb{Z}_p^\wedge)$ .

*Proof.* The main theorem of Dwyer and Wilkerson, [12, Theorem 2.4], asserts that  $H^*(BX; \mathbb{F}_p)$  is Noetherian. Lemma 4.1 allows us to apply Theorem 2.4 to prove the first claim. The second claim follows then from Theorem 3.2 because  $\pi_1 BX$  is a finite  $p$ -group, [12, Lemma 2.1]. Finally Theorem 3.6 implies the third claim. ▀

**Remark 4.3.** Let us consider the case of  $BO(n)$  at the prime 2 (the fundamental group is cyclic of order 2). E.H. Brown made in [9] an explicit computation of the integral cohomology. He actually proves that the square of any even Stiefel-Whitney class  $w_{2i}^2$  belongs to the image of  $\rho$  and the technique we use in Lemma 2.3 is somewhat inspired by his computations. Even though the relations in the mod  $p$  cohomology of an arbitrary  $p$ -compact group (one which is not  $p$ -torsion free) make it difficult to exhibit explicit generators for the  $p$ -adic cohomology, Theorem 4.2 (1) gains qualitative control on it.

As for twisted coefficients, let  $\mathbb{Z}^\vee$  be a free abelian group of rank one, endowed with the sign action of the fundamental group  $C_2$ . In [10, Theorem 1] Čadek exhibits an explicit *finite* set of generators of  $H^*(BO(n); \mathbb{Z}^\vee)$ , as a module over  $H^*(BO(n); \mathbb{Z})$ . This is one of the few available explicit computations illustrating our results.

Broto, Levi, and Oliver defined in [8] the concept of  $p$ -local finite group. It consists of a triple  $(S, \mathcal{F}, \mathcal{L})$  where  $S$  is a finite  $p$ -group and,  $\mathcal{F}$  and  $\mathcal{L}$  are two categories whose objects are subgroups of  $S$ . The category  $\mathcal{F}$  models abstract conjugacy relations among the subgroups of  $S$ , and  $\mathcal{L}$  is an extension of  $\mathcal{F}$  with enough information to define a classifying space which behaves like the  $p$ -completed classifying space of a finite group. The classifying space of the  $p$ -local finite group is defined to be  $|\mathcal{L}|_p^\wedge$ .

**Lemma 4.4.** *Let  $(S, \mathcal{F}, \mathcal{L})$  be a  $p$ -local finite group. The  $p$ -torsion in  $H^*(|\mathcal{L}|_p^\wedge; \mathbb{Z}_p^\wedge)$  is then bounded.*

*Proof.* Ragnarsson proved in [22] that the suspension spectrum  $\Sigma^\infty(|\mathcal{L}|_p^\wedge)$  is a retract of  $\Sigma^\infty BS$ . Since the order of  $S$  annihilates all cohomology groups of  $BS$ , the same holds for  $H^*(|\mathcal{L}|_p^\wedge; \mathbb{Z}_p^\wedge)$ . ■

**Theorem 4.5.** *Let  $(S, \mathcal{F}, \mathcal{L})$  be a  $p$ -local finite group, let  $M$  be a finite  $\mathbb{F}_p[\pi_1(|\mathcal{L}|_p^\wedge)]$ -module, and let  $N$  be a  $\mathbb{Z}_p^\wedge[\pi_1(|\mathcal{L}|_p^\wedge)]$ -module which is finitely generated over  $\mathbb{Z}_p^\wedge$ . Then*

- (1) *the  $\mathbb{Z}_p^\wedge$ -algebra  $H^*(|\mathcal{L}|_p^\wedge; \mathbb{Z}_p^\wedge)$  is Noetherian;*
- (2) *the module  $H^*(|\mathcal{L}|_p^\wedge; M)$  is Noetherian over  $H^*(|\mathcal{L}|_p^\wedge; \mathbb{F}_p)$ ;*
- (3) *the module  $H^*(|\mathcal{L}|_p^\wedge; N)$  is Noetherian over  $H^*(|\mathcal{L}|_p^\wedge; \mathbb{Z}_p^\wedge)$ .*

*Proof.* We follow the same steps we took for  $p$ -compact groups in Theorem 4.2. The first ingredient is [8, Theorem B], which shows that  $H^*(|\mathcal{L}|_p^\wedge; \mathbb{F}_p)$  is computed by stable elements and Noetherian. We just proved that the torsion in  $H^*(|\mathcal{L}|_p^\wedge; \mathbb{Z}_p^\wedge)$  is bounded. Moreover, the fundamental group of  $|\mathcal{L}|_p^\wedge$  is a finite  $p$ -group by [8, Proposition 1.12]. ■

## APPENDIX A

This short appendix deals with Noetherianity in the graded case over the  $p$ -adics. We start however with a more general result, probably well-known to the experts: the graded Eakin-Nagata Theorem. The non-graded version can be found for example in Matsumura's book [18, Theorem 3.7(i)].

**Proposition A.1.** *Let  $A^*$  be a graded subring of  $B^*$ . Assume that  $B^*$  is Noetherian as a ring and finitely generated as an  $A^*$ -module. Then  $A^*$  is also a Noetherian ring.*

*Proof.* By [18, Theorem 13.1],  $B^0$  is Noetherian and  $B^*$  is a finitely generated  $B^0$ -algebra. Moreover,  $B^0$  is a finitely generated  $A^0$ -module and therefore  $B^*$  is

a finitely generated  $A^0$ -algebra. Also,  $A^0$  is Noetherian by the classical Eakin-Nagata theorem [18, Theorem 3.7(i)]. Applying [4, Proposition 7.8] to the inclusions  $A^0 \subset A^* \subset B^*$  we obtain that  $A^*$  is a finitely generated  $A^0$ -algebra. Again, by [18, Theorem 13.1],  $A^*$  is a Noetherian ring.  $\blacksquare$

The following technical proposition allows us to deduce Noetherianity over the  $p$ -adics from the Noetherianity of the mod  $p$  reduction.

**Proposition A.2.** *Let  $A^*$  be a graded  $\mathbb{Z}_p^\wedge$ -algebra such that in each degree  $A^k$  is complete for the  $p$ -adic topology. Let  $N^*$  be a graded  $A^*$ -module such that for all  $k$ ,  $N^k$  is Hausdorff for the  $p$ -adic topology. If  $N^* \otimes \mathbb{F}_p$  is a Noetherian  $A^*$ -module, then so is  $N^*$ .*

*Proof.* Let us choose homogeneous elements  $\nu_1, \dots, \nu_n \in N^*$  such that  $\nu_1 \otimes 1, \dots, \nu_n \otimes 1$  generate  $N^* \otimes \mathbb{F}_p$  as  $A^*$ -module. We claim that  $\nu_1, \dots, \nu_n$  generate  $N^*$  as an  $A^*$ -module. Given  $n \in N^*$  we may write  $n \otimes 1 = \sum a_i^0(\nu_i \otimes 1)$  for some  $a_i^0 \in A^*$ . Define  $n_0 = \sum a_i^0 \nu_i$  and notice that  $n - n_0 \in pN^*$ . Thus, there exists an element  $m_1 \in N^*$ , homogeneous of degree  $\leq \deg n$ , such that  $n - n_0 = pm_1$ . We iterate the procedure and find elements  $a_i^1 \in A^*$  such that  $m_1 \otimes 1 = \sum a_i^1(\nu_i \otimes 1)$ . We define  $n_1 = n_0 + p \sum a_i^1 \nu_i = \sum (a_i^0 + pa_i^1) \nu_i$ . In this way we construct, for any  $i$ , Cauchy sequences of coefficients  $(a_i^0 + pa_i^1 + \dots + p^k a_i^k)_k$  in  $A^*$ . By completeness this sequence converges to some  $a_i \in A^*$ . Since  $N^*$  is Hausdorff, the element  $\sum a_i \nu_i$  is equal to  $n$ .  $\blacksquare$

In the following corollary, the assumption that  $A^*$  be connected, i.e.  $A^0 = \mathbb{Z}_p^\wedge$ , is important. The referee kindly provided a non-connected counter-example, namely  $A^* = \mathbb{Q}_p^\wedge(t)$ , which is Noetherian, but not finitely generated as a  $\mathbb{Z}_p^\wedge$ -algebra.

**Corollary A.3.** *Let  $A^*$  be a graded connected Hausdorff  $\mathbb{Z}_p^\wedge$ -algebra. If  $A^* \otimes \mathbb{F}_p$  is a Noetherian  $\mathbb{F}_p$ -algebra, then  $A^*$  is a Noetherian  $\mathbb{Z}_p^\wedge$ -algebra.*

*Proof.* Since  $\mathbb{Z}_p^\wedge$  is Noetherian and  $A^*$  is connected,  $A^*$  is a Noetherian  $\mathbb{Z}_p^\wedge$ -algebra if and only if  $A^*$  is a finitely generated  $\mathbb{Z}_p^\wedge$ -algebra, [18, Theorem 13.1]. Note that

$A^* \otimes \mathbb{F}_p$  is also a Noetherian  $\mathbb{Z}_p^\wedge$ -algebra via the mod  $p$  reduction  $\mathbb{Z}_p^\wedge \rightarrow \mathbb{F}_p$ . Let us choose homogeneous elements  $\gamma_1, \dots, \gamma_n \in A^*$  such that  $\gamma_1 \otimes 1, \dots, \gamma_n \otimes 1$  generate  $A^* \otimes \mathbb{F}_p$  as  $\mathbb{Z}_p^\wedge$ -algebra. For a fixed  $k \geq 0$ ,  $A^k \otimes \mathbb{F}_p$  is generated as a  $\mathbb{Z}_p^\wedge$ -module by the monomials  $(\gamma_1 \otimes 1)^{e_1} \cdots (\gamma_n \otimes 1)^{e_n}$  with  $\sum_{i=1}^n |\gamma_i| e_i = k$ . Since  $A^k$  is a Hausdorff  $\mathbb{Z}_p^\wedge$ -module, the proof of Proposition A.2 shows that  $A^k$  is generated by the monomials  $\gamma_1^{e_1} \cdots \gamma_n^{e_n}$  with  $\sum_{i=1}^n |\gamma_i| e_i = k$ . This shows that  $A^*$  is generated as a  $\mathbb{Z}_p^\wedge$ -algebra by the elements  $\gamma_1, \dots, \gamma_n \in A^*$  and therefore  $A^*$  is a Noetherian  $\mathbb{Z}_p^\wedge$ -algebra. ■

## REFERENCES

1. J. Aguadé, C. Broto, and D. Notbohm, *Homotopy classification of spaces with interesting cohomology and a conjecture of Cooke, part I*, *Topology* **33** (1994), 455–492.
2. K. K. S. Andersen and J. Grodal, *The classification of 2-compact groups*, *J. Amer. Math. Soc.* **22** (2009), no. 2, 387–436.
3. K. K. S. Andersen, J. Grodal, J. M. Møller, and A. Viruel, *The classification of  $p$ -compact groups for  $p$  odd*, *Ann. of Math. (2)* **167** (2008), no. 1, 95–210.
4. M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.
5. D. J. Benson and N. Habegger, *Varieties for modules and a problem of Steenrod*, *Proceedings of the Northwestern conference on cohomology of groups* (Evanston, Ill., 1985), vol. 44, 1987, pp. 13–34.
6. A. K. Bousfield, *On the  $p$ -adic completions of nonnilpotent spaces*, *Trans. Amer. Math. Soc.* **331** (1992), no. 1, 335–359.
7. A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Springer-Verlag, Berlin, 1972, *Lecture Notes in Mathematics*, Vol. 304.
8. C. Broto, R. Levi, and B. Oliver, *The homotopy theory of fusion systems*, *J. Amer. Math. Soc.* **16** (2003), no. 4, 779–856 (electronic).
9. E. H. Brown, Jr., *The cohomology of  $BSO_n$  and  $BO_n$  with integer coefficients*, *Proc. Amer. Math. Soc.* **85** (1982), no. 2, 283–288.
10. M. Čadek, *The cohomology of  $BO(n)$  with twisted integer coefficients*, *J. Math. Kyoto Univ.* **39** (1999), no. 2, 277–286.
11. W. G. Dwyer, *Transfer maps for fibrations*, *Math. Proc. Cambridge Philos. Soc.* **120** (1996), no. 2, 221–235.

12. W. G. Dwyer and C. W. Wilkerson, *Homotopy fixed-point methods for Lie groups and finite loop spaces*, Ann. of Math. (2) **139** (1994), no. 2, 395–442.
13. L. Evens, *The cohomology ring of a finite group*, Trans. Amer. Math. Soc. **101** (1961), 224–239.
14. ———, *The spectral sequence of a finite group extension stops*, Trans. Amer. Math. Soc. **212** (1975), 269–277.
15. E. Golod, *The cohomology ring of a finite  $p$ -group*, Dokl. Akad. Nauk SSSR **125** (1959), 703–706.
16. R. Greenblatt, *Homology with local coefficients and characteristic classes*, Homology, Homotopy Appl. **8** (2006), no. 2, 91–103 (electronic).
17. R. Levi and K. Ragnarsson,  *$p$ -local finite group cohomology*, arxiv preprint 0910.5639.
18. H. Matsumura, *Commutative ring theory*, second ed., Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1989, Translated from the Japanese by M. Reid.
19. P. A. Minh and P. Symonds, *Cohomology and finite subgroups of profinite subgroups*, Preprint available at <http://www.maths.abdn.ac.uk/bensondj/html/archive/minh-symonds.html>.
20. I. Moerdijk and J.-A. Svensson, *The equivariant Serre spectral sequence*, Proc. Amer. Math. Soc. **118** (1993), no. 1, 263–278.
21. J. M. Møller,  *$N$ -determined 2-compact groups. II*, Fund. Math. **196** (2007), no. 1, 1–90.
22. Kári Ragnarsson, *Classifying spectra of saturated fusion systems*, Algebr. Geom. Topol. **6** (2006), 195–252.
23. E. H. Spanier, *Algebraic topology*, Springer-Verlag, New York, 1981, Corrected reprint.
24. B. B. Venkov, *Cohomology algebras for some classifying spaces*, Dokl. Akad. Nauk SSSR **127** (1959), 943–944.