
Fluvial and eolian ichnofaunas from the Lower Permian of South America (Patquía Formation, Paganzo Basin)

V. KRAPOVICKAS^{|1|} A.C. MANCUSO^{|2|} A. ARCUCCI^{|3|} A. CASELLI^{|1|}

^{|1|} **Departamento de Ciencias Geológicas, FCEN, UBA**

Ciudad Universitaria Pabellón II, 1428 Buenos Aires, Argentina. Krapovickas E-mail: vkrapovickas@gl.fcen.uba.ar
Caselli E-mail: acaselli@gl.fcen.uba.ar

^{|2|} **IANIGLIA, CCT-CONICET, Mendoza**

Adrián Ruiz Leal s/n, Parque Gral. San Martín, 5500 Mendoza CC330. Argentina E-mail: amancu@lab.cricyt.edu.ar

^{|3|} **Área de Zoología, Universidad Nacional de San Luis**

Chacabuco 917, 5700 San Luis Argentina. E-mail: andrea.arcucci@gmail.com

| A B S T R A C T |

The Lower Permian Patquía Formation is the youngest unit of the Paganzo Basin, western Argentina. The lower section consists of red mudstones, and fine- and coarse-grained sandstones deposited in fluvial systems with extensive and thick floodplain deposits. These rocks contain a low-diversity and relatively abundant association of trace fossils suggesting the activity of a sub-superficial to superficial fauna. The association is characterized by *Rusophycus carbonarius* (Cubichnia), *Cruziana problematica* (Repichnia), and *Palaeophycus tubularis* (Domichnia) of the *Scoyenia* ichnofacies. Disarticulated fish remains are also present. The upper section is dominated by red, cross-bedded, medium- to fine-grained sandstones deposited in eolian systems that host a low-diversity and low-abundance association of trace fossils that indicates the activity of a mainly superficial fauna. Tetrapod footprints (*Chelichnus duncani*, oval digit imprints, short parallel grooves, and sinusoidal grooves), horizontal to vertical burrows (*Palaeophycus tubularis* and *Skolithos* isp. respectively), and arthropod trackways are the typical components of these deposits. The association shows elements of the three presently proposed eolian ichnofacies (*Chelichnus*, *Octopodichnus* and *Entradichnus* ichnofacies) suggesting the necessity of revision and the possible integration of these separate ichnofacies into a single model. The record of fossil vertebrate tracks is uncommon in Lower Permian strata of South America. Therefore, the ichnologic record of the Patquía Formation is a significant contribution to the understanding of Lower Permian South American ichnofaunas.

KEYWORDS | *Scoyenia* ichnofacies. Eolian ichnofacies. Tetrapod footprints. Early Permian. Patquía. Formation.

INTRODUCTION

Upper Paleozoic ichnofaunas have been extensively recorded in the Northern Hemisphere including both ver-

tebrate and invertebrate trace fossils (e.g. Gilmore, 1926; Hunt et al., 1995; Haubold and Lucas, 2003; Minter, et al., 2007). The records of trace fossils mostly correspond to strata assigned by different authors to the Lower Perm-

ian (Artinskian-Kungurian) including the well-known occurrences from the Robledo Mountains in the USA. The tetrapod trackway associations have been divided in two groups, recently formalized as the *Chelichnus* and *Batrachichnus* ichnofacies, corresponding to eolian and redbed depositional settings respectively (Hunt et al., 1995; Hunt and Lucas, 2007).

Continental body fossils and trace fossils are very rare in South America considering the widespread exposure of Permian rocks in various basins. The great majority of Permian tetrapods and fishes are recorded in Brazil, although a precise chronostratigraphic framework is still lacking (e.g. Richter and Langer, 1998). Recently, new outcrops of undetermined Permian age have been found in the Buena Vista Formation, Uruguay. A diverse but fragmentary record of tetrapod body fossils has been recently recovered and is currently under study (e.g. Piñeiro et al., 2007).

In Argentina, the Permian record of tetrapod body fossils is scarce, consisting only in sparse occurrences (Siano, 1990; Cisterna and Simanaukas, 2000). Tetrapod footprints are known in three geological units from three different late Paleozoic basins in Argentina. Eolian facies in the Yacimiento Los Reyunos Formation at Southern Mendoza province yields a tetrapod footprint association dominated by *Chelichnus* (Cei and Gargiulo, 1977; Melchor, 2001). $^{40}\text{Ar}/^{39}\text{Ar}$ analysis indicates a Wordian/Capitanian (Kazanian/Tatarian) age of 266.31 ± 0.82 Ma. for the Yacimiento Los Reyunos Formation (Melchor, 2000). The Carapacha basin records diverse invertebrate and tetrapod trackway associations that were compared with what was later called the *Batrachichnus* ichnofacies, and corresponds to shallow-lake facies (Melchor and Sarjeant, 2004). Paleofloristic elements indicate a late Early Permian to early Late Permian age for the Carapacha Formation (Melchor and Césari, 1991). Finally, there are several isolated occurrences of tetrapod footprints in eolian facies of the Lower Permian Patquía Formation (Paganzo Basin), in La Rioja Province. These tetrapod footprints were compared provisionally with *Dromopus* and *Gilmoreichnus* by Caselli and Arcucci (1999), and are re-analyzed in this contribution.

Permian invertebrate trace fossils from Argentina have been documented mainly in fluvial and shallow-lacustrine deposits (Aceñolaza and Buatois, 1991; 1993), floodplains of high-sinuosity rivers (Buatois et al., 1996), low-energy ponds in floodplains of braided fluvial systems (Buatois et al., 1997), and playa-lake complexes (Zhang et al., 1998). Little is known of invertebrate traces in eolian environments. Globally, trace fossil assemblages reported from eolian facies are scarce in the geologic record and more examples are needed to improve our knowledge of such ichnofaunas. Several eolian ichnofacies models have been

proposed: the *Octopodichnus* ichnofacies (Hunt and Lucas, 2007), *Entradichnus* ichnofacies (Morrissey, 2006; Ekdale et al., 2007) and the *Chelichnus* ichnofacies (Hunt and Lucas, 2007), although they seem to be superposed.

The ichnologic record of Argentina provides the only evidence of the distribution of tetrapods for the Early Permian. There has been little crossover between the studies of vertebrate and invertebrate ichnofossils in nonmarine ichnology (cf. Lockley, 2007). Herein we use a more integrated approach to study tetrapod footprints and invertebrate trace fossils from the Lower Permian Patquía Formation. These findings, although fragmentary, are the first record of tetrapods in the Paganzo basin and may be the oldest in the Permian of Argentina. The main purpose of this contribution is to document invertebrate and vertebrate trace fossils in the Lower Permian Patquía Formation. In addition, we attempt to discuss this evidence in a paleobiologic context, and to compare the ichnofauna with other Lower Permian associations.

Specimens collected are preserved at (CRILAR) Centro Regional de Investigaciones Científicas y Transferencia Tecnológica (CRILAR Ic), Anillaco, La Rioja Province, Argentina.

GEOLOGIC SETTING

The Paganzo Basin was developed as a foreland basin in western Argentina, bounded to the east by the Sierras Pampeanas cratonic area (Limarino et al., 2002). The basin includes one of the most important Late Paleozoic successions of South America. The Carboniferous successions record the Gondwana glaciation (Limarino et al., 2002; Geuna et al., 2010). The thick and geographically extensive Permian redbed successions record the arid and semiarid continental events observed globally (Spalletti et al., 2010; Tedesco et al. 2010). Outcrops of this basin are documented in Sierras Pampeanas, Precordillera, and the Famatina System (La Rioja, San Juan, and San Luis provinces).

The succession outcropping nearby the town of Paganzo is the type section of the Carboniferous-Permian “Estratos de Paganzo” (Bodenbender, 1911, 1912), subsequently referred to as the “Paganzo Group” (Azcuy and Morelli, 1970). This succession is located in the eastern domain of the basin (Limarino et al., 2002) in the central-south area of La Rioja province (Fig.1). The upper Paleozoic strata unconformably overlie the crystalline basement (Azcuy et al., 1979). They include grey sandstones and mudstones of the Carboniferous Lagares Formation, and Permian red beds of the Patquía Formation. This upper Paleozoic succession is overlain by the Triassic Talampaya Formation (Fig. 1).

The sedimentary facies of the Patquía Formation were studied in detail by Caselli (1998). This author included several Permian redbed successions that occur in the basin at the Patquía Formation (see also Caselli and Limarino, 2002). Two intervals within the Patquía Formation were recognized in the Paganzo area (Fig. 2). The lower interval is 240m-thick, and is characterized by red mudstones and fine-grained sandstones interbedded with light red pebbly and coarse-grained sandstones. The tabular mudstones and fine-grained sandstone beds are either massive or show parallel- or ripple cross-lamination. Individual beds are 0.20-1.5m thick. These deposits contain invertebrate trace fossils (*Rusophycus carbonarius*, *Cruziana problematica*, and *Palaeophycus tubularis*) and disarticulated fish remains (Fig.2). The lenticular pebbly and coarse-grained sandstone beds are 0.20- 0.80m thick and display parallel, planar, and trough cross-stratification. The lower interval was interpreted as deposited in fluvial environments, in which floodplain areas dominated over channel subenvironments (Caselli, 1998).

The upper interval is 160 m-thick and is dominated by red, high-angle planar cross-bedded, well-sorted, ma-

ture, medium- to fine-grained sandstones. The basal succession is dominated by 0.10-0.20m thick, parallel-laminated, ripple cross-laminated, and planar cross-stratified sandstones interpreted as low-angle sand-sheet deposits (Caselli, 1998). This facies records invertebrate trace fossils (*Palaeophycus tubularis*, *Skolithos* isp., and arthropod trackway), and a variety of tetrapod fossil footprints (*Chelichnus duncani*, oval digit imprints, short parallel grooves, and sinusoidal grooves) (Fig.2). The succession passes upwards into thick-bedded, high-angle cross-stratified sandstone beds. This interval has been interpreted as dune and interdune deposits of an extensive eolian system developed under arid conditions (Caselli, 1998).

The age of the Patquía Formation is poorly constrained. The fossil content suggests an Early Permian age (e.g. Frenguelli, 1949; Limarino and Césari, 1985). Paleomagnetic information and absolute ages (K-Ar) of basalt interbedded near the base of the Patquía Formation in the Paganzo area indicate a Late Carboniferous to Early Permian age (263 ± 4 a 302 ± 6 M.a.) (Thompson and Mitchell, 1972; Valencio, 1972; Valencio and Mitchell, 1972; Valencio et al., 1977; Sinito et al., 1979).




FIGURE 1 | Geologic map of the Paganzo Group in the Paganzo area.

INVERTEBRATE TRACE FOSSILS

Ichnogenera are placed in alphabetic order, and are followed by those described in open nomenclature. The preservational nomenclature of Seilacher (1964) is followed.

Ichnogenus *Cruziana* d’Orbigny, 1842
Cruziana problematica (Schindewolf, 1928)
 Figure 3A, 3C, 3D, 3E

Material. CRILAR Ic-2, CRILAR Ic-3, CRILAR Ic-4 and material preserved in the field.

Description. Two parallel rounded ridges separated by a central groove. The ridges are generally featureless, al-

though some traces display fine, nearly transverse, striae. External width is from 1.5-8.0mm. Preserved as convex hypichnia.

Remarks. This ichnospecies displays two main size ranges in the Patquía Formation: 1.5-3mm (highly abundant) and 5-8mm (Fig. 3C-D).

The distinction between *Isopodichnus*, *Rusophycus*, and *Cruziana* has been largely discussed (e.g. Bromley and Asgaard, 1972, 1979; Trewin, 1976; Pollard, 1985; Keighley and Pickerill, 1996). A few authors (e.g. Hantzschel, 1975; Seilacher, 1955) have regarded *Rusophycus* as a junior synonym of *Cruziana*, although others have argued that *Rusophycus* must be retained for short bilobate resting traces (e.g. Osgood, 1970). The ichnogenus *Isopodichnus* has also been traditionally applied to small bilobate trace fossils in continental environments, particularly redbeds (Trewin, 1976; Seilacher, 1985; Pollard, 1981, 1985), covering both crawling and resting traces. Bromley and Asgaard (1979) placed these bilobate traces in the ichnogenera *Cruziana* and *Rusophycus*, a decision endorsed here. It is highly accepted between ichnologists that paleoenvironment is not a valid ichnotaxobase (Bertling et al., 2006). The presence of *Cruziana problematica* has been documented for the Patquía or equivalent formations (e.g. Aceñolaza and Buatois, 1991, 1993; Zhang et al., 1998). *Cruziana* is known from the Cambrian to the Recent.

Ichnogenus *Palaeophycus* Hall, 1847
Palaeophycus tubularis Hall, 1847
 Figure 4A, 4C

Material. CRILAR Ic-5 and material preserved in the field.

Description. Horizontal to sub-horizontal, straight to slightly curved, unbranched and branched tubular trace fossils. Circular to elliptical sections are 5-10mm in diameter. Infill similar to the host rock and smooth lining. Preserved as full relief.

Remarks. *Palaeophycus* differs from *Planolites* by the presence of a wall lining and infill similar to that of the host rock (Pemberton and Frey, 1982). The systematics of *Palaeophycus* has been reviewed by Pemberton and Frey (1982). *Palaeophycus tubularis* is distinguished from the other ichnospecies of *Palaeophycus* by its thin burrow lining and lack of conspicuous wall ornamentation. The presence of *Palaeophycus tubularis* was documented for the Patquía or equivalent formations (e.g. Aceñolaza and Buatois, 1991, 1993; Zhang et al., 1998). The ichnogenus is present from the Ediacaran to the recent (Pemberton and Frey, 1982).




FIGURE 2 | Sedimentologic section of the Patquía Formation in the Paganzo area. F: mudstone; Sf: fine-grained sandstone; Sm: medium-grained sandstone; Sg: coarse-grained sandstone; G: pebble conglomerate.




FIGURE 3 | *Rusophycus carbonarius* (black arrow), note transverse scratch marks, and *Cruziana problematica* (white arrow). A) CRILAR Ic-2, general view. B) CRILAR Ic-3, note abundant small *Cruziana problematica*. C) CRILAR IC-4, *Cruziana problematica* of medium size. D) CRILAR Ic-2, note small *Cruziana problematica*. E) CRILAR Ic-4. Scale bars are 1 cm.

Ichnogenus *Rusophycus* Hall 1852

Rusophycus carbonarius Dawson, 1864

Figure 3A, 3B, 3D, 3E

Material. CRILAR Ic-2, CRILAR Ic-3, CRILAR Ic-4 and material preserved in the field.

Description. Subcircular trace fossil consisting of two parallel lobes separated by a central groove and resembling a coffee bean. Some specimens display an anterior subtriangular gap (see Fig. 3D). Lobes are covered by fine transverse striae that extend almost to the margin. Lengths are 5-9mm and widths are 5-10mm. Preserved as convex hypichnia.

Remarks. Schlirf et al. (2001) noted that, in contrast to the widely held view, *Rusophycus (Isopodichnus) eutendorfensis* does not in fact possess transverse to oblique striations because the original description by Linck (1942) identifies the presence of mostly smooth lobes, separated

by a longitudinal furrow. Thus, *R. carbonarius* is distinguished from *R. eutendorfensis* by the presence of transverse striae (Schlirf et al., 2001), from *R. stromnessi* by the absence of expanded, smooth and lily-like ends (Trewin, 1976; Keighley and Pickerill, 1996), from *R. furcosus* in the arrow shape of the latter (Gand, 1994), and from *R. minutus* in its triangular shape (Debriette and Gand, 1990). This ichnogenus is present from the Cambrian to the recent.

Ichnogenus *Skolithos* Haldeman, 1840

Skolithos isp.

Figure 4A, 4B

Material. Material preserved in the field.

Description. Simple, elongate, cylindrical vertical trace fossils. Wall is smooth and infill similar to the host rock. Diameter is 8-11mm. Short tubes with unknown lower end. Preserved as full relief.

Remarks. Specimens are typically preserved in bedding-plane view with maximum observed depths up to 30mm. No ichnospecific assignment can be made for the material from the Patquía Formation due to its partial preservation. *Skolithos* ranges from the Cambrian to the recent (Alpert, 1974; Seilacher et al., 2005).

Arthropod trackway

Figure 5A, 5D, 5H

Material. One trackway with nine tracks preserved in the field.

Description. Partially preserved trackway consisting of one row of tracks grouped in a series of two to four tracks with a medial impression. Individual imprints are oval or bifurcating tracks of approximately 13-20mm long. The trackway presents series of tracks partially preserved. It varies from two tracks arranged in a straight line, three tracks arranged in a triangular pattern, and four tracks arranged in a rhomboidal parallelogram. The distance between the medial impression and the outer track range from 50-64mm.

Remarks. This trackway resemble *Octopodichnus didactylus* in the large size of the trackway, the presence of bifurcated tracks, and tracks grouped in a series of four. On the other hand, it shows a medial impression, characteristic of specimens assigned to *Paleohelcura*. The ichnogenera *Paleohelcura* and *Octopodichnus* are similar enough in their morphology that difficulty in distinguishing between the two may arise (Sadler, 1993). At present, the two clearly distinguishing features of these two ichnogenera are the presence or absence of both a bifurcated track impression and a medial impression (Sadler, 1993). Unfortunately, this trackway can not be assigned with certainty to either ichnogenera (*Paleohelcura* or *Octopodichnus*) due to the fact that it presents characteristics mutually exclusive between them (bifurcated tracks and medial impression).

TETRAPOD TRACE FOSSILS

Tetrapod tracks and trackways are preserved on three closely spaced surfaces. All three levels comprise a single trampling surface where numerous imprints of appendages and tails are preserved (Fig. 5). Ichnogenera are placed in alphabetic order and are followed by those described in open nomenclature. The preservational nomenclature of Seilacher (1964) is followed. Material was studied directly in the field, and is preserved as concave epichnia.

Ichnogenus *Chelichnus* Jardine 1850

Chelichnus duncani Jardine 1850

Figure 5A, 5B, 5E




FIGURE 4 | *Palaeophycus tubularis* and *Skolithos* isp. A) *Palaeophycus tubularis* and *Skolithos* isp. preserved in eolian deposits. B) *Skolithos* isp. preserved in eolian deposits. C) CRILAR Ic-5, *Palaeophycus tubularis* preserved in fluvial deposits. White areas on scales are 1cm.




FIGURE 5 | Tetrapod trampling surfaces. A) Diagram of the trampling surfaces showing all the specimens. Scale bar is 5cm. B) Detail of the upper right area of the diagram showing *Chelichnus duncani*, oval digit imprints, short parallel grooves, and sinusoidal grooves. Scale bar is 3cm. C) Detail of the lower left area of the diagram showing oval digit imprints, short parallel grooves, and sinusoidal grooves. Scale bar is 3cm. D) Detail of the upper left area of the diagram showing an arthropod trackway. Note bifurcated track (black arrow) and medial impression (white arrow). Scale bar 10cm. E) *Chelichnus duncani*. Scale bar 3cm. F) Short parallel grooves. Scale bar 3cm. G) Oval digit imprints. Two consecutive steps, note digit drag mark (arrow). Scale bar 3cm. H) Upper left area of the diagram, note ripple marks (arrow).

Material. Three isolated tracks.

Description. Plantigrade tracks that show a rounded sole imprint and four subparallel, relatively short, anteriorly directed digit imprints. Footprint length is 32-37mm and footprint width is 28-35mm. Digit imprints are of approximately the same length as the sole imprint and lateral digit imprints (7-12mm) are shorter than central digit imprints (12-15mm). There is no clear trackway pattern of these footprints expressed on the trampling surface.

Remarks. According to McKeever and Haubold (1996) *Chelichnus* consists of rounded manus and pes tracks of nearly equal size. Complete manus and pes tracks are pentadactyl with a round sole imprint and five short digit imprints, although commonly only three or four digit imprints are visible. There are four recognized ichnospecies of *Chelichnus* differing principally in size. *Chelichnus bucklandi* has a pes track length of 10-25mm, *C. duncani* has a pes track length of 25-75mm, *C. gigas* has a pes track length of 75-125 mm, and *C. titan* displays a pes track length exceeding 125mm (McKeever and Haubold, 1996). The Patquía specimens are assigned to *Chelichnus duncani* on the basis of the plantigrade tracks with short and subparallel digit imprints that are oriented anteriorly.

Oval digit imprints

Figure 5A, 5B, 5C, 5G

Material. Eighteen individual tracks.

Description. Circular to oval digit imprints (4-7mm in width) parallel to each other and forming an arc. Distances between digit imprints are 2-3mm. In well-preserved specimens there are five digit imprints, although the number of digit imprints is commonly lower. It is not possible to observe their relation to forefoot or hind foot. Trackway pattern unclear.

Remarks. These imprints are most likely a differential preservation of *Chelichnus duncani* because the digit imprints of that ichnospecies are in the same size range.

Short parallel grooves

Figure 5A, 5B, 5C, 5F

Material. Six individual tracks.

Description. Parallel slightly curved grooves with pointed tips. Well-preserved specimens consist of four parallel grooves, although more partially preserved specimens display only two parallel grooves. Individual groove length is 10-20mm. They are sharply incised and are more similar to digit scratch marks than to digit imprints. No distinct trackways can be observed.

Remarks. Parallel furrows probably consist of scratch marks of a plantigrade quadruped with clawed tips. The Patquía parallel furrows strongly suggest a similar morphological pattern to that expressed by tracks where plantigrady and number of digits is typically reduced. Under somewhat dry conditions or in undertrack preservation, the number of digit imprints could gradually disappear.

Sinusoidal grooves

Figure 5A, 5B, 5C, 5G

Material. Six sinusoidal grooves.

Description. Sinusoidal grooves are present all over the tracking surface. They consist of sharp, curved impressions that locally show two parallel grooves but commonly consist of only one simple groove. The length of the grooves is 40-140mm. Small lobes commonly occur on the sides of the grooves, and represent sand deformation features.

Remarks. These sinusoidal grooves probably represent digit and tail drag marks.

DISCUSSION

Trace-fossil producers

Freshwater examples of *Cruziana* and *Rusophycus* have been interpreted as cubichnia and repichnia, respectively, of branchiopod crustaceans (notostracans) by Bromley and Asgaard (1972). These are mostly benthic organisms that crawl and burrow along the bottom of ephemeral ponds in continental environments. According to Bromley and Asgaard (1979), the morphology of very short freshwater *Rusophycus* with transverse striations could be compared with head-down burrowing behavior of trilobites of Seilacher (1970). As noted by Minter et al. (2007), anostracan and lipostracan branchiopods cannot be ruled out as possible producers of these striated trails. Some anostracans have been observed to feed by scraping the surface, and lipostracans, although extinct, are inferred to have had the same behavior on the basis of comparative functional morphology. Both scrape the substrate rather than ploughing through it, and so they are more likely to have produced superficial scratch traces rather than furrowed traces (Minter et al., 2007). Lipostracans are very small and so they can only be considered as potential tracemakers of the smallest (1.5-3mm) trace fossils.

Pemberton and Frey (1982) proposed several potential producers for *Palaeophycus*, in the marine realm. These structures were interpreted as passively infilled dwelling burrows of worm-like organisms. As suggested by Krapovickas et al. (2008 and references therein), some




FIGURE 6 | Idealized reconstruction of trace fossil associations of the Patquía Formation. Above, trace fossil association 2 developed in eolian sand-sheet deposits. Note its similarities with the three presently known eolian ichnofacies. Below, trace fossil association 1 developed in floodplain deposits. A: *Arenicolites* isp.; Ar: Arthropod trackway; C: *Chelichnus duncani*; Cr: *Cruziana problematica*; D: *Diplocraterion parallelum*; E: *Entradichnus meniscus*; O: *Octopodichnus* isp.; P: *Paleohelcura* isp.; Pa: *Palaeophycus tubularis*; Pl: *Planolites beverleyensis*; R: *Rusophycus carbonarius*; S: *Skolithos* isp.; T: *Taenidium serpentinum*.

modern producers of *Palaeophycus*-like structures in continental floodplains are semiaquatic organisms such as orthopterans (Tridactylidae), hemipterans (Salidae), beetles (Heteroceridae) and, non-aquatic cicindelinae beetles.

In non-marine environments, *Skolithos* is commonly recorded in subaerial and permanently subaqueous sediments (Gierlowski-Kordesch, 1991). Among recent organisms that construct vertical excavations are numerous groups of insects and arachnids (Smith and Hein, 1971, Ratcliffe and Fagerstrom, 1980, Bown, 1982).

The presence of two to four tracks in each series of imprints in the arthropod trackway indicates at least four appendages on each side of the body of the trackmaker, suggesting a member of the class Arachnida. In addition, the presence of a medial impression suggests a tail drag left by a scorpionid.

The poor preservation of the tetrapod tracks obscures diagnostic morphologic details of tracks and trackways, making it difficult to determine potential tracemakers. Specimens assigned to *Chelichnus* are interpreted to be

produced by a non-mammalian synapsid (McKeever and Haubold, 1996).

Paleoecology and taphonomy of trace fossils

Fluvial deposits in the Patquía Formation comprise scarce fluvial channels and thick floodplain deposits with shallow-water bodies that contain an abundant invertebrate ichnofauna. The assemblage (trace-fossil association 1) includes crawling, resting and dwelling trace fossils (Fig. 6, 7) corresponding to *Rusophycus carbonarius* (Cubichnia), *Cruziana problematica* (Repichnia), and *Palaeophycus tubularis* (Domichnia). Disarticulated fish scales are also present. This low-diversity ichnofauna suggests the activity of a sub-superficial to superficial benthic invertebrate fauna. The taphonomic features of both invertebrate trace fossils and fish remains (*i.e.* disarticulate and dispersed scales and fish bones, of which some show a patterned arrangement and are densely packed and well sorted) suggest that they were preserved in a floodplain environment characterized by an alternation of overflow and fall-out deposits (Mancuso, 2008).

Eolian sand-sheet deposits of the Patquía Formation include tetrapod trace fossils (*Chelichnus duncani*, oval digit imprints, short parallel grooves, and sinusoidal grooves) and invertebrate locomotion and dwelling trace fossils (arthropod trackway, *Palaeophycus tubularis*, and *Skolithos* isp.) (Fig. 6, 7). This low-diversity and low-abundance association (trace-fossil association 2) represents the activity of a terrestrial superficial ichnofauna produced by cursorial tetrapods and shallow burrowing desert-dwelling arthropods. The absence of morphologic details and the preservation of only the deepest features suggest that the tetrapod footprints in association 2 are preserved as “undertracks” and/or that the surface was partially eroded before final burial. The sediment grain-size (medium- to fine-grained sand) compared to the size of the footprints prevents the preservation of fine morphologic details. In addition, the production of tracks is linked to sediment grain size and substrate water content. Thus, in eolian environments moist sand is needed to preserve tracks (Fornós et al., 2002; Loope, 2006). The Patquía tetrapod trackway assemblage must have therefore been emplaced in a moist substrate. Therefore, the taphonomic characteristics allow us to interpret the track-bearing levels as humid sand-sheet deposits buried by small dune migration, after an exposure time long enough to record a succession of different overprinted trackways.

Early Permian tetrapod-track assemblages

The Permian vertebrate body-fossil and footprint record of Europe and North America is abundant and well studied (e.g. Olson, 1989; Milner, 1993; Hunt et al., 1995; Tverdokhlebov et al., 2005; Minter, et al., 2007). In the last



FIGURE 7 | Reconstruction of the Permian Paganzo fluvio-eolian environment with the putative vertebrate and invertebrate trackmakers. Illustration by Boris Budiša.

decades several authors have proposed different schemes to group trace-fossil associations from the Early Permian (Haubold, 1996; Lucas, 1998). These associations were mainly grouped in two principal clusters, originally named as ichnofacies. First, the association of eolian facies dominated by *Chelichnus* and, less abundantly, *Dromopus* (Lockley et al., 1994; McKeever and Haubold, 1996), and second, the association of red-beds in fluvio-lacustrine facies, characterized mainly by *Batrachichnus*, *Limnopus*, *Gilmoreichnus*, and *Dimetropus* (Hunt et al., 1995). In Argentina, tracks from the Yacimiento Los Reyunos Formation were assigned to the typical *Chelichnus* association (Melchor, 2001), and those from the Carapacha Formation were compared to the “redbeds” association, with a late (early Late Permian) representation in South America (Melchor and Sarjeant, 2004). The material of the Patquía Formation reveals some resemblance to those that were referred to the *Chelichnus* ichnofacies by Lockley et al. (1994).

Ichnofacies

Nonmarine ichnofacies are now receiving more attention by ichnologists (*e.g.* Buatois and Mángano, 2004,

2007; Hunt and Lucas, 2007; Ekdale et al., 2007; Minter et al., 2007; Melchor et al. 2006). Three archetypal nonmarine ichnofacies are widely accepted: *Scoyenia* (Seilacher, 1964, 1967), *Mermia* (Buatois and Mángano, 1995), and *Coprinsphaera* (Genise et al., 2000). Additionally, the typically marine *Skolithos* ichnofacies is recognized in nonmarine settings (Buatois and Mángano, 2004). Various eolian ichnofacies have been proposed: *Octopodichnus*, *Chelichnus*, and *Entradichnus* ichnofacies (Hunt and Lucas, 2007; Ekdale et al., 2007). And, a new model of tetrapod ichnofacies was recently presented (Hunt and Lucas, 2007), suggesting five ichnofacies (*Grallator*, *Brontopodus*, *Batrachichnus*, *Characichnos*, and *Chelichnus* ichnofacies) which correspond with aspects of invertebrate ichnofacies.

There is a strong need for integration of vertebrate and invertebrate trace fossils in continental ichnofacies models. There are several ichnofacies models proposed for eolian environments, but these are either based on invertebrate or vertebrate trace fossils. Hunt and Lucas (2007) proposed the *Chelichnus* ichnofacies for tetrapod associations recurrent in dune facies of eolian environments (Fig. 6). These authors also erected the *Octopodichnus* ichnofacies for eolian trace-fossil associations of the Coconino Sandstone (Fig. 6), whilst Ekdale et al. (2007) defined the *Entradichnus* ichnofacies for eolian trace-fossil associations based on the Navajo Sandstone (Fig. 6). The close correspondence between the *Chelichnus* and *Octopodichnus* ichnofacies in dune environments was suggested by Hunt and Lucas (2007). In the same vein, Buatois and Mángano (2008) attempted to integrate eolian ichnofacies models, and combined both ichnofacies in the *Octopodichnus-Entradichnus* ichnofacies. The Patquía trace-fossil association 2 has aspects in common with the *Octopodichnus*, *Chelichnus*, and *Entradichnus* ichnofacies (Fig. 6). It particularly resembles the *Entradichnus* ichnofacies in containing trace fossils that have been produced by shallow burrowing, desert-dwelling arthropods, such as beetles and others insects, that kept pace with dune migration (Ekdale et al., 2007). Also, it shares the presence of arthropod trackways with the *Octopodichnus* ichnofacies, which consists exclusively of locomotion trace fossils (Hunt and Lucas, 2007). Finally, it also reveals some resemblance to the *Chelichnus* ichnofacies. The material described here demonstrates the need to integrate both invertebrate and vertebrate trace fossils in ichnofacies models.

Trace-fossil association 1 from the Patquía Formation developed in floodplain deposits and is ascribed to the *Scoyenia* ichnofacies (Fig. 6). It displays many of the archetypal characteristics of the *Scoyenia* ichnofacies, including the presence of small horizontal bilo-

bate trace fossils, such as *Cruziana* and *Rusophycus*. The *Scoyenia* ichnofacies occurs in low-energy settings periodically exposed to air or periodically inundated (Frey et al., 1984; Buatois and Mángano, 1995). In this case there is no clear evidence of subaerial exposure, although a progressive desiccation and stiffening of the substrate is probably indicated by the presence of striae in some *Rusophycus* specimens.

CONCLUSIONS

The vertebrate ichnofauna documented here represents the oldest record of tetrapods in Argentina. Floodplain deposits of the Patquía Formation record invertebrate trace fossils (trace-fossil association 1) that are an example of the *Scoyenia* ichnofacies displaying mostly arthropod trace fossil. Eolian sand-sheet deposits of the Patquía Formation record a second trace-fossil association that has aspects in common with the three proposed eolian ichnofacies, suggesting the need for revision and their potential integration. The *Octopodichnus-Entradichnus* ichnofacies (*sensu* Buatois and Mángano, 2008) seems to be characterized by horizontal (e.g., *Palaeophycus*, *Planolites*) and vertical burrows (e.g., *Skolithos*, *Digitichnus*), arthropod trackways (e.g., *Octopodichnus*, *Paleohelcura*), and vertebrate tracks chiefly produced by a non-mammalian synapsids and mammals (e.g. *Chelichnus*, *Brasilichnus*).

ACKNOWLEDGMENTS

This work is part of the Doctoral Thesis of V. Krapovickas, financially supported by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). We specially thank Nicholas Minter and Ricardo Melchor for their careful review and useful suggestions, and Luis Buatois and Gabriela Mángano for their helpful comments to improve the quality of the first version of this contribution. Field research was supported by the Natural Sciences and Engineering Research Council (NSERC) Discovery Grant (311727-05 and 08) to M. Gabriela Mángano (VK), and UBACyT-X133 and PICT 07-1902 to Claudia Marsicano (VK), PICT 32236 (AM), PIP CONICET 0535/98 (AA).

REFERENCES

- Aceñolaza, F.G., Buatois, L.A., 1991. Trazas fósiles del Paleozoico Superior continental Argentino. *Ameghiniana*, 28(1-2), 89-108.
- Aceñolaza, F.G., Buatois, L.A., 1993. Nonmarine perigondwanic trace fossils from the late Paleozoic of Argentina. *Ichnos*, 2, 183-201.
- Alpert, S.P., 1974. Systematic review of the genus *Skolithos*. *Journal of Paleontology*, 48, 661-669.
- Azcuy, C.L., Morelli, J.R., 1970. Geología de la Comarca de Paganzo - Amaná. El Grupo Paganzo, formaciones que lo componen y sus relaciones. *Revista de la Asociación Geológica Argentina*, 25(4), 405-429.
- Azcuy, C.L., Valencio, D.A., Vilas, J.F., 1979. Estratigrafía de la comarca de Amaná - Talampaya. Neuquén, 1978, VII Congreso Geológico Argentino, 1(Actas), 243-246.
- Bertling, M., Braddy, S.J., Bromley, R.G., Demathieu, G.R., Genise, J., Mikuláš, R., Nielsen, J.K., Nielsen, K.S.S., Rindsberg, A.K., Schlirf, M., Uchman, A., 2006. Names for trace fossils: a uniform approach. *Lethaia*, 39(3), 265-286.
- Bodenbender, G., 1911. Constitución geológica de la parte meridional de la provincia de La Rioja y regiones limítrofes. República Argentina. Córdoba, Boletín de la Academia Nacional de Ciencias, 19(1), 2-211.
- Bodenbender, G., 1912. Parte meridional de la provincia de La Rioja y regiones limítrofes. Constitución geológica y productos minerales. Buenos Aires, Anales del Ministerio de Agricultura de la Nación, Sección Geología, Mineralogía y Minería, 7(3), 1-161.
- Bown, T.M., 1982. Ichnofossils and rhizoliths of the near shore fluvial Jebel Qatrani Formation (Oligocene), Fayum Province, Egypt. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 40, 255-309.
- Bromley, R.G., Asgaard, U., 1972. Notes on Greenland trace fossils: freshwater *Cruziana* from the Upper Triassic of Jameson Land, east Greenland. *Grønlands Geologiske Undersøgelse, Rapport*, 49, 7-13.
- Bromley, R.G., Asgaard, U., 1979. Triassic freshwater ichno-coenoses from Carlsberg Fjord, east Greenland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 28, 39-80.
- Buatois, L.A., Mángano, M.G., 1995. The paleoenvironmental and paleoecological significance of the lacustrine *Mermia* ichnofacies: an archetypical subaqueous nonmarine trace fossil assemblage. *Ichnos*, 4, 151-161.
- Buatois, L.A., Mángano, M.G., 2004. Animal-substrate interactions in freshwater environments: applications of Ichnology in facies and sequence stratigraphic analysis of fluvio-lacustrine successions. In: McIlroy, D.M. (ed.). *The Application of Ichnology to Palaeoenvironmental and Stratigraphic Analysis*. London, Geological Society, 228(Special Publications), 311-333.
- Buatois, L.A., Mángano, M.G., 2007. Invertebrate ichnology of continental freshwater environments. In: Miller, W. III (ed.). *Trace Fossils*. Amsterdam, Elsevier, 285-323.
- Buatois, L.A., Mángano, M.G., 2008. Continental Ichnology: Conceptual Framework and Prospects. 33rd International Geological Congress. Oslo (Norway), Abstract in CD-ROM.
- Buatois, L.A., Jalfin, G., Aceñolaza, F.G., 1997. Permian non-marine invertebrate trace fossils from southern Patagonia, Argentina: Ichnologic signatures of substrate consolidation and colonization sequences. *Journal of Paleontology*, 71(2), 324-336.
- Buatois, L.A., Mángano, M.G., Aceñolaza, F.G., 1996. Ichnofaunas paleozoicas en sustratos firmes no marinos: evidencias del Pérmico de la Cuenca Paganzo. *Ameghiniana*, 33(3), 265-270.

- Caselli, A.T., 1998. Estratigrafía y Sedimentología de las Formaciones Patquía (Pérmico) y Talampaya (Triásico Inferior), en las Sierras Pampeanas Noroccidentales y Precordillera Central (Provincias De La Rioja Y San Juan). Doctoral Thesis. Universidad de Buenos Aires, 437pp.
- Caselli, A.T., Arcucci, A.B., 1999. Huellas de tetrápodos en eolianitas de la Formación Patquía, Cuenca Paganzo, Provincia de La Rioja. I Simposio Argentino del Paleozoico Superior. *Ameghiniana*, 36(4), 20.
- Caselli, A., Limarino, C.O., 2002. Sedimentología y evolución paleoambiental de la Formación Patquía (Pérmico) en el extremo sur de la sierra de Maz y cerro Bola, provincia de La Rioja, Argentina. *Revista de la Asociación Geológica Argentina*, 57(4), 415-436.
- Cei, R.L., Gargiulo, J., 1977. Icnitas de tetrápodos Pérmicos del sur de Mendoza. *Ameghiniana*, 14, 127-132.
- Cisterna, G.A., Simanaukas, T., 2000. Brachiopods from the Río del Peñón Formation, Río Blanco Basin, Upper Paleozoic of Argentina. *Revista Española de Paleontología*, 15(2), 129-151.
- Dawson, J.W., 1864. On the fossils of the genus *Rusophycus*. *Canadian Naturalist and Geologist*, 1, 363-367.
- Debriette, P., Gand, G., 1990. Stratigraphic and paleoenvironmental results from new paleontological observations made in the Permian of the western part of the Lodève basin (South Massif Central). *Géologie de la France*, 19-32.
- d'Orbigny, A., 1842. Voyage dans l'Amerique méridionale (le Bresil, la République argentine, le Patagonia, le République de Chile, le Républiiue de Bolivie, le République de Perou) executé pendant les anées 1826, 1827, 1828, 1829, 1830, 1831, 1832 et 1833. París, *Paleontologie*, 3(4), 1-188.
- Ekdale, A.A., Bromley, R.G., Loope, D.B. 2007. Ichnofacies of an ancient erg: A climatically influenced trace fossil association in the Jurassic Narvajo Sandstone, Southern Utah, USA. In: Miller, W. III (Ed.). *Trace Fossils: Concepts, Problems, Prospects*. Amsterdam, Elsevier, 562-574.
- Fornós, J.J., Bromley, R.G., Clemmensen, L.B., Rodríguez-Perea, A., 2002. Tracks and trackways of *Myotrachus balearicus* Bate (Artiodactyla, Caprinae) in Pleistocene aeolianites from Mallorca (Balearic Islands, Western Mediterranean). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 180, 277-313.
- Frenguelli, J., 1949. Acerca de un nuevo descubrimiento de plantas en los Estratos del Arroyo Totoral, en la sierra de Los Llanos, La Rioja. *Revista de la Asociación Geológica Argentina*, 4(3), 153-164.
- Frey, R.W., Pemberton, S.G., Fagerstron, J.A., 1984. Morphological, ethological and environmental significance of the ichnogenera *Scoyenia* and *Ancorichnus*. *Journal of Paleontology*, 58, 511-528.
- Gand, G., 1994. *Isopodichnus furcosus* nov. ichnosp. Ichnocoenoses from the Permian Lodève basin (Massif Central, France). *Geobios*, 27, 73-86.
- Genise, J.F., Mángano, M.G., Buatois, L.A., Laza, J., Verde, M., 2000. Insect trace fossil associations in paleosols: the *Coprinisphaera* ichnofacies. *Palaaios*, 15, 33-48.
- Geuna, S.E., Escosteguy, L.D., Limarino, C.O., 2010. Paleomagnetism of the Carboniferous-Permian Patquía Formation, Paganzo basin, Argentina: implications for the apparent polar wander path for South America and Gondwana during the Late Palaeozoic. *Geologica Acta*, 8(4), 373-397.
- Gierlowski-Kordesch, E., 1991. Ichnology of an ephemeral lacustrine alluvial plane system. Jurassic East Berlin Formation, Hartford Basin, U.S.A. *Ichnos*, 6, 221-232.
- Gilmore, C.W., 1926. Fossil footprints from the Grand Canyon. *Smithsonian Miscellaneous Collections*, 77(9), 1-41.
- Haldeman, S.S., 1840. Supplement to Number One of "A monograph of the Limniades, or fresh-water univalve shell of North America", containing descriptions of apparently new animals in different classes, and the name and characters of the subgenera in *Paludina* and *Anculosa*. Philadelphia, J. Dobson, 3pp.
- Hall, J., 1847. Palaeontology of New York. Volume I. Containing descriptions of the organic remains of the Lower Division of the New York System (Equivalent to the Lower Silurian rocks of Europe). Albany (New York), C. van Benthuyssen, 338pp.
- Hall, J., 1852. Palaeontology of New York, vol. 2. State of New York, Albany, 362pp.
- Häntzschel, W., 1975. Trace fossils and problematica. In: Teichert, C. (ed.). *Treatise on Invertebrate Paleontology*, W (Supplement 1). Lawrence (Kansas), University of Kansas Press, 1-269.
- Haubold, H., 1996. Ichnotaxonomie und Klassifikation von Tetrapodenfährten aus dem Perm. *Hallesches Jahrbuch für Geowissenschaften*, 18(B), 23-88.
- Haubold, H., Lucas, S.G., 2003. Tetrapod footprints of the Lower Permian Choza Formation at the Castle Peak, Texas. *Paläontologische Zeitschrift*, 77(2), 247-261.
- Hunt, A., Lockley, M., Lucas, S.G., Haubold, H., Braddy, S.J., 1995. Tetrapod ichnofacies in early Permian red beds of the American Southwest. In: Lucas, S.G., Heckert, A.B. (eds.). *Early Permian footprints and facies*. New Mexico Museum of Natural History and Science Bulletin, 6, 295-301.
- Hunt, A., Lucas, S.G., 2007. Tetrapod ichnofacies: a new paradigm. *Ichnos*, 14, 59-68.
- Jardine, W., 1850. Note to Mr. Harness's paper "On the position of the impressions of footsteps in the Bunter sandstones of Dumfriesshire". *Annals and Magazine of Natural History*, 6, 208-209.
- Keighley, D.G., Pickerill, R.K., 1996. Small *Cruziana*, *Rusophycus*, and related ichnotaxa from eastern Canada: the nomenclatural debate and systematic ichnology. *Ichnos*, 4, 261-285.
- Krapovickas, V., Mángano, M.G., Mancuso, A., Marsicano, C.A., Volkheimer, W., 2008. Ichnofaunas triásicas en abanicos aluviales distales: evidencias de la Formación Cerro Puntudo, cuenca cuyana, Argentina. *Ameghiniana*, 45(2), 463-472.
- Limarino, C.O., Césari, S.N., 1985. Primer registro paleoflorístico de la Formación La Colina (Paleozoico Superior), Cuenca Paganzo, República Argentina. *Universidade de São Paulo, Boletim Instituto de Geociências*, 15, 32-37.
- Limarino, C.O., Césari, S.N., Net, L.I., Marensi, S.A., Guitierrez, R.P., Tripaldi, A., 2002. The Upper Carboniferous postglacial

- transgression in the Paganzo and Río Blanco basins (north-western Argentina): facies and stratigraphic significance. *Journal of South American Earth Sciences*, 15, 445-460.
- Linck, O., 1942. Die Spur *Isopodichnus*. *Senckenbergiana*, 25, 232-255.
- Lockley, M.G., Hunt, A.P., Meyer, C., 1994. Vertebrate Tracks and the Ichnofacies Concept: Implications for Paleoecology and Palichnostratigraphy. In: Donovan, S. (ed.). *The Paleobiology of Trace Fossils*. New York, Wiley & Sons, 241-268.
- Lockley, M., 2007. A tale of two ichnologies: the different goal and potential of invertebrate and vertebrate (Tetrapod) ichnotaxonomy and how they relate to ichnofacies analysis. *Ichnos*, 14, 39-57.
- Loope, D.B., 2006. Dry-season tracks in dinosaur-triggered grainflows. *Palaaios*, 21, 132-142.
- Lucas, S.G., 1998. Toward a tetrapod biochronology of the Permian. In: Lucas, S.G., Estep, J.W., Hoffer, H. (eds.). *Permian Stratigraphy and Paleontology of the Robledo Mountains, New Mexico*. New Mexico Museum of Natural History and Science Bulletin, 12, 71-91.
- Mancuso, A.C., 2008. Tafonomía de peces y su relación con los paleoambientes de depositación: dos ejemplos de Argentina. Buenos Aires (Argentina), XII Reunión Argentina de Sedimentología, Abstract, 104.
- McKeever, P.J., Haubold, H., 1996. Reclassification of vertebrate trackways from the Permian of Scotland and related forms from Arizona and Germany. *Journal of Paleontology*, 70, 1011-1022.
- Melchor, R.N., 2000. Stratigraphic and biostratigraphic consequences of a new $^{40}\text{Ar}/^{39}\text{Ar}$ date for the base of the Cochicó Group (Permian), Eastern Permian Basin, San Rafael, Mendoza, Argentina. *Ameghiniana*, 37(3), 271-282.
- Melchor, R.N., 2001. Permian tetrapod footprints from Argentina. *Hallesches Jahrbuch für Geowissenschaften*, 23(B), 35-43.
- Melchor, R., Cesari, S., 1991. Algunos elementos paleoflorísticos de la Formación Carapacha (Pérmico Inferior), Prov. de La Pampa, Rep. Argentina. *Ameghiniana*, 28(3-4), 347-352.
- Melchor, R.N., Sarjeant, W.A.S., 2004. Small amphibian and reptile footprints from the Permian Carapacha Basin, Argentina. *Ichnos*, 11, 57-78.
- Melchor, R.N., Bedatou, E., De Valais, S., Genise, J.F., 2006. Lithofacies distribution of invertebrate and vertebrate trace-fossil assemblages in an Early Mesozoic ephemeral fluvio-lacustrine system from Argentina: implications for the Scoyenia ichnofacies. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 239, 253-285.
- Milner, A.R., 1993. The biogeography of salamanders in the Mesozoic and early Cenozoic: a cladistic-vicariance model. In: Sims, R.W., Price, J.H., Whalley, P.E.S. (eds.). *Evolution, Time and Space: The Emergence of the Biosphere*. London, Academic Press, 431-468.
- Minter, N.J., Krainer, K., Lucas, S.G., Braddy, S.J., Hunt, A.P., 2007. Palaeoecology of an Early Permian playa lake trace fossil assemblage from Castle Peak, Texas, USA. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 246, 390-423.
- Morrissey, L.B., 2006. The ichnology of terrestrial palaeoenvironments: Old Red Sandstone, British Isles. Doctoral Thesis. Bristol, University of the West of England, 399pp.
- Olson, E.C., 1989. Problems of Permo-Triassic terrestrial vertebrate extensions. *Historical Biology*, 2, 17-35.
- Osgood, R.G., 1970. Trace fossils of the Cincinnati area. *Palaeontographica Americana*, 6, 281-439.
- Pemberton, S.G., Frey, R.W. 1982. Trace fossil nomenclature and the *Planolites Palaeophycus* dilemma. *Journal of Paleontology*, 56, 843-881.
- Piñeiro, G., Marsicano, C., Lorenzo, N., 2007. A new temnospondyl from the Permo-Triassic Buena Vista Formation of Uruguay. *Palaeontology*, 50(3), 627-640.
- Pollard, J.E., 1981. A comparison between the Triassic trace fossils of Cheshire and south Germany. *Palaeontology*, 24, 555-588.
- Pollard, J.E., 1985. *Isopodichnus*, related arthropod trace fossils and notostracans from Triassic fluvial sediments. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 76, 273-285.
- Ratcliffe, B.C., Fagerstrom, J.A., 1980. Invertebrate lebensspuren of Holocene floodplains: their morphology, origin and palaeoecological significance. *Journal of Paleontology*, 54, 614-630.
- Richter, M., Langer, M., 1998. Fish remains from the Upper Permian Rio do Rastro Formation (Paraná Basin) of southern Brazil. *Journal of African Earth Sciences*, 27, 158-159.
- Sadler, C.J., 1993. Arthropod trace fossils from the Permian de Chelly Sandstone, Northeastern Arizona. *Journal of Paleontology*, 67(2), 240-249.
- Schindewolf, O.H., 1928. Studien aus dem Marburger Buntsandstein III-VI. *Senckenbergiana*, 10, 16-54.
- Schlirf, M., Uchman, A., Kümmel, M., 2001. Upper Triassic (Keuper) non-marine trace fossils from the Haßberge area (Franconia, southeastern Germany). *Paläontologische Zeitschrift*, 75, 71-96.
- Seilacher, A., 1955. Spuren und Lebensweise der Trilobiten. In: Schindewolf, O.H., Seilacher, A. (eds.). *Beiträge zur Kenntnis des Kambriums in der Salt Range (Pakistan)*. Abhandlungen der Mathematisch-Naturwissenschaftlichen Klasse, Mainz, Akademie der Wissenschaften und der Literatur, 10, 342-372.
- Seilacher, A., 1964. Biogenic sedimentary structures. In: Imbrie, J., Newell, N. (eds.). *Approaches to paleoecology*. New York, John Wiley and Sons, 296-316.
- Seilacher, A., 1967. Bathymetry of trace fossils. *Marine Geology*, 5, 413-428.
- Seilacher, A., 1970. *Cruziana* stratigraphy of "non-fossiliferous" Palaeozoic sandstones. In: Crimes, T.P., Harper, J.C. (eds.). *Trace fossils*. *Geological Journal*, 3(Special Issue), 447-476.
- Seilacher, A., 1985. Trilobite palaeobiology and substrate relationships. *Royal Society of Edinburgh Transactions: Earth Sciences*, 76, 231-237.
- Seilacher, A., Buatois, L.A., Mángano, M.G. 2005. Trace fossils in the Ediacaran-Cambrian transition: Behavioral Diversifi-

- cation, Ecological turnover and Environmental Shift. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 227, 323-356.
- Siano, C.A., 1990. Últimos eventos depositacionales de la Formación La Colina (Patquía), sierra de Los Llanos, La Rioja, Argentina. San Juan, 11° Congreso Geológico Argentino, 2(Actas), 187-191.
- Sinito, A.M., Valencia, D.A., Vilas, J.F., 1979. Palaeomagnetism of a sequence of Upper Palaeozoic-Lower Mesozoic red beds from Argentina. *Geophysical Journal of the Royal Astronomical Society*, 58, 237-247.
- Smith, N.D., Hein, F.J., 1971. Biogenic reworking of fluvial sediments by staphylinid beetles. *Journal of Sedimentary Petrology*, 41, 598-602.
- Spalletti, L.A., Limarino, C.O., Colombo Piñol, F., 2010. Internal anatomy of an erg sequence from the aeolian-fluvial system of the De La Cuesta Formation (Paganzo Basin, northwestern Argentina). *Geologica Acta*, 8(4), 431-447.
- Tedesco, A., Ciccioli, P., Suriano, J., Limarino, C., 2010. Changes in the architecture of fluvial deposits in the Paganzo Basin (Upper Paleozoic of San Juan province): an example of sea level and climatic controls on the development of coastal fluvial environments. *Geologica acta*, 8(4), 463-482.
- Thompson, R., Mitchel, J.C., 1972. Paleomagnetic and radiometric evidence for the age of the lower boundary of the Kiama magnetic interval. *London, Geophysics Journal*, 27, 207-214.
- Trewin, N.H., 1976. *Isopodichnus* in a trace fossil assemblage from the Old Red Sandstone. *Lethaia*, 9, 29-37.
- Tverdokhlebov, V.P., Tverdokhlebova, G.I., Minikh, A.V., Surkov, M.V., Benton, M.J., 2005. Upper Permian vertebrates and their sedimentological context in the South Urals, Russia. *Earth-Science Reviews*, 69, 27-77
- Valencio, D.A., 1972. International correlation of Late Paleozoic South American rocks on the basis of their magnetic remanences. *Anais de Academia Brasileira de Ciencias*, 44, 357-364.
- Valencio, D.A., Mitchell, J.G., 1972. Edad Potasio-Argón y paleomagnetismo de rocas igneas de las formaciones Quebrada del Pimiento y Las Cabras, Provincia de Mendoza. *Revista de la Asociación Geológica Argentina*, 27(2), 170-178.
- Valencio, D.A., Vilas, J.F. Mendia, J.E., 1977. Palaeomagnetism of a sequence of red beds of the middle and upper sections of Paganzo Group (Argentina) and the correlation of Upper Paleozoic-Lower Mesozoic rocks. *Geophysical Journal of the Royal Astronomical Society*, 51, 59-74.
- Zhang, G., Buatois, L.A., Mángano, M.G., Aceñolaza, F.G., 1998. Sedimentary facies and environmental Ichnology of a Permian playa-lake complex in western Argentina. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 138, 221-243.

Manuscript received June 2009;
revision accepted October 2009;
published Online July 2010.