
Lower Paleozoic Biostratigraphy of South America: New contributions and advances

G.L. ALBANESI and G. ORTEGA

CONICET - Museo de Paleontología, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba
CC 1598, 5000 Córdoba, Argentina. E-mail Albanesi: galbanes@com.uncor.edu E-mail Ortega: gcortega@arnet.com.ar

The early Paleozoic was the most critical time interval for the diversification of the biota on planet Earth. This phenomenon was driven by important paleocontinental and paleoceanographic changes, caused by intensive plate tectonic and climatic processes. The Cambrian explosion of life providing bauplans, the great Ordovician biodiversification event, the second most important mass extinction of life in the Phanerozoic that occurred in the Late Ordovician, and the appearance of land plants represent benchmarks in this evolutionary process (e.g., Albanesi et al., 2003; Webby et al., 2004; Munneke and Servais, 2007; Li et al., 2007)

The chronostratigraphic status of the lower Paleozoic, indispensable to correlate globally all of these important events, has changed dramatically in the last decade. The International Commission on Stratigraphy (ICS) of the International Union of Geological Sciences (IUGS), through the work of respective subcommissions for each time period, has made significant advances in establishing new global chronostratigraphic references. The Ordovician, Silurian and Devonian systems have been fully established and the Cambrian System is being rapidly reorganized in terms of stages, series and respective intra-systemic boundaries to accomplish the international time chart (see most recent advances at ICS-IUGS web site: www.stratigraphy.org)

This *Geologica Acta* issue is included in a series of them that have been published thanks to the editorial efforts carried out by research members of the CONICET who focus its activity on the lower Paleozoic

record (see Aceñolaza, 2003 and 2005, and cites herein). A number of contributions on the lower Paleozoic basin research, biostratigraphy and paleogeography have been gathered in these issues and makes up a significant deal of basic knowledge on this subject. The present issue consists of a series of selected papers that were presented as abstracts in the “9° Congreso Argentino de Paleontología y Bioestratigrafía” held in Córdoba, Argentina, on September 18-22, 2006. These contributions were invited papers from the total presented in the 2nd symposium of the congress, entitled “Bioestratigrafía del Paleozoico inferior” (i.e. Lower Paleozoic Biostratigraphy). Sixteen co-authors contributed six papers that compose this varied volume on lower Paleozoic paleontological and biostratigraphic subjects.

The paper by Bordonaro et al. (this issue) proposes a biostratigraphic model for the early Marjumian (Middle Cambrian) of the Argentine Precordillera involving trilobite biofacies and biozonation. The trilobite assemblages represent three biofacies that respond to segregated environmental conditions within an essentially synchronous interval that correspond to three biozones defined previously for the Middle Cambrian of western Laurentia.

Zeballo et al. (this issue) provide new paleontological data for the Eastern Cordillera of northwestern Argentina, in particular on conodont and graptolite faunas, which allow for the refinement of regional biostratigraphic schemes. The analysis of the conodont fauna reveals a succession of the conodont zones *Cordylodus angulatus*, *Paltodus deltifer*, and *Acodus deltatus-Parois-*

todus proteus. Graptolites of the *Aorograptus victoriae* Zone are linked with conodonts of the *Acodus deltatus-Paroistodus proteus* Zone, and trilobites of the *Noto-peltis orthometopa* Zone. The authors conclude that the Early Ordovician conodont fauna of northwestern Argentina represent a separate province in the southern South American margin of Gondwana.

An Early Ordovician (Floian) conodont fauna is reported for the first time from the Eastern Cordillera of Peru by Gutiérrez-Marco et al. (this issue). The conodont assemblage is referred to the upper part of the *Oepikodus evae* Zone and characterizes a South Gondwana Province of the Cold Domain in the Shallow-Sea Realm. Regarding previous biostratigraphic reports on the San José Formation, this conodont record in the study area indicates a highly diachronic lower boundary of the formation at a regional scale.

A conodont-graptolite high resolution biostratigraphy is presented by Ortega et al. (this issue) for Middle-Upper Ordovician siliciclastic facies of the Central Precordillera, Argentina. These records verify that the lower strata of the Sierra de La Invernada Formation are located to the east, contrary to conclusions of previous studies. The deposition of this unit developed as a response to sea level changes, which are also made evident by variations on the graptolite-conodont faunas along the revisited section. Six graptolite biozones and corresponding conodont zones are determined in this section, allowing for regional and global correlations of the upper Darriwilian, Sandbian and lower Katian stages (Middle-Upper Ordovician).

The lower Silurian Itacurubí Group exposed in the eastern Paraguay yielded an interesting graptolite fauna that is described and discussed firstly by Uriz et al. (this issue) The Llandovery monograptids of the Vargas Peña Formation from this group allows the correlation of units from the Eastern Cordillera and the Precordillera of Argentina. Regarding paleoenvironments, the association of graptolites and invertebrate macrofossils of the Vargas Peña Formation suggest an oxygen-rich environment in a shallow shelf.

Finally, an association of plants and palynomorphs from the Lower Devonian Santa Rosa Formation at Angosto de Alarache, Cinco Picachos Range in South Bolivia, is presented in the contribution by Di Pasquo and Noetinger (this issue). Taphonomic evidences from the early plant remains and preserved palynomorphs added to the absence of rooting systems allow interpreting the fossil remains as para-autochthonous to allochthonous, probably due to deposition under very low energy currents in shallow ponds, such as flooded alluvial plains or protected bays.

ACKNOWLEDGEMENTS

We thank the CONICET, ANPCyT (FONCyT), and Universidad Nacional de Córdoba, Argentina, for their continuing support on our research work, and especially to the contributors of this volume, who made it all possible. The editors are grateful to the colleagues and friends who kindly helped with the reviewing of the papers included in this *Geologica Acta* issue. These reviewers made important contributions by improving critically the final version of the papers. We thank all of them for this responsibility: Allison Palmer, Franco Tortello, Claudia Rubinstein, Dianne Edwards, Stig Bergström, Svend Stouge, Yong Yi Zhen, Juan Carlos Gutiérrez-Marco, Daniel Goldman, Edsel Brussa, Alfred Lenz, Anita Löfgren, Enrico Serpagli, and Christopher Barnes. We also acknowledge gratefully the editorial board of *Geologica Acta*, in particular the scientific editor Lluís Cabrera and the managing editor Angeles Santos, and also the whole managing scientific committee of the journal for supporting us with continuous professional assistance and enthusiasm throughout the editorial process of this publication.

REFERENCES

- Aceñolaza, G.F. (ed.), 2003. Advances in the knowledge of the Cambrian System. *Geologica Acta*, 1(1), 1-150.
- Aceñolaza, G.F. (ed.), 2005. Ordovician revisited: Reconstructing a unique period in Earth history. *Geologica Acta*, 3(4), 307-409.
- Albanesi, G.L., Beresi, M.S., Peralta, S.H. (eds.), 2003. Ordovician from the Andes. Proceedings of the 9th International Symposium on the Ordovician System, Serie Correlación Geológica 17, INSUGEO, Tucumán: 1-552.
- Bordonaro, O.L., Banchig, A.L., Pratt, B.R., Raviolo, M.M., 2008. Trilobite-based biostratigraphic model (biofacies and biozonation) for the Middle Cambrian carbonate platform of the Argentina Precordillera. *Geologica Acta*, 6(2), 115-129.
- Di Pasquo, M.M., Noetinger, S., 2008. First record of Early Devonian (Lochkovian) flora from the Santa Rosas Formation – Alarache, Southern Bolivia. *Geologica Acta*, 6(2), 191-210.
- Gutiérrez-Marco, J.C., Albanesi, G.L., Sarmiento, G.N., Carlotto, V., 2008. An Early Ordovician (Floian) Conodont Fauna from the Eastern Cordillera of Peru (Central Andean Basin). *Geologica Acta*, 6(2), 147-160.
- Li, J., Fan, J., Percival, I. (eds.), 2007. The global Ordovician and Silurian. Proceedings of the 10 International Symposium on the Ordovician System. *Acta Palaeontologica Sinica*, 46, 1-566.
- Munnecke, A., Servais, T. (eds.), 2007. Early Palaeozoic palaeogeography and palaeoclimate. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 245, 1-316.

Ortega, G., Albanesi, G.L., Banching, A.L., Peralta, G.L., 2008. High resolution conodont-graptolite biostratigraphy of the Mid-Late Ordovician in Sierra de La Invernada Formation (Central Precordillera, Argentina). *Geologica Acta*, 6(2), 161-180.

Uriz, N.J., Alfaro, M.B., Galeano Inchausti, F.C., 2008. Silurian (Llandovery) monograptids from the Vargas Formation (Parana Basin, Eastern Paraguay). *Geologica Acta*, 6(2), 181-190.

Webby, B.D., Paris, F., Droser, M.L., Percival, I.G. 2004. The great Ordovician biodiversification event. New York, Columbia University Press, 1-484.

Zeballo, F.J., Albanesi, G.L., Ortega, G., 2008. New late Tremadocian (Early Ordovician) conodont and graptolite records from the southern South American Gondwana margin (Eastern Cordillera, Argentina). *Geologica Acta*, 6(2), 131-145.



Guillermo L. Albanesi is a researcher of the National Council of Sciences (CONICET-CICTERRA) since 1999, and currently works at the Museum of Paleontology (FCEfYN, National University of Córdoba, Argentina). He finished his Geology graduation (1991) and PhD (1997) in the same university, holding from 1992 to 1998 a CONICET fellowship at the National Universities of Córdoba

and La Rioja, Argentina, and in 1999 in the University of Victoria, B.C., Canada (1999). He also held a Fulbright-Antorchas fellowship at the Ohio State University, USA from 2001 to 2002). From 1999 to 2008 Dr. Albanesi was the Secretary of the International Subcommittee on Ordovician Stratigraphy (ICS-IUGS) and in 2003 edited the proceedings of the 9th International Symposium on the Ordovician System held in San Juan, Argentina. He has developed extensive field studies in Argentina and published numerous papers and book chapters on conodont faunas and lower Paleozoic biostratigraphy. Director and participant in several research projects; his interest is mainly focused on conodont biostratigraphy, paleoenvironments, bioevents and paleothermometry of Paleozoic basins in South America.



Gladys Ortega is Geologist (1981) and Doctor in Geological Sciences (1987)

from the National University of Córdoba, Argentina. Since 1989 she became a researcher of the National Council of Sciences (CONICET) and worked at the Museum of Paleontology (FCEfYN, National University of Córdoba, Argentina). From 1995 to 2000 she performed as vice-director of the Museum of Paleontology. In 2003 Dr. Ortega chaired the 7th International Graptolite Conference held in San Juan, Argentina, and edited the volume of proceedings. Her research interest focuses mainly on Ordovician and Silurian graptolite faunas, having produced numerous contributions, which resulted from several projects that she led on lower Paleozoic biostratigraphy from South America. She continues currently her work in new projects on high resolution biostratigraphy, graptolite faunas and paleoenvironments from Paleozoic basins of Argentina.