

Isotopic characterization of Jurassic evaporites. Aconcagua-Neuquén Basin, Argentina

G.L. LO FORTE^{|1|} F. ORTÍ^{|2|} and L. ROSELL^{|2|}

^{|1|}UBA-CONICET. Departamento de Ciencias Geológicas, Facultad de Ciencias Exactas y Naturales
Pabellón II, Ciudad Universitaria. 1428 Ciudad Autónoma de Buenos Aires, Argentina. E-mail: galof@gl.fcen.uba.ar

^{|2|}Departament de Geoquímica, Petrologia i Prospecció Geològica, Universitat de Barcelona
Zona Universitària de Pedralbes, 08028 Barcelona, Spain. Ortí E-mail: f.ortí@ub.edu Rosell E-mail: lrosell@ub.edu

ABSTRACT

Isotopic analysis can be used to interpret the origin of evaporitic sediments. A preliminary isotopic study of strontium, oxygen and sulphur has been carried out in Ca-sulphate facies of Jurassic marine evaporites (Tábanos Formation and Auquilco Formation) outcropping in southern Mendoza, Aconcagua-Neuquén Basin (Argentina), as a part of a comprehensive sedimentologic study. The analysed sections are located at arroyo Las Leñitas, Cañada Ancha and arroyo Blanco. Sampled units include laminated, banded, and nodular lithofacies, made up of anhydrite, secondary gypsum and calcite. The mineralogy was studied by conventional petrographic analysis and X-ray diffraction. The $^{87}\text{Sr}/^{86}\text{Sr}$ ratio was obtained in six samples, with values ranging from 0.706793 to 0.706839, which match marine calcium-sulphate data of the same age. A similar conclusion may be derived from ten samples analysed for oxygen ($\delta^{18}\text{O}$) and sulphur ($\delta^{34}\text{S}$) isotopic composition: the obtained values are between +11.55‰ and +14.42‰, and between +17.25‰ and +18.48‰ respectively. The sedimentologic-stratigraphic evidence and the isotopic data both suggest a marine origin for the Tábanos and Auquilco evaporites, without an analytically detectable contribution of continental waters or hydrothermal solutions. The results also suggest that no isotope fractionation occurred during the primary gypsum-to-anhydrite-to-secondary gypsum transformations.

KEYWORDS | Marine evaporites. Isotopic composition. Jurassic. Aconcagua-Neuquén Basin. Argentina.

INTRODUCTION

Stable isotope geochemistry applied to evaporitic sediments may provide specific information about depositional and diagenetic conditions. Due to the large variability of environmental parameters (such as salinity, temperature, pH, Eh, etc.), common early diagenetic reactions (by the action of either allochthonous solutions or in

situ bacterial activity), and rock-solution interactions during later diagenesis, the study of stable isotopes is essential to properly interpret the sedimentary history of evaporite sediments.

The sedimentary infilling of the Aconcagua-Neuquén Basin (West-central Argentina, Mendoza province) is well known due to academic and industry research carried out

since the end of the 19th century. However, the detailed sedimentologic features of the evaporitic units, mainly represented by Ca-sulphates (anhydrite and secondary gypsum) and subordinate Na-chloride, are largely unknown. At present, there is some isotopic data on CaSO₄ samples from quarries located in the southern part of the basin (Brodtkorb et al., 1997, and references herein; Zappettini, 1999), in south Mendoza and Neuquén provinces. Recently, new results from outcrops and quarries located in southern Mendoza were presented by Lo Forte et al. (2003), and shortly after by Linares et al. (2003).

A preliminary isotopic study of strontium, oxygen, and sulphur was carried out on the Ca-sulphate evaporites of the marine Jurassic sequences cropping out in the Aconcagua-Neuquén Basin (Fig. 1). The aim of this paper is to characterize and discuss the isotopic composition of these sulphate deposits, as a part of a larger sedimentologic and stratigraphic study on the palaeogeographic evolution of these sequences.

GEOLOGICAL SETTING

Since Early Jurassic times the geodynamic evolution of the western border of southern South America has been closely related to an east-deepening subduction zone with variable rates of convergence and segmentation along latitude (Mpodozis and Ramos, 1989; Ramos, 1992). In this context, the Aconcagua-Neuquén Basin represents a back-arc depocenter generated by pericratonic extensional processes which started during the Middle Triassic (Ramos, 1992; Manceda and Figueroa, 1995) due to the fragmentation of Gondwanaland (Uliana et al., 1989). The Jurassic evaporative sequences were deposited during a thermal decay episode representing a broad epeiric sag, after a synrift event (Manceda and Figueroa, 1995).

The sediments of the Aconcagua-Neuquén Basin are siliciclastic, volcanoclastic, carbonatic and evaporitic deposits, up to 7000 metres in thickness. They are assigned to nine mesosequences controlled by tectono-eustatic sea level changes ranging from Triassic to Cenozoic (Legarreta and Uliana, 1996). During Late Jurassic the sedimentary accumulation was influenced by important changes in the average position of the shoreline and the main depositional trends were driven by early-middle Callovian emergence, late Callovian to early Oxfordian flooding, and late Oxfordian to early Kimmeridgian basin shallowing and desiccation (Legarreta, 1991; Legarreta and Uliana, 1996). The stratigraphic framework of the basin during Late Jurassic in Malargüe area is briefly summarised in Figure 2.

The Tábanos and Auquilco formations are basin-filling evaporitic deposits (up to 80 and 400 metres respec-




FIGURE 1 | Location map of the study area and simplified stratigraphic columns of the sampled sites. Key to columns: To: Tordillo Fm.; Au: Auquilco Fm.; LM: La Manga Fm.; Lo: Lotena Fm.; Ta: Tábanos Fm.; Ca: Calabozo Fm. Legend: 1) anhydrite; 2) secondary gypsum + carbonate; 3) carbonate + secondary gypsum; 4) limestones and lutites; 5) sandstones and conglomerates; 6) sampled level.

tively) that cover an extended area of hundred of kilometres across the basin (for detailed palaeogeography see Legarreta and Uliana, 1996). They are transgressive




FIGURE 2 | Late Jurassic stratigraphic framework of the basin at Malargüe area (modified from Legarreta and Uliana, 1996 and Legarreta and Uliana, 1999). VM: Vaca Muerta Fm.; To: Tordillo Fm.; Au: Auquilco Fm.; LM: La Manga Fm.; Ta: Tabanos Fm.; Ca: Calabozo Fm.; TE: Tres Esquinas Fm.

Legend: 1) basinal to slope clastics; 2) slope to inner shelf clastics; 3) nearshore to non-marine clastics; 4) shelf carbonates; 5) marine evaporites; 6) mud flat to playa lake shales and sandstones; 7) distal fluvial to eolian sandstones; 8) proximal fluvial sandstones and conglomerates.

deposits of a preexisting carbonate-platform palaeotopography (Calabozo Formation and La Manga Formation and equivalents, respectively; Fig. 2) after a sudden “Messinian-style” desiccation event (Legarreta, 1991; Legarreta and Uliana, 1996): the basal Tábanos evaporite is palaeogeographically confined to the Calabozo slope and basinal domains revealing that the onset of sulphate deposition occurred within an almost completely desiccated basin. The evaporite beds onlap the Calabozo carbonates and locally overstep older Callovian strata. These deposits represent a transgressive event under negative hydrologic balance (Legarreta, 1991; Legarreta and Uliana, 1996). The older Auquilco evaporite is areally confined to the depositional basin of the older Oxfordian carbonates (La Manga Formation) revealing a base level fall of at least 150m. These strata reflect an aggradational system that filled the deeper part of the desiccated basin, lapping on the former slope front. The younger Auquilco evaporite strata expanded updip just beyond the proximal edge of the Oxfordian La Manga carbonate wedge (Legarreta, 1991; Legarreta and Uliana, 1996). In several of the sampled profiles herein, the sudden transition from normal marine to hypersaline conditions is documented by shallow subtidal to intertidal carbonates and evaporites overlying hemipelagic sediments, as previously stated by Legarreta and Uliana (1996). The end of both evaporitic episodes resulted from a

major base level fall, evidenced by erosional incisions, that led to the establishment of an arid to fluvial environment; the subsequent fluvial to eolian deposits are assigned to the lower Lotena Formation (and equivalent units) and Tordillo Formation (and equivalent units), respectively.

By the end of the Eocene and during the Middle Miocene a fold-thrust belt was developed, which produced the present structure of the basin. This fold and thrust belt is characterized by a series of east-verging folds and related thrusts, which developed a complex thrust front with duplexes and underthrusts, bounding to the east with an area of gentle folds (Ramos, 1992; Manceda and Figueroa, 1995).

ANALYSED SAMPLES

The analysed samples belong to four outcrop and quarry sections located at arroyo Las Leñitas, Cañada Ancha and arroyo Blanco localities (Fig. 1). These sections were measured and surveyed at 1:100 and 1:200 scale, paying special attention to the lithologic composition, sedimentary structures, and bedding patterns.

The carbonate phase is mainly represented by calcite (LMC). Ca-sulphate phases are anhydrite and secondary

gypsum, the latter derived from the hydration of a precursor anhydrite phase as revealed by the petrographic study of the samples. Gypsum lithofacies include laminated, banded and micronodular secondary gypsum. Secondary gypsum crystalline fabrics are mostly represented by the alabastrine and megacrystalline varieties (Lo Forte, 2003). Although the original crystalline fabrics have been changed with the successive mineralogical transformations, the macroscopic structures (lithofacies) of the rocks remain little altered or unchanged because the anhydrite-to-secondary gypsum conversion is considered to be mainly an isovolumetric process.

The isotopic analyses were carried out in selected secondary gypsum samples; in all of them the mineralogical composition was controlled by conventional petrographic analysis and by X-ray diffraction (XRD).

ANALYTICAL PROCEDURES

Oxygen ($\delta^{18}\text{O}$) and sulphur ($\delta^{34}\text{S}$) isotopic compositions were analysed in ten samples of secondary gypsum following the procedures of Longinelli and Craig (1967) for oxygen and Robinson and Kusakabe (1975) for sulphur. In order to eliminate the CO_2 produced by the carbonate mineral matrix, samples were washed with 2N HCl. The CO_2 and SO_2 were analysed on a mass-spectrometer VG ISOGAS SIRA series 10 and VG ISOGAS SIRA series II respectively. Oxygen isotopic values are referred to the SMOW standard (Standard Mean Ocean Water; Craig, 1961) while sulphur isotopic values are referred to the CDT standard (Cañon Diablo Troilite; Jensen and Nakai, 1962). In all cases the analytical precision is $\pm 0.1\%$. Isotopic measurements were carried out at the Servicio General de Análisis de Isótopos Estables of the Universidad de Salamanca (Spain).

The $^{87}\text{Sr}/^{86}\text{Sr}$ ratios were determined in six gypsum samples at the Laboratorio de Geocronología y Geo-

química Isotópica of the Universidad Complutense de Madrid (Spain). Powdered samples were diluted and washed with 2.5 N HCl. Sr was separated and concentrated using a chromatographic column with DOWEX 50W*8 200/400 resin. Strontium isotopic analyses were performed on a TIMS VG SECTOR 54 mass-spectrometer. Values were corrected for the presence of ^{87}Rb and normalized to $^{86}\text{Sr}/^{88}\text{Sr} = 0.1194$. The value of measured isotopic standard referred to NBS-987 was $^{87}\text{Sr}/^{86}\text{Sr} = 0.71026 \pm 0.00002$ (2σ ; $n=7$). The analytic error in the $^{86}\text{Sr}/^{88}\text{Sr}$ ratio, referred to two standard deviations, is 0.01%.

ISOTOPIC GEOCHEMISTRY

The oxygen and the sulphur isotope compositions were analysed in three samples from the Tábanos Formation and seven samples from the Auquilco Formation; individual results and averages for each sample are summarised in Table 1. The oxygen and sulphur isotopic compositions are relatively uniform, averaging $+13.25\%$, with a $+11.55\%$ minimum value and a $+14.42\%$ maximum value for oxygen, and $+17.80\%$ in average, with a $+17.25\%$ minimum value and a $+18.48\%$ maximum value for sulphur. The comparison between the $\delta^{18}\text{O}$ and $\delta^{34}\text{S}$ values, shown in Figure 3, reflects this homogeneity without an evident data set clustering, even between both formations.

Six of these samples, one from the Tábanos Formation and five from the Auquilco Formation, were also analysed for strontium isotopes. The results gave an average of 0.706815 and minimum and maximum values ranging from 0.706793 to 0.706839; the detailed results for each sample are presented in Table 2. These isotopic values are also relatively uniform, as can be appreciated comparing the strontium isotopic ratios with the oxygen and sulphur isotopic compositions (Fig. 4).

TABLE 1 | Isotopic composition ($\delta^{18}\text{O}$, $\delta^{34}\text{S}$) of secondary gypsum samples

Sample	Fm.	$\delta^{34}\text{S} \text{‰}$	$\delta^{34}\text{S} \text{‰}$	Average $\delta^{34}\text{S} \text{‰}$	$\delta^{18}\text{O} \text{‰}$	$\delta^{18}\text{O} \text{‰}$	$\delta^{18}\text{O} \text{‰}$	Average $\delta^{18}\text{O} \text{‰}$
A1	Tab.	+18,28	+18,69	+18,48	+13,27	+13,68	+13,83	+13,59
A20	Auq.	+18,24	+18,51	+18,37	+14,27	+14,70	+14,30	+14,42
A*3	Tab.	+17,56	+17,84	+17,70	+13,13	+13,11	+13,03	+13,09
C1	Auq.	+18,16	+17,96	+18,06	+13,08	+13,15	N/V	+13,11
C11	Auq.	+17,23	+17,34	+17,28	+13,14	+13,12	+13,49	+13,25
C5	Auq.	+18,06	+18,16	+18,11	+11,47	+11,69	+11,55	+11,57
C7	Auq.	+17,14	+17,36	+17,25	+11,52	+11,27	+11,85	+11,55
D4	Auq.	+17,60	+17,00	+17,30	+14,06	+14,14	+14,16	+14,12
D6	Auq.	+17,43	+17,43	+17,43	+13,97	+13,64	+14,08	+13,90
E7	Tab.	+18,00	+18,01	+18,00	+13,84	+13,85	+13,95	+13,88




FIGURE 3 | $\delta^{34}\text{S}$ versus $\delta^{18}\text{O}$ plot of analysed gypsum samples. Open circles: Tábanos Fm; filled circles: Auquilco Fm. Dashed area represents Jurassic values of marine evaporites (from Claypool et al., 1980).

DISCUSSION

Since the 1970's considerable effort has been focused on the isotopic variation of marine sediments and seawater composition through time. Analyses of the $\delta^{18}\text{O}$, $\delta^{34}\text{S}$ and $^{87}\text{Sr}/^{86}\text{Sr}$ ratios in different marine carbonates and evaporites have shown a considerable variation for all three isotopes (Claypool et al., 1980; Burke et al., 1982).

Regarding the isotopic composition of the dissolved sulphate anion, sulphur values exhibit a greater variation than oxygen through geologic time due to its larger time-oscillation and to its low exchange ratio with

TABLE 2 | Strontium isotopic composition ($^{87}\text{Sr}/^{86}\text{Sr}$) of secondary gypsum samples

Sample	Formation	$^{87}\text{Sr}/^{86}\text{Sr}$
C1	Auq.	0,706825
C5	Auq.	0,706837
C7	Auq.	0,706798
C11	Auq.	0,706839
A*3	Tab.	0,706793
A20	Auq.	0,706794

seawater. Strontium isotope ratios are more sensitive than the sulphur isotopic composition to the incorporation of non-marine waters (hydrothermal or meteoric) to marine brines, since the strontium content is relatively low in marine brines but can be higher in non-marine waters. Thus, the contribution of small amounts of non-marine waters to restricted seawater bodies can alter the marine strontium isotopic signal and give higher values.

The relationships between the $\delta^{18}\text{O}$ and $\delta^{34}\text{S}$ values, shown in Figure 3 and Table 1, reflect homogeneity and clustering within the range of Jurassic evaporites. A number of Upper Jurassic formations analysed by Claypool et al. (1980; appendix) have $\delta^{34}\text{S}$ comprised between +15.4 and +17.5‰. Although the range of $\delta^{34}\text{S}$ values obtained in the present paper is slightly higher (between +17.25‰ and +18.48‰), it fits the Jurassic uncertainty area of the sulphur curve of Claypool et al. (1980) and the Jurassic values presented by Strauss (1997). Moreover, $\delta^{34}\text{S}$ values up to +19.2‰ have been reported in the literature (Utrilla et al., 1992) for marine anhydrites of Liassic age,




FIGURE 4 | A) $^{87}\text{Sr}/^{86}\text{Sr}$ versus $\delta^{34}\text{S}$, and B) $^{87}\text{Sr}/^{86}\text{Sr}$ versus $\delta^{18}\text{O}$ plots of analysed gypsum samples. Open circles: Tábanos Fm; filled circles: Auquilco Fm. Dashed areas represent Jurassic values of marine evaporites (from Claypool et al., 1980, and Burke et al., 1982).




FIGURE 5 | Geohistoric curves of marine $\delta^{18}\text{O}$ (A), $\delta^{34}\text{S}$ (B) and $^{87}\text{Sr}/^{86}\text{Sr}$ (C) variation through Phanerozoic times. The age (horizontal dotted lines) and the minimum and maximum values (vertical dotted lines) of the analysed gypsum samples are indicated. $\delta^{18}\text{O}$ and $\delta^{34}\text{S}$ curves after Claypool et al. (1980) and $^{87}\text{Sr}/^{86}\text{Sr}$ curve after Burke et al. (1982).

as also pointed out by Strauss (1997). Concerning the strontium isotope ratios, they also reflect certain homogeneity (Fig. 4 and Table 2) as well as a marine origin, as they are almost identical to the Upper Jurassic marine curve.

The comparison between the oxygen, sulphur, and strontium isotopic values obtained from our gypsum samples, and those values characteristic of Jurassic marine evaporites, is shown in Figure 5. The minimum and maximum values of $\delta^{18}\text{O}$, $\delta^{34}\text{S}$ and $^{87}\text{Sr}/^{86}\text{Sr}$ have been plotted on the geohistoric curves of Claypool et al. (1980) for oxygen and sulphur, and of Burke et al. (1982) for strontium ratio. These results (Fig. 5) are also in agreement with biostratigraphic constraints (Riccardi, 1983; Riccardi et al., 2000).

We conclude that the analysed samples of secondary gypsum retained their original isotopic (marine) signal throughout the complete diagenetic cycle of calcium sulphate.

Our results are in agreement with the isotopic $\delta^{34}\text{S}$ data and $^{87}\text{Sr}/^{86}\text{Sr}$ ratios reported by Brodtkorb et al. (1997) in the southern part of the basin (Neuquén province). Brodtkorb et al. (1997) analysed samples from secondary gypsum, anhydrite and celestite, and obtained values that were consistent with those of the Jurassic marine evaporites. Gypsum samples had $^{87}\text{Sr}/^{86}\text{Sr}$ values ranging from 0.70684 to 0.70728 and $\delta^{34}\text{S}$ mean values of +16‰ (Tábanos Formation) and +16.8‰ (Auquilco Formation). Recently, Linares et al. (2003) have contributed new $\delta^{34}\text{S}$ data for the Auquilco Formation, sampled at arroyo Negro near the Malargüe city, although in this case the $\delta^{34}\text{S}$ values show a slightly lower mean (+15.3‰).

CONCLUDING REMARKS

The sulphur isotopic composition of the Ca-sulphate facies of the Jurassic Tábanos and Auquilco evaporitic sequences cropping-out in the Aconcagua-Neuquén Basin shows mean values of +13.25‰ for $\delta^{18}\text{O}$ and +17.80‰ for $\delta^{34}\text{S}$, all of them consistent with those of the Jurassic marine evaporites, according to published data (Claypool et al., 1980). Likewise, the $^{87}\text{Sr}/^{86}\text{Sr}$ ratios, ranging from 0.706793 to 0.706839, depict a good correspondence between our values and those corresponding to the Jurassic marine evaporites in the geohistoric curve of Burke et al. (1982). Despite their secondary character (i.e. the provenance from a precursor anhydritic phase), the analysed gypsum samples are valuable material for isotopic studies, because they have retained the original marine isotopic signal apparently unaffected by diagenesis. These results also suggest the absence of a significant contribution of continental waters or hydrothermal solutions, at least in an analytically detectable proportion, in the studied area of the basin.

ACKNOWLEDGEMENTS

This study has been supported by H.O.L.F., by the project DGI BTE2001-3201 (Spain), and by a FOMEC (UBA, Argentina) fellowship to G. Lo Forte during 2001. C.G. Asato provided field assistance and L. Legarreta help with basin stratigraphy. The authors also like to thank M.K. Brodtkorb (UBA, Argentina) for bibliography and J. Illa (UB, Spain) for preparing the petrographic collection. Revision supplied by T.K. Lowenstein and J.M. Rouchy provided helpful comments which improved the manuscript.

REFERENCES

- Brodtkorb, M.K., Barbieri, M., Zák, K., Hladiková, J., Tassinari, C., Ametrano, S., Etcheverry, R., de Barrio, R.E., Del Blanco, M., Danieli, J.C., 1997. Isotope data on barite and celestite deposits relate to three Mesozoic evaporitic sequences of the Neuquén Basin, Argentina. *International Geology Review*, 39, 307-316.
- Burke, W.H., Denison, R.E., Hetherington, E.A., Koepnick, R.B., Nelson, H.F., Otto, J.B., 1982. Variation of seawater $^{87}\text{Sr}/^{86}\text{Sr}$ throughout Phanerozoic time. *Geology*, 10, 516-519.
- Claypool, G.E., Holser, W.T., Kaplan, Y.R., Sakai, H., Zak, I., 1980. The age of sulfur and oxygen isotopes in marine sulfate and their mutual interpretation. *Chemical Geology*, 28, 199-260.
- Craig, H., 1961. Standard for reporting concentrations of deuterium and oxygen 18 in natural waters. *Science*, 133, 1833-1834.
- Jensen, M.L., Nakai, N., 1962. Sulfur isotope meteorite standards, results, and recommendations. In: Jensen, M.L. (ed.). *Biogeochemistry of sulfur isotopes*. New Haven, Yale University Press, 30-35.
- Legarreta, L., 1991. Evolution of a Callovian-Oxfordian carbonate margin in the Neuquén basin west-central Argentina: facies, architecture, depositional sequences and global sea-level changes. *Sedimentary Geology*, 70, 209-240.
- Legarreta, L., Uliana, M.A., 1996. The Jurassic succession in west-central Argentina: stratal patterns, sequences and paleogeographic evolution. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 120, 303-330.
- Legarreta, L., Uliana, M.A., 1999. El Jurásico y Cretácico de la Cordillera Principal y la Cuenca Neuquina. In: Caminos, R. (ed.). *Geología Argentina*. Buenos Aires, SEGEMAR, Anales, 29, 399-432.
- Linares, E., Armella, C., Cabaleri, N., Ostera, H., Haller, M., Iglesias, V., 2003. Sulfur isotopes in evaporites of the Neuquén basin, southern Mendoza, Argentina. IV South American Symposium on Isotope Geology. Short Papers 1, 361-363.
- Lo Forte, G.L., 2003. Evolución paleambiental y diagenética de la Formación Auquilco en el depocentro Atuel-Valenciana, Cuenca Neuquino-Aconcagiina. Primer Simposio Argentino del Jurásico. *Ameghiniana*, 40(4R), 43-44.
- Lo Forte, G.L., Ortí, F., Rosell, L., 2003. Análisis isotópico de las evaporitas jurásicas aflorantes en el sector surmendocino de la cuenca Neuquino-Aconcagiina. Primer Simposio Argentino del Jurásico. *Ameghiniana*, 40(4R), 44.
- Longinelli, A., Craig, H., 1967. Oxygen-18 variations in sulfate ions in sea water and saline lakes. *Science*, 156, 1431-1438.
- Maceda, R., Figueroa, D., 1995. Inversion of the Mesozoic Neuquén rift in the Malargüe fold-thrust belt, Mendoza, Argentina. In: Tankard, A.T., Suárez, R., Welsink, H.J. (eds.). *Petroleum basins of south America*. Tulsa, American Association of Petroleum Geologists, Memoir, 62, 369-382.
- Mpodozis, C., Ramos, V.A., 1989. The Andes of Chile and Argentina. In: Ericksen, G.E., Cañas Pinochet, M.T., Reine-mud, J.A. (eds.). *Geology of the Andes and its relation to Hydrocarbon and Mineral Resources*. Circun Pacific Council for Energy and Mineral Resources, Houston, Earth Sciences Series, 11, 59-90.
- Ramos, V.A., 1992. Control geotectónico de las cuencas triásicas de Cuyo. *Boletín de Informaciones Petroleras*. Tercera época, 9, 2-9.
- Riccardi, A.C., 1983. The Jurassic of Argentina and Chile. In: Moullade, M., Nairn, A.E.M. (eds.). *The Phanerozoic Geology of the World, II, Mesozoic B*, Amsterdam, Elsevier, 201-263.
- Riccardi, A.C., Leanza, H., Damborenea, S., Manceñido, M., Ballent, S., Zeiss, A., 2000. Marine Mesozoic Biostratigraphy of the Neuquén Basin. In: Miller, H., Hervé, F. (eds.). *Zeitschrift für Angewandte Geologie*. Hannover, 31st International Geological Congress, SH 1, 103-108.
- Robinson, B.E., Kusakabe, M., 1975. Quantitative preparation of sulfur dioxide, for $^{34}\text{S}/^{32}\text{S}$ analysis, from sulfides by combustion with cuprous oxide. *Analytical Chemistry*, 47, 1179-1181.
- Strauss, H., 1997. The isotopic composition of sedimentary sulfur through time. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 132, 97-118.
- Uliana, M., Biddle, K.T., Cerdán, J., 1989. Mesozoic extension and the formation of Argentine sedimentary basin. In: Tankard, A.J., Balkwill, H.R. (eds.). *Extensional tectonics and stratigraphy of the north Atlantic margins*, Tulsa, American Association of Petroleum Geologist, Memoir 46, 509-604.
- Utrilla, R., Pierre, C., Ortí, F., Pueyo, J.J., 1992. Oxygen and sulphur isotope compositions as indicators of the origin of Mesozoic and Cenozoic evaporites from Spain. *Chemical Geology (Isotope Geoscience Section)*, 102, 229-244.
- Zappettini, E. (ed.), 1999. *Recursos Minerales de la República Argentina II*. Buenos Aires, SEGEMAR, Anales, 35, 1150 pp.

Manuscript received October 2003;
revision accepted July 2004.