

University of Massachusetts Amherst
ScholarWorks@UMass Amherst

Cranberry Station Extension meetings

Cranberry Station Outreach and Public Service
Activities

2010

2010 Cranberry Management Update: New Planting Pest Management Techniques

Carolyn J. DeMoranville

University of Massachusetts - Amherst, carolynd@umext.umass.edu

Frank Caruso

UMass Cranberry Station, fcarus@umext.umass.edu

Hilary A. Sandler

University of Massachusetts - Amherst, hsandler@umass.edu

Anne L. Averill

University of Massachusetts - Amherst, averill@eco.umass.edu

Martha Sylvania

UMass Cranberry Station, martys@umext.umass.edu

Follow this and additional works at: https://scholarworks.umass.edu/cranberry_extension

 Part of the [Horticulture Commons](#)

Recommended Citation

DeMoranville, Carolyn J.; Caruso, Frank; Sandler, Hilary A.; Averill, Anne L.; and Sylvania, Martha, "2010 Cranberry Management Update: New Planting Pest Management Techniques" (2010). *Cranberry Station Extension meetings*. 95.

Retrieved from https://scholarworks.umass.edu/cranberry_extension/95

This Article is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Station Extension meetings by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.



New Planting Management Tips

Hilary Sandler, Anne Averill, Frank Caruso,
Carolyn DeMoranville, Marty Sylvia
UMass Amherst Cranberry Station




Nutrient Management

- Roots take up nutrients
 - Plugs can be fertilized right away but...
 - May look dormant first 2-3 weeks
 - Cuttings: all slow-release after a week or wait ~3 weeks
- Use slow release N
- Limit use of complete N-P-K
- Do not use high P materials.
 - Use 1N to 1P or less than 1P



Nutrient Management

- Other than slow release, use low rates every 2-3 weeks.
- Total N rate depends on materials used.



Weed Management

- Probably one of most imp't tasks to ensure good colonization!
- Vines should be as weed-free as possible.
- Hand-weeding - easier after watering.
 - Walking the bog is good opportunity to look for other pest issues.



Weed Management

- PRE herbicides once roots established.
 - Devrinol (up to 18 lb/A; multiple applications permitted)
 - Evital – some growers using this.
- POST herbicides – Callisto low-end rates.
 - Poast / Select: OK, but only control grasses.
- No Casoron until planting well-established.



Disease Management

- Prevent buildup of fungi that are causal agents of fruit rot, root rot, upright dieback.
- Several fungi (*Phyllosticta*, *Phomopsis*, *Fusicoccum*) can cause leaf drop & URD in addition to fruit rot – several diseases will be managed.
- Controlling leaf spot/drop will allow vines to colonize the bed more quickly.



Disease Management

- Scout for leaf spot/drop.
- Apply fungicide to entire bed once symptoms are first noticed during planting year.
- Rotate chlorothalonil, fenbuconazole, mancozeb, azoxystrobin at 10-day intervals, using no more than 5 applications.
- Light sanding between 1st & 2nd seasons buries inoculum & anchors runners.



Disease Management

- Initially, poor drainage will not be an issue for Phytophthora root rot, but low spots may eventually develop.
- Ridomil should be used during 1st season if root rot has been an issue in the past.
- Phostrol, Prophyt can be effectively used in subsequent seasons.



Insect Management

EVEN ON A NEW PLANTING...

- Keep an eye on the bog.
- Especially in the spring and summer
 - Sweep net intermittently
 - Walk the bog



Insect Management

EVEN ON A NEW PLANTING...

WATCH FOR VORACIOUS FOLIAGE FEEDERS

- Black-headed fireworm
 - And even yellow-headed fireworm
- Sparganothis fruitworm
- These foliage feeders can do significant damage in a short period!!



Insect Management

REMEMBER, new varieties are different...

- Larger bodied fruit are more attractive.
- CFW could move in by 2nd and 3rd year.

When ready to manage CFW

- Use % out of bloom.
- New planting and new varieties are earlier.
- Berries size up faster.

