University of Massachusetts Amherst ScholarWorks@UMass Amherst

Cranberry Station Extension meetings

Cranberry Station Outreach and Public Service Activities

2010

Pesticide Safety 2010 - Plant Disease Predictions for the 2010 Growing Season

Frank Caruso UMass Cranberry Station, fcaruso@umext.umass.edu

Follow this and additional works at: https://scholarworks.umass.edu/cranberry_extension Part of the Horticulture Commons



Recommended Citation

Caruso, Frank, "Pesticide Safety 2010 - Plant Disease Predictions for the 2010 Growing Season" (2010). Cranberry Station Extension meetings. 99.

Retrieved from https://scholarworks.umass.edu/cranberry_extension/99

This Article is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Station Extension meetings by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

Plant disease predictions for the 2010 growing season

Frank L. Caruso
UMass Cranberry Station

Diseases to mention

- Fruit rot
- Upright dieback
- Fairy ring
- Phytophthora root rot
- Other minor diseases



Fruit rot

- Wasn't bad in 2009 mostly in Ben Lear, Stevens
- Sanding should help in reducing inoculum for infection this year
- Preliminary Keeping Quality Forecast
 - possibly 5/10 points = good
 - possibly 7/10 points = very good to excellent (waiting for sunshine hours)

Fruit rot fungicide comments

- One new registration, but not in the Chart Book
- Will be testing it and one other new active ingredient this summer
- In order of efficacy (best to worst): chlorothalonil, EBDC's, fenbuconazole, azoxystrobin, ferbam, coppers



Upright dieback

- Higher incidence than usual in 2009, particularly in Early Black
- Some beds had 20% affected uprights, although more vegetative uprights were affected
- No prolonged periods of drought stress in 2009, so should not be much UD present in 2010
- If you have upright dieback this summer, CONTACT ME!



Fairy ring

- Increasing in incidence and severity, as more Ben Lear and Stevens beds come into production
- Causal agent has been identified in both MA and NJ
- May treatments (Abound, Indar) are more effective than June treatments (Ferbam)
- Should run sprinklers before and after application for 30 minutes
- Susceptibility of new hybrid cultivars?

Plans for 2010

- Collaborating with Peter Oudemans and Jim Polashock, Rutgers on the complicated life cycle of the causal agent which includes a rust on several weed species
- Will evaluate several unregistered sterol inhibitor fungicides for disease control
- Eventually hope to easily culture the fungus to help identify the problem, as opposed to drought stress, etc.
- Presence of fungal infection structures on the affected tissue makes diagnosis easier







Phytophthora root rot

- Disease is prominently present, especially in poorly-drained beds
- Many beds have been flooded or presently have standing water
- When renovating a bed that has had root rot, you MUST treat the soil with Basamid or the disease will quickly return
- Ridomil, Aliette, Phostrol are all very effective fungicides, but you should improve the drainage first

Other minor diseases

- Who knows what we'll see in 2010?
- Pyrenobotrys leaf spot
- Protoventuria leaf spot
- Red shoot
- Red leaf spot
- Call me if you have something worth a look!



Protoventuria leaf spot



Healthy

Red Shoot



Red leaf spot



Questions on cranberry diseases???