

University of Massachusetts Amherst  
**ScholarWorks@UMass Amherst**

---

Cranberry Chart Book - Management Guide

Cranberry Station Outreach and Public Service  
Activities

---

2007

## 2007 Chart Book: Insect Management At-A-Glance

Anne L. Averill

*University of Massachusetts Amherst Cranberry Station, [averill@eco.umass.edu](mailto:averill@eco.umass.edu)*

Martha M. Sylvia

*University of Massachusetts Amherst Cranberry Station, [martys@umass.edu](mailto:martys@umass.edu)*

Follow this and additional works at: <https://scholarworks.umass.edu/cranchart>

 Part of the [Agriculture Commons](#), and the [Plant Sciences Commons](#)

---

Averill, Anne L. and Sylvia, Martha M., "2007 Chart Book: Insect Management At-A-Glance" (2007). *Cranberry Chart Book - Management Guide*. 78.

Retrieved from <https://scholarworks.umass.edu/cranchart/78>

This Public Service and Outreach is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Chart Book - Management Guide by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

## INSECT MANAGEMENT 2007 AT A GLANCE

Prepared by Anne L. Averill and Martha M. Sylvia

**University of Massachusetts, Cranberry Station  
PO Box 569, East Wareham, MA 02538**

### *ALWAYS READ THE LABEL and REFER TO 2007 CRANBERRY CHART BOOK*

PRACTICE	RATE/A	INSECT TARGETED	NOTES
Actara*	2-4 oz	cranberry weevil, flea beetle	Do not exceed 8 oz per season, low rate works well, beware bee toxicity.
Admire Pro Admire 2F, Alias 2F	7-14 oz 16-32 oz	striped colaspis, oriental beetle	Best against just hatched larvae. Use drench Use drench application.
<i>Bacillus thuringiensis</i> based products		depending on product: gypsy moth, armyworms cutworms, spanworms	Best against early stage larvae, coverage must be thorough, combination with insecticides may control larger caterpillars.
Intrepid 2F* Confirm 2F	10-16 oz 16 oz	blackheaded fireworm, cutworms ** <i>Sparganothis</i> fruitworm gypsy moth, spanworms Cranberry fruitworm**	Insect growth regulators. Intrepid is more active and has greater residual than Confirm. However, Intrepid is restricted use and Zone II restricted.
Diazinon 50 W* Diazinon AG500* Diazinon AG600* & 5 WP	4-6 lb 4-6 pt 51-76.5 oz	fireworms, cranberry fruitworm cutworms**, flea beetle** striped colaspis**	<i>For all diazinon formulations:</i> <i>Highly toxic to birds.</i> Hold water at least 3 days, 7 day PHI, 24 hr/5 day REI, 3, 4 or 6 apps depending on product and rate.
Diazinon 14G*	21 lb	cranberry girdler	SLN 2007 + 2008, limited availability.
Flood, fall		cranberry girdler black vine & strawberry root weevil cranberry fruitworm	Sept. 20-30, hold 6 days no later than Sept. 25. 10-14 days after harvest; best if water is warm. Sept. 20-30, 3-4 weeks.
Flood, late water		CB fruitworm, southern red mite gypsy moth and cutworms	April 15-20, hold for 30 days.
Flood, summer		cranberry root & white grub	<i>Loss of crop</i> , May 12-July 20.
Imidan 70W	1.33-4 lbs	fireworms, CB fruitworm	Little or varied efficacy data available in MA.
Lorsban 4E*, Nufos 4E* & Hatchet*, Chlorpyrifos 4E AG* Lorsban 75 WG*	3 pt 2 lbs	cranberry weevil, cutworms fireworms, spanworms, <i>Sparganothis</i> & CB fruitworms	<i>Extremely toxic to fish.</i> Hold water at least 5 days, use only 2 times/season, 60 day PHI, 24 hr REI. Most populations of weevil and <i>Sparganothis</i> are resistant to Lorsban.
Nematodes		black vine & strawberry root weevil cranberry girdler	Apply in early evening in May and/or Sept. Apply 2 weeks after end of moth flight.
Orthene 97, Acephate 97UP Orthene 75S Acephate 75 WSP	1 lb 1 1/3 lb 1 1/3 lb	gypsy moth, fireworms, spanworms, cutworms,	Only 1 app per season, 90 day PHI, beware bee toxicity. 24 hr REI
Nexter	4.4-7.0 oz	southern red mite	Low-end rates usually gives excellent control, 2 apps/season, no aerial application.
Pyronyl or pyrenone	up to 12 oz	spanworms	
Sevin XLR Plus Sevin 4F & Carbaryl 4L Sevin 80 WSP & 80S	1.5-2 qt 1.5-2 qt 1.88-2.5 lb	fireworms, cutworms, gypsy moth, CB fruitworm CB flea beetle, striped colaspis	Avoid application within 10 days of start of bloom, 7 day PHI, no more than 5 app/season, 7 day application interval.
SpinTor 2SC (Spinosad)	4-10 oz	<i>Sparganothis</i> fruitworm, cutworms blackheaded fireworm, spanworms	Do not exceed 29 oz/season. Use lower rate only with good systems (6 min or better).

\* Restricted Use    \*\* FIFRA 2ee recommendation