



University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): Jon F. Fallaha, Brendan B. Spooner and Gavin D. Perkins

Article Title: Does Dual Operator CPR help minimize interruptions in chest compressions?

Year of publication: 2009

Link to published article:

<http://dx.doi.org/10.1016/j.resuscitation.2009.04.048>

Publisher statement: NOTICE: this is the author's version of a work that was accepted for publication in Resuscitation. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Resuscitation, Vol. 82, Issue 9, (September 2009), doi:10.1016/j.resuscitation.2009.04.048

1 **Does Dual Operator CPR help minimize interruptions in chest compressions?**

2 Jon F. Fallaha¹, Brendan B. Spooner¹, Gavin D. Perkins².

3 ¹The Medical School, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK.

4 ²University of Warwick Medical School, University of Warwick, Coventry, CV4 7AL,
5 UK

6

7 **Corresponding author**

8 Gavin D Perkins

9 Tel: +44 (0) 121 424 2966

10 E-mail: g.d.perkins@warwick.ac.uk

11

12 **Keywords**

13 Basic Life Support (BLS);

14 Chest Compression;

15 Training;

16 Bystander CPR;

17 Adult;

18 Cardiopulmonary resuscitation

19

20 **Word count**

21 2167

22

23

24 **Abstract**

25 Aims: Basic Life Support Guidelines 2005 emphasise the importance of reducing
26 interruptions in chest compressions (no-flow duration) yet at the same time stopped
27 recommending Dual operator CPR. Dual Operator CPR (where one rescuer does
28 ventilations and one chest compressions) could potentially minimize no-flow duration
29 compared to Single Operator CPR. This study aims to determine if Dual Operator CPR
30 reduces no-flow duration compared to Single Operator CPR.

31

32 Methodology: This was a prospective randomized controlled crossover trial. Medical
33 students were randomised into 'Dual Operator' or 'Single Operator' CPR groups. Both
34 groups performed 4 minutes of CPR according to their group allocation on a resuscitation
35 manikin before crossing over to perform the other technique one week later.

36

37 Results: Fifty participants were recruited. Dual Operator CPR achieved slightly lower
38 no-flow durations than the Single Operator CPR (28.5%(S.D. = 3.7) versus 31.6%(S.D. =
39 3.6), $P < 0.001$). Dual Operator CPR was associated with slightly more rescue breaths
40 per minute (4.9 (S.D. = 0.5) versus 4.5(S.D. = 0.5), $P = 0.009$). There was no difference in
41 compression depth, compression rate, duty cycle, rescue breath flow rate or rescue breath
42 volume.

43 Conclusions: Dual Operator CPR with a compression to ventilation rate of 30: 2 provides
44 marginal improvement in no-flow duration but CPR quality is otherwise equivalent to
45 Single Operator CPR. There seems little advantage to adding teaching on Dual Operator
46 CPR to lay / trained first responder CPR programs.

47 **Introduction**

48 Several factors may affect the quality of CPR. Performance of chest compressions is
49 known to be tiring. It has been shown that rescuer fatigue occurs within 1 minute of
50 chest compressions and that fatigue results in less efficacious chest compressions.¹⁻³
51 Hightower et al.¹ found that the percentage of adequate compressions decreased from
52 93% to 39% after 3 minutes and only 18% were satisfactory after 5 minutes.
53 Performance of CPR by a single rescuer may also be affected by pauses between
54 compressions (also known as the no-flow duration). Interruption in chest compressions
55 reduce coronary perfusion pressure and the chance of successful defibrillation.

56

57 Guidelines 2005 increased the compression to ventilation ratio from 15:2 to 30:2 for dual
58 operator CPR with the aim of reducing interruptions. In addition, if two rescuers are
59 present, then the problems of fatigue and increased no-flow duration can be minimized.
60 If the tiring compressions are shared between two or more rescuers then fatigue may be
61 less of an issue. Furthermore, if one rescuer can perform chest compressions
62 immediately after the other performs rescue breaths then it is possible to reduce the 'no-
63 flow duration'. Although not the primary aim of a study in 2004, Handley and Handley
64 suggested that 2 person CPR was more efficient than Single Operator CPR in avoiding
65 long pauses for ventilation,⁴ effectively resulting in a reduced 'no-flow duration'.

66

67 European Resuscitation Council (ERC) Guidelines in 2000⁵ and 2005⁶ offer different
68 suggestions on how to approach the situation if 2 rescuers are present. The former ERC
69 Guidelines 2000 suggest that when two or more trained rescuers are present they should

70 perform Dual Operator CPR. This is when one rescuer performs chest compressions
71 whilst the other performs rescue breaths, swapping if required (see Figure 1). ERC
72 Guidelines 2005 recommend that when two or more rescuers are present Single Operator
73 CPR should be performed, where one rescuer swaps with the other every 1-2 minutes.
74 Both sets of guidelines stress that any changeover of rescuers should be undertaken with
75 a minimum of delay. However, despite changing the guidelines between 2000 and 2005
76 there seems to have been no published research comparing the two methods. There have
77 been studies examining how best to provide advanced resuscitation with multiple
78 professional providers; in the pre-hospital setting⁷ and the in-hospital advanced life
79 support setting⁸, but none of these have focused on the provision of basic life support
80 alone or examined the effect on interruptions in chest compressions.

81

82 The aim of the present study was to determine if Dual Operator CPR, performed by a
83 team of trained first responders reduces no-flow duration compared to Single Operator
84 CPR with two rescuers.

85

86 **Material and Methods**

87 **Setting**

88 The study was carried out at the University of Birmingham Medical School, UK. We
89 recruited 58 Basic Life Support (BLS) instructors that taught BLS on a peer-led BLS and
90 Automated External Defibrillator (AED) course Ethical approval was granted by the
91 South Birmingham Research Ethics Committee. Verbal consent was obtained from the
92 candidates.

93

94 **Study design**

95 The present study was a randomised controlled crossover trial. All candidates were
96 randomly allocated to work in pairs. The pairs were then randomised into a ‘Single
97 Operator’ group (n=24) and a ‘Dual Operator’ group (n=26) using the random number
98 generation feature of Microsoft® Excel.

99

100 The pairs allocated to the Single Operator group were instructed to complete 4 minutes of
101 CPR by performing Single Operator CPR; swapping CPR provider after 2 minutes
102 according to current guidelines. The pairs allocated to the Dual Operator group were
103 instructed to perform 4 minutes of Dual Operator CPR (see Figure 1); where one BLS
104 provider performs chest compressions only and the other BLS provider performs rescue
105 breaths only, swapping roles every 2 minutes.

106

107

108 Before the session, the researcher instructed the candidates on how to perform Dual
109 Operator or Single Operator CPR. Candidates were instructed to use current European
110 Resuscitation Council BLS Guidelines to perform chest compressions and rescue breaths,
111 i.e. a ratio of 30 chest compressions (at a rate of 100min^{-1}) to 2 rescue breaths (each
112 delivered over 1 second).**Error! Bookmark not defined.** The instructions were
113 consistent throughout the study, always asking the candidates to start with chest
114 compressions. The only other input the researcher had was to begin and end the session.

115 The researcher did not indicate at any point during the 4 minutes how far through the
116 session the candidates were.

117

118 After completing the initial CPR assessment, candidates returned 1 week later to perform
119 the crossover part of the study. Upon their return candidates who were initially in the
120 'Dual Operator' group were asked to perform 4 minutes of Single Operator CPR and
121 candidates initially in the 'Single Operator' group were asked to perform 4 minutes of
122 Dual Operator CPR.

123

124 **Data collection**

125 CPR performance was assessed objectively using a Laerdal Skillmeter Manikin Resusci®
126 Anne with PC Skillmeter VAM software which allows data variables about the quality of
127 CPR to be downloaded to a laptop computer. The variables collected were: no-flow
128 duration, number of compressions, number of correct compressions, compression rate,
129 compression depth, duty cycle, number of rescue breaths, rescue breath volume, rescue
130 breath flow rate and minute volume. The PC Skillmeter VAM software was programmed
131 to take into account the ERC BLS Guidelines 2005 in order to count a 'correct' chest
132 compression (depth 38-51 mm, correct hand position, complete release). During the
133 assessment, both the candidate and researcher were blinded to the VAM software output.

134

135 **Statistical methods**

136 Using data from our previous study⁹ we calculated that we would require 22 patients to
137 detect a 10% difference in no-flow times with 90% power at a significance level of
138 0.05. We aimed to recruit 25 participants to allow for any loss to follow-up.

139 Data were analysed by SPSS 13 (SPSS Inc). Data were checked for normality using the
140 Shapiro-Wilk Tests. Data were normally distributed and therefore analysed using paired
141 t-tests. For all statistical testing a P-value <0.05 was considered statistically significant.

142

143

144 **Results**

145 58 candidates were assessed for eligibility and 8 were excluded before randomisation
146 (n=8 due to personal commitments making them unable to attend CPR assessments). At
147 initial CPR assessment there were 50 candidates (Dual Operator group n=26, Single
148 Operator group n=24). At the crossover assessment 1 week after the initial assessment
149 there were no candidates lost to follow up (Dual Operator group n=26, Single Operator
150 group n=24). Figure 2 shows the flow of candidates through the study.

151

152 **Participant characteristics**

153 All participants were 2nd year medical students who had completed the ERC BLS/AED
154 Instructor course in the preceding 3 months. All participants were involved in teaching
155 on our peer led instructor programme at the time of the study.^{10, 11} The initial Dual
156 Operator group consisted of 18 women (69%) and 8 men (31%). The initial Single
157 Operator group consisted of 15 women (63%) and 9 men (38%). The mean age (years) in
158 the groups was very similar (20.5 in the Single Operator group versus 20.6 in the Dual
159 Operator group).

160

161

162 **CPR Performance**

163 Data from the assessment of CPR performance are presented in Table 1. Dual Operator
164 CPR achieved lower no-flow duration than Single Operator CPR. This improvement,
165 although small, was statistically significant (28.5% (S.D. = 3.7) versus 31.6% (S.D. =
166 3.6), P<0.001). In absolute values, these percentages equate to 68 and 76 seconds without

167 compressions for Dual and Single Operator CPR respectively, a difference of 8 seconds
168 over the four-minute CPR session. To explore the impact of cohort averaging reducing
169 the magnitude of difference between techniques the data on no flow proportions were
170 dichotomised into two group – those where dual operator CPR increased no flow time
171 and those where it reduced it. No flow duration for dual operator CPR decreased in 20
172 out of 25 of participants (32.3%(3.4) versus 27.8% (3.5), P=0.0001) and increased in five
173 participants (28.9%(3.1) versus 31.2%(3.2), P=0.005).

174

175 There was no other difference in the performance of chest compressions. Specifically
176 compression depth, compression rate, percentage of correct chest compressions and duty
177 cycle were all very similar.

178

179 There were some small but statistically significant differences for the performance of
180 rescue breaths. Dual Operator CPR achieved more rescue breaths per minute (4.9 (S.D. =
181 0.5) versus 4.5(S.D. = 0.5), P=0.009) and a higher minute volume (3730ml (S.D. =
182 490)versus 3387ml (S.D. = 414), P=0.006). There was no difference in rescue breath
183 flow rate or rescue breath volume.

184

185 **Discussion**

186 The principal finding of this study was that compared to Single Operator CPR, Dual
187 Operator CPR achieved a statistically significant reduction in no-flow duration when
188 compared with Single Operator CPR (28.53% versus 31.62%, P=<0.001). However, the
189 magnitude of the improvement (3%) was small and would be unlikely to have any major

190 effect on patient outcome were they to be reproduced in an actual resuscitation attempt.
191 There were no other clinically significant differences in CPR (rescue breath or chest
192 compression) performance between techniques.
193
194 The quality of CPR is an important determinant of survival from cardiac arrest.¹²⁻¹⁵
195 Studies have stressed the importance of minimising interruptions in chest compressions in
196 order to maintain coronary perfusion pressure and improve the chance of successful
197 defibrillation.¹⁶ Observational studies in humans in cardiac arrest have reported
198 prolonged interruptions in chest compression in clinical practice. Valenzuela et al.
199 reported that chest compressions were not performed 57% of the time during pre-hospital
200 resuscitation attempts.¹⁷ Wik et al. showed in series of 176 out of hospital CPR attempts
201 no chest compressions were performed 38% of the time even allowing for the time
202 necessary for electrocardiographic analysis, pulse checks and defibrillation.¹⁸ When two
203 or more trained rescuers are present (e.g. a lifeguard team; community first responders),
204 one strategy for minimizing interruptions in chest compressions is to undertake Dual
205 Operator CPR, which theoretically would reduce interruptions in chest compressions as a
206 result of the rescuer switching between ventilations and chest compressions.
207
208 The ERC Basic Life Support (BLS) guidelines from 2000 recommended Dual Operator
209 CPR when 2 trained rescuers were present. However, this recommendation was
210 withdrawn in the 2005 revision of the Guidelines. The decision to change the guidelines
211 was taken on a pragmatic basis rather than as a result of new evidence specifically in this
212 scenario. Underpinning the change in BLS guidelines between 2000 and 2005 was the

213 idea of simplifying the algorithm to simplify teaching, in order to improve retention of
214 skills¹⁹. However, another objective of Guidelines 2005 was to improve the quality of
215 CPR and reduce interruptions in chest compressions. At the time of these changes, only
216 one study had indirectly looked at the effect of Dual as opposed to Single Operator CPR.
217 In a manikin study investigating the performance of CPR in confined spaces, Handley
218 and Handley demonstrated a reduction in no-flow times (10 versus 6 seconds per CPR
219 cycle) when Dual Operator CPR was performed. Therefore, it was possible that dropping
220 Dual Operator CPR from the guidelines could have inadvertently led to a reduction in the
221 quality of CPR.

222

223 The present study differs from the Handley and Handley study in that it was conducted in
224 accordance with Guidelines 2005 which recommends a compression to ventilation ratio
225 of 30:2 as opposed to 15:2. The change in compression to ventilation ratio has been
226 associated with a significant reduction in no-flow duration.²⁰ We hypothesise that the
227 difference in compression to ventilation ratio between the two studies explains why this
228 study found that the improvement in no-flow duration was marginal with the two operator
229 approach. The present study therefore supports the decision to drop Dual Operator CPR
230 from the BLS curriculum as Dual Operator CPR increases the complexity of the
231 guidelines without any meaningful benefits in terms of quality of CPR.

232

233 This study has several limitations. Firstly, the study set out to evaluate the impact of
234 Dual Operator CPR used by a team of trained first responders. These findings and the
235 recommendation that Dual Operator CPR should not be taught to first responders / lay

236 persons applies only to this group of trainees. Dual Operator CPR should continue to
237 form part of the training pathways for pre or in-hospital advanced life support
238 resuscitation teams as these teams are tasked with multiple interventions (e.g.
239 defibrillation, advanced airway management) in contrast to the first responder group in
240 this study. Secondly, we used an “expert group” of BLS CPR providers (trained
241 instructors). The quality of CPR in the study although better than has been seen in other
242 clinical studies, still only yielded moderate compression performance (% correct
243 compressions circa 40%) so these results may not necessarily extrapolate to clinical
244 practice. Thirdly, the study period was relatively short at only 4 minutes and thus did not
245 fully examine the impact of fatigue on performance. In many areas ambulance response
246 times are in the region of 8 minutes. Whether any differences would have been seen over
247 a longer duration of CPR was not investigated. Finally, it was not possible to blind
248 candidates to the technique they were performing. Whilst we have no reason to suspect
249 that this influenced the results, we cannot exclude this as a possibility.

250

251

252 **Conclusion**

253 Dual Operator CPR provides marginal improvement in minimising interruptions in chest
254 compressions when compared to Single Operator CPR performed by BLS resuscitation
255 teams. There are no other differences in the performance of CPR between Dual Operator
256 and Single Operator CPR when 2 rescuers are present. There seems little advantage in
257 adding teaching of Dual Operator CPR to trained first responder/BLS CPR programs in
258 view of the added complexities.

259

260 **Conflicts of interest**

261 None declared.

262

263 **Acknowledgements**

264 Thanks to Rebecca Taylor, Andy Owen, Laura Kocierz and the RMD teachers for
265 participating. Finally to Professor Julian Bion and Dr Jon Hulme for their continued
266 support.

267

268 **Funding:**

269 The manikins in this study were paid for by a grant from the Resuscitation Council (UK).

270 GDP is supported by a Department of Health NIHR Clinician Scientist Award.

271

272

273 **References**

274

- 275 1. Hightower D, Thomas SH, Stone CK, Dunn K, March JA. Decay in quality of
276 closed-chest compressions over time. *AnnEmergMed* 1995;26:300-3.
- 277 2. Perkins GD, Augre C, Rogers H, Allan M, Thickett DR. CPREzy: an evaluation
278 during simulated cardiac arrest on a hospital bed. *Resuscitation* 2005;64:103-8.
- 279 3. Ochoa FJ, Ramalle-Gomara E, Lisa V, Saralegui I. The effect of rescuer fatigue
280 on the quality of chest compressions. *Resuscitation* 1998;37:149-52.
- 281 4. Handley AJ, Handley JA. Performing chest compressions in a confined space.
282 *Resuscitation* 2004;61:55-61.
- 283 5. Handley AJ, Monsieurs KG, Bossaert LL. European Resuscitation Council
284 Guidelines 2000 for Adult Basic Life Support. A statement from the Basic Life Support
285 and Automated External Defibrillation Working Group(1) and approved by the Executive
286 Committee of the European Resuscitation Council. *Resuscitation* 2001;48:199-205.
- 287 6. Handley AJ, Koster R, Monsieurs K, Perkins GD, Davies S, Bossaert L. European
288 Resuscitation Council guidelines for resuscitation 2005. Section 2. Adult basic life
289 support and use of automated external defibrillators. *Resuscitation* 2005;67 Suppl 1:S7-
290 23.
- 291 7. Hackman BB, Kellermann AL, Everitt P, Carpenter L. Three-rescuer CPR: the
292 method of choice for firefighter CPR? *Ann Emerg Med* 1995;26:25-30.
- 293 8. Dunkley CJ, Thomas AN, Taylor RJ, Perkins RJ. A comparison of standard and a
294 modified method of two resuscitator adult cardiopulmonary resuscitation: description of a
295 new system for research into advanced life support skills. *Resuscitation* 1998;38:7-12.
- 296 9. Spooner BB, Fallaha JF, Kocierz L, Smith CM, Smith SC, Perkins GD. An
297 evaluation of objective feedback in basic life support (BLS) training. *Resuscitation*
298 2007;73:417-24.
- 299 10. Perkins GD, Hulme J, Bion JF. Peer-led resuscitation training for healthcare
300 students: a randomised controlled study. *Intensive Care Med* 2002;28:698-700.
- 301 11. Bucknall V, Sobic EM, Wood HL, Howlett SC, Taylor R, Perkins GD. Peer
302 assessment of resuscitation skills. *Resuscitation* 2008;77:211-5.
- 303 12. Kramer-Johansen J, Edelson DP, Losert H, Kohler K, Abella BS. Uniform
304 reporting of measured quality of cardiopulmonary resuscitation (CPR). *Resuscitation*
305 2007;74:406-17.
- 306 13. Edelson DP, Litzinger B, Arora V, et al. Resuscitation with actual performance
307 integrated debriefing (RAPID) improves CPR quality and initial patient survival.
308 *Resuscitation Science Symposium* 2008;A61:29.
- 309 14. Edelson DP, Abella BS, Kramer-Johansen J, et al. Effects of compression depth
310 and pre-shock pauses predict defibrillation failure during cardiac arrest. *Resuscitation*
311 2006;71:137-45.
- 312 15. Leary M, Abella BS. The challenge of CPR quality: improvement in the real
313 world. *Resuscitation* 2008;77:1-3.
- 314 16. Eftestol T, Sunde K, Steen PA. Effects of interrupting precordial compressions on
315 the calculated probability of defibrillation success during out-of-hospital cardiac arrest.
316 *Circulation* 2002;105:2270-3.

- 317 17. Valenzuela TD, Kern KB, Clark LL, et al. Interruptions of chest compressions
318 during emergency medical systems resuscitation. *Circulation* 2005;112:1259-65.
- 319 18. Wik L, Kramer-Johansen J, Myklebust H, et al. Quality of cardiopulmonary
320 resuscitation during out-of-hospital cardiac arrest. *JAMA: The Journal of the American*
321 *Medical Association* 2005;293:299-304.
- 322 19. Nolan J. European Resuscitation Council Guidelines for Resuscitation 2005:
323 Section 1. Introduction. *Resuscitation* 2005;67:S3-S6.
- 324 20. Perkins GD, Boyle W, Bridgestock H, et al. Quality of CPR during advanced
325 resuscitation training. *Resuscitation* 2008;77:69-74.
- 326
- 327