

Technical University of Denmark



Does an onion-enriched diet beneficially affect the microbial composition in healthy human subjects?

Bergström, Anders; Borg, Birgitte; Marin, Eduvigis Roldán; Damsgaard, Camilla Trab; Dragsted, Lars Ove; Licht, Tine Rask

Publication date:
2009

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Bergström, A., Borg, B., Marin, E. R., Damsgaard, C. T., Dragsted, L. O., & Licht, T. R. (2009). Does an onion-enriched diet beneficially affect the microbial composition in healthy human subjects?. Poster session presented at 7th Symposium of Food Microbiology, Helsingør, Denmark.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Does an onion-enriched diet beneficially affect the microbiotal composition in healthy human subjects?

Anders Bergström¹, Birgitte Borg², Eduvigis Roldán Marin³,
Camilla Trab Damsgaard², Lars O Dragsted², Tine Rask Licht¹


¹DTU Food, ²Dept. Human Nutrition, University of Copenhagen;
³Instituto del Frio, University of Madrid;

Corresponding author: adbe@food.dtu.dk

1) Background:

Onions are rich in fructo-oligosaccharides (FOS), which are well acknowledged prebiotic substances. FOS consumption have previously been associated with an increased level of fermenting bacterial genera e.g. Lactobacillus and Bifidobacterium. Generally, these groups of bacteria are considered to have beneficial effects on the intestinal environment. The aim of the present study was to use quantitative PCR and DGGE to analyze the effects of onion consumption on the gut microbiotal profile.

3) Study timeline for one intervention period




5) DGGE results:

DGGE was run on PCR amplified universal bacterial 16S rRNA genes of the isolated stool DNA.

Gel representing DGGE profiles of 3 subjects on Yellow diet is shown here.

Note the band present in the Before (sample days 1+2), but not the After (sample days 3+4) samples for individuals 1 and 2. Gels for subjects 4-6 needs to be run again due to insufficient PCR reaction (not shown)

Gels for the Red diet showed no obvious differences



2) Study design:

- 10 human subjects (5 male + 5 females)
- 24-67 years old
- BMI: 25-35 kg/m²
- No other restriction diets allowed
- Randomized double-blind crossover design
- Two 14 day dietary intervention periods

- A) Onion enriched (Yellow/Red)
- B) Onion not enriched (Yellow/Red)

- 8 days restriction diet before each intervention period excluding onions and many other possibly confounding vegetables etc

4) qPCR results:

Total DNA was isolated from 4 stool samples (Sample Days 1+2, Before diet, and Sample Days 3+4, After diet) for each individual, for each of the two diets, using the Qiagen Stool Kit and subsequently quantitative real-time PCR was performed with primers representing the Genera:

Lactobacillus, Bifidobacterium, Bacteroidetes, and Clostridium

Note that no differences between Sample Days are present for any of the tested primers.

However, there are large individual differences in both the Yellow and Red diet groups.

t-test of differential expression in each individual showed no systematic effects (not shown)

Strain/Test	Yellow		Red	
	2-way ANOVA Days (1) x Individuals (10)	1-way ANOVA Time Period (Before/After)	2-way ANOVA Days (4) x Individuals (5)	1-way ANOVA Time Period (Before/After)
Bifido	$F_{3,15}=0.197, p=0.99$ (no effect of Day) $F_{3,15}=3.73, p=0.022^*$ (clear Individual effect)	$F_{1,22}=0.10, p=0.75$ (no treatment effect)	$F_{3,15}=1.11, p=0.38$ (no effect of Day) $F_{3,15}=5.15, p=0.0027^*$ (clear Individual effect)	$F_{1,22}=0.08, p=0.78$ (no treatment effect)
Lactobacillus	$F_{3,15}=0.53, p=0.67$ (no effect of Day) $F_{3,15}=1.52, p=0.21$ (no Individual effect)	$F_{1,22}=1.06, p=0.31$ (no treatment effect)	$F_{3,15}=1.36, p=0.24$ (no effect of Day) $F_{3,15}=6.30, p=0.0016^*$ (clear Individual effect)	$F_{1,22}=1.015, p=0.32$ (no treatment effect)
Clostridia	$F_{3,15}=1.49, p=0.26$ (no effect of Day) $F_{3,15}=26.25, p<<0.0001^*$ (clear Individual effect)	$F_{1,22}=0.42, p=0.52$ (no treatment effect)	$F_{3,15}=1.62, p=0.23$ (no effect of Day) $F_{3,15}=22.46, p<<0.0001^*$ (clear Individual effect)	$F_{1,22}=0.08, p=0.93$ (no treatment effect)
Bacteroides	Failed – to many missing obs. $F_{1,21}=0.113, p=0.91$ (no treatment effect)	Failed – to many missing obs.	Failed – to many missing obs. $F_{1,21}=0.113, p=0.91$ (no treatment effect)	Failed – to many missing obs.

6) Conclusions:

• qPCR of selected primers showed no effect of treatment (Red/Yellow)

• Significant differences in individual Bifido, Lactobacillus, Clostridia and Bacteroides found

• DGGE of universal primers showed no or few obvious changes, but it would be a good idea to repeat the PCR, since gels representing half the test population were less efficient

• It is unfortunate that only 6 of 10 subjects finished the protocol. Makes it difficult to draw conclusions (or expect) findings

• The gel bands differing between Before and After in the DGGE gel merits some attention