Technical University of Denmark



In silico evolution leads to green chemical factory

Vanillin green cell factory

Brochado, Ana Rita; Matos, Claudia; Mortensen, Uffe Hasbro; Patil, Kiran Raosaheb

Publication date: 2009

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA):

Brochado, A. R., Matos, C., Mortensen, U. H., & Patil, K. R. (2009). In silico evolution leads to green chemical factory: Vanillin green cell factory. Poster session presented at SYSTEMS BIOLOGY AS A DRIVER FOR INDUSTRIAL BIOTECHNOLOGY, Istanbul, Turkey, .

DTU Library Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



In silico evolution leads to green chemical factory

A R Brochado¹, C Matos¹, J Hansen², B L Møller³, U H Mortensen¹ and K R Patil¹

¹Center for Microbial Biotechnology, Technical University of Denmark, ²Evolva A/S, Denmark, ³ KU-LIFE, Copenhagen University, Denmark

Motivation

The beneficial metabolic properties of naturally occurring microorganisms are a result of an evolutionary process that increases their fitness in the natural habitat. Phenotype conferring to the fitness is often inclined towards growth and less so towards the trait of interest, e.g. high productivity of a desired compound. Random mutagenesis followed by selection pressure is a widely used way to obtain high-producing strains by exploiting the genetic plasticity. The space of possible genetic alterations, however, is too large to explore in either random or in a brute-force manner. We propose an alternative solution to this problem, by *mimicking* the Darwinian evolution process *in silico*. The motivation for this attempt lies in the availability of complete genome sequences and modeling concepts capable of simulating cellular phenotype. We explore the possibility of using such *in silico* evolution for obtaining metabolic engineering targets towards high productivity. The applicability of this approach is demonstrated with a case study of heterologous production of

vanillin, one of the most popular flavor compound obtained from an Orchidea, in the baker's yeast – Saccharomyces cerevisiae.



Figure 1 - Biosynthetic pathway for *de novo* biosynthesis of vanillin glucoside in *S. cerevisiae*. 3DSD: 3dehydroshikimate dehydratase from *Podospora pausiceta*; ACAR: aromatic carboxylic acid reductase from Nocardia sp; PPTase: phophopantetheinyl transferase from *E. coli;* OMT: O-methyltransferase from *Homo Sapiens*; UGT: UDP-glycosyltransferase from *Arabidopsis Thaliana* (Hansen, et al., 2009).



Figure 2: Schematic representation of the OptGene algorithm. The algorithm starts with a initial population producing low amount of vanillin that evolves through several generations where the best producers are progressively selected leading to the prediction of a high producing genotype.



cultivation. **B**. Substrate specific yield of vanillin glucoside, PAC and PAL in batch cultivation. **C**. Substrate specific yield for biomass profiles in batch production in continuous cultivation. D: Substrate specific yield of vanillin glucoside, PAC and PAL in glucoside, PAC and PAL in continuous cultivation.

Conclusion and Future perspectives

A significant improvement of glucovanillin production was observed for the model driven obtained mutants. Genome-wide *omics* detailed investigation of these strains, as transcriptome and fluxome analysis will be performed and incoming results will provide an excellent platform for further strain design applying model based strategies. The resulting data may serve as a starting point for incorporating regulatory information into the yeast metabolic model. In summary, *in silico* design by using an evolutionary approach was found to be a good platform for rational improvement of microbial cell factories – a promising step towards green chemical industry.

Hansen EH [et al.] De novo biosynthesis of vanillin in fission and baker's yeast [Journal]. - [s.l.] : Applied Environmental Microbiology, 2009. Patil KR, Rocha I, Forster J and Nielsen, J. 2005. "Evolutionary programming as aplatform for *in silico* metabolic engineering". BMC Bioinformatics. Vol. 6, p. 308. Forster J, Famili L, Fu P, Palsson BO and Nielsen J. 2003. "Genome-scale reconstruction of the *Saccharomyces cerevisiae* metabolic network". Genome Research. Vol.13, p. 244-253.