

Technical University of Denmark



## Transcription factors as targets for improving *Aspergillus niger* as cell factory

Poulsen, Lars; Bruno, K.S.; Thykær, Jette; Baker, S.E.; Eliasson Lantz, Anna

*Publication date:*  
2009

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Poulsen, L., Bruno, K. S., Thykær, J., Baker, S. E., & Eliasson Lantz, A. (2009). Transcription factors as targets for improving *Aspergillus niger* as cell factory. Poster session presented at SYSTEMS BIOLOGY AS A DRIVER FOR INDUSTRIAL BIOTECHNOLOGY, Istanbul, Turkey, .

**DTU Library**  
Technical Information Center of Denmark

---

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Transcription factors as targets for improving *Aspergillus niger* as cell factory

L. Poulsen<sup>1\*</sup>, K. S. Bruno<sup>2</sup>, J. Thykaer<sup>1</sup>, S. E. Baker<sup>2</sup>, A. E. Lantz<sup>1</sup>

(1) Center for Microbial Biotechnology, Department of Systems Biology, Technical University of Denmark, Lyngby, Denmark

(2) Fungal Biotechnology Team, Pacific Northwest National Laboratory, Richland, Washington 99352, USA  
\*lap@bio.dtu.dk


## Introduction

Altering fluxes for overcoming metabolic bottlenecks have traditionally been approached by genetic engineering of a single or few metabolic genes. This strategy struggles to overcome the subjacent regulation thus the outcome has frequently shown to be of limited success. Transcription factors have the potential of controlling several fluxes in an organism, hence manipulating expression of these proteins can provide an alternative tool for overcoming metabolic bottlenecks. This approach has previously been demonstrated in yeast with great success for production of ethanol (Schuurmans et al., 2008).

In the present study the effect of modulation of TFs in *A. niger*, which is an industrially important micro-organism. The strategy described in this work focuses on regulation connected to pH. It was chosen as an important process parameter, due to its significant influences on both organic acid and enzyme production.


## Locating targets

In order to locate targets for knockout, data from a previous transcription analysis (Andersen et al., 2009) was utilized. Several putative TF with pH responding behavior was identified and formed the basis for gene-knockout.



## Strain construction


The *A. niger* ATCC 1015 was used as Wild-type strain and was the basis for all strain constructions.



## Physiological characterisation

### Screening


Three novel TF mutants were screened for protease activity and acid production and compared against Wild-type and a previously described protease deficient mutant  $\Delta$ prtT (P. J Punt et al., 2008). The experiments were performed for 4 days as 6 mL stationary cultures in minimal- and complex medium.




### Batch fermentations

To further characterize MUT 1, which expressed an oxalic acid overproducing phenotype (OOP), this mutant was cultivated in 2L scale bioreactors.



Parameters	Germination	Fermentation
Aeration	0.10 vvm	1 vvm
Stir speed	200 RPM	1000 RPM
pH	3.0	6.0



### Protease activity



### Acid production



## References

- Andersen MR, Lehmann L, Nielsen J. (2009) Systemic analysis of the response of *Aspergillus niger* to ambient pH. *Genome Biol.* 10 (45): R47.  
Schuurmans JM, Rossell SL, van Tuijl A, Bakker BM, Hellingwerf KJ, Teixeira de Mattos MJ. (2008) Effect of hxx2 deletion and HAP4 overexpression on fermentative capacity in *Saccharomyces cerevisiae*. *FEMS Yeast Res.* 8(2):195-203.  
Punt PJ, Schuren FH, Lehmebeck J, Christensen T, Hjort C, van den Hondel CA. (2008) Characterization of the *Aspergillus niger* prtT, a unique regulator of extracellular protease encoding genes. *Fungal Genet Biol.* 2008 Dec;45(12):1591-9

## Conclusion

- Several pH responding TF deficient mutants were constructed and screened for protease activity and acid production
- One mutant had an oxalic acid overproducing phenotype (OOP) and one mutant had indications of being protease deficient.
- The OOP mutant was further characterized in 2L scale bioreactors, and a 90% ( $\pm 25\%$ ) increase of oxalic acid on glucose was seen.