

Technical University of Denmark



Demonstration of quadrature squeezed surface-plasmons in a gold waveguide

Huck, Alexander; Smolka, Stephan; Krivitsky, Leonid; Lodahl, Peter; Sørensen, A. S.; Boltasseva, Alexandra; Andersen, Ulrik Lund

Published in:

Conference abstract series, CLEO/Europe - EQEC

Link to article, DOI:

[10.1109/CLEOE-EQEC.2009.5191638](https://doi.org/10.1109/CLEOE-EQEC.2009.5191638)

Publication date:

2009

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Huck, A., Smolka, S., Krivitsky, L., Lodahl, P., Sørensen, A. S., Boltasseva, A., & Andersen, U. L. (2009). Demonstration of quadrature squeezed surface-plasmons in a gold waveguide. In Conference abstract series, CLEO/Europe - EQEC IEEE. DOI: 10.1109/CLEOE-EQEC.2009.5191638

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Demonstration of Quadrature Squeezed Surface-Plasmons in a Gold Waveguide

Alexander Huck,¹ Stephan Smolka,² Leonid Krivitsky,^{1,3} Peter Lodahl,² Anders S. Sørensen,⁴ Alexandra Boltasseva,² and Ulrik L. Andersen¹

¹Department of Physics, Technical University of Denmark, Building 309, 2800 Kgs. Lyngby, Denmark

²DTU Fotonik, Department of Photonics Engineering, Technical University of Denmark, Ørsted Plads 343, 2800 Kgs. Lyngby, Denmark

³Data Storage Institute, Agency for Science, Technology and Research (A-Star), 117608 Singapore

⁴QUANTOP, Danish Quantum Optics Center and Niels Bohr Institute, 2100 Copenhagen, Denmark

The first experiment demonstrating the quantum optical properties of SPPs was the preservation of entanglement under plasmon assisted transmission through sub-wavelength holes in a conductor [1,2].

In this contribution we present an experiment demonstrating the generation of non-classical SPPs by exciting them with a squeezed optical light field generated using a bow-tie shaped optical parametric oscillator operating below threshold. Free space optics and end-fire coupling are used for the excitation of long-range SPPs (LR-SPPs) [3] on gold stripes embedded in lossless transparent polymer BCB. The gold stripes have a length, width, and height of 2mm, 1 μ m, and 14nm, respectively. At maximum, we measure a total transmission of 33.2 \pm 0.5% through the sample. The squeezed vacuum input mode and the LR-SPP output mode are characterized using an optical homodyne detection system. We employ the maximum likelihood method to reconstruct the density matrices of the squeezed vacuum input state and the LR-SPP output state. The noise power of the LR-SPP output state as a function of relative phase is presented in Fig. 1(a). We simulate the impact of the LR-SPP mode by applying a beam-splitter operation on the squeezed vacuum input mode. The overlap between the calculated output mode and the LR-SPP mode is evaluated by calculating the Fidelity between the two states, which peaks for a transmission of η =33%, as shown in Fig. 1(b). This result is in very good agreement with the transmission measured classically and proves the validity of the beam-splitter model in the limit of low photon number excitation.




Fig. 1 (a) Noise variance of the LR-SPP output state calculated from the time resolved data (black solid line) and from the reconstructed LR-SPP states density matrix (red dashed line). (b) Calculated overlap between the squeezed vacuum input state and the LR-SPP output state for various transmissions η through a beam splitter.

Furthermore, we also demonstrate the excitation of LR-SPP modes with a priori two photon Fock states generated by spontaneous parametric down-conversion.

In conclusion, we demonstrated the excitation of non-classical LR-SPPs and verified, that the transmission of LR-SPP modes in a metallic waveguide can be described by a beam-splitter operation.

References

- [1] E. Altewischer, M.P. van Exter, and J.P. Woerdeman, "Photon-assisted transmission of entangled photons", *Nature* **418**, 304-306 (2002).
- [2] E. Moreno, F. J. Garcia-Vidal, D. Erni, J. Ignacio Cirac, and L. Martín-Moreno, "Theory of Plasmon-Assisted Transmission of Entangled Photons," *Phys. Rev. Lett* **92**, 236801 (2004).
- [3] J.J. Burke, G.I. Stegeman, and T. Tamir, "Surface-polariton-like waves guided by thin, lossy metal films," *Phys. Rev. B* **33**, 5186 (1986).