

Technical University of Denmark



Continuous Variable Entanglement and Squeezing of Orbital Angular Momentum States

Lassen, Mikael Østergaard; Leuchs, Gerd; Andersen, Ulrik Lund

Published in:
Physical Review Letters

Link to article, DOI:
[10.1103/PhysRevLett.102.163602](https://doi.org/10.1103/PhysRevLett.102.163602)

Publication date:
2009

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Lassen, M. Ø., Leuchs, G., & Andersen, U. L. (2009). Continuous Variable Entanglement and Squeezing of Orbital Angular Momentum States. *Physical Review Letters*, 102(16), 163602. DOI: 10.1103/PhysRevLett.102.163602

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Continuous Variable Entanglement and Squeezing of Orbital Angular Momentum States

M. Lassen,^{1,2} G. Leuchs,^{2,3} and U. L. Andersen¹

¹*Department of Physics, Technical University of Denmark, Fysikvej, 2800 Kongens Lyngby, Denmark*

²*Max Planck Institute for the Science of Light, Günther Scharowskystrasse 1, 91058 Erlangen, Germany*

³*University Erlangen-Nürnberg, Staudtstrasse 7/B2, 91058 Erlangen, Germany*

(Received 19 January 2009; published 23 April 2009)

We report the first experimental characterization of the first-order continuous variable orbital angular momentum states. Using a spatially nondegenerate optical parametric oscillator (OPO) we produce quadrature entanglement between the two first-order Laguerre-Gauss modes. The family of orbital angular momentum modes is mapped on an orbital Poincaré sphere, where the mode's position on the sphere is spanned by the three orbital parameters. Using a nondegenerate OPO we produce squeezing of these parameters, and as an illustration, we reconstruct the “cigar-shaped” uncertainty volume on the orbital Poincaré sphere.

DOI: 10.1103/PhysRevLett.102.163602

PACS numbers: 42.50.Dv, 03.67.-a

Propagating light beams have a spin angular momentum and an orbital angular momentum (OAM), which are associated with the polarization and the phase distribution of the light state, respectively [1]. The *quantum* spin angular momentum (or polarization) has been thoroughly investigated both for discrete and for continuous variables (CV) [2–4]. Recently the *quantum* OAM of light has attracted a lot of attention, in the single photon regime, due to its unique capabilities in tailoring the dimensionality of the Hilbert space [5–7]. In contrast, there has been very little work devoted to the OAM of intense modes described by CV; the only account in this regime is the very recent work on the creation of CV entanglement between two spatially separated OAM modes generated in atomic vapor exploiting the nonlinear process of four-wave mixing [8,9]. The interest in CV OAM states stems from the further increase of the dimensionality of the Hilbert space. OAM of single photons has already found uses in various quantum information protocols, and we envisage that the CV OAM states hold similar potentials in CV quantum information and also in quantum imaging and quantum metrology. The most promising application of CV OAM states is their connectivity with atoms, thus allowing for storage of CV quantum information [10,11].

The simplest spatial mode that can carry OAM is the first-order Laguerre-Gaussian (LG) mode which produces either a left-handed or right-handed corkscrewlike phase front and a ring-structured intensity profile. They are denoted $LG_{p=0}^{l=\pm 1}$, where l and p are the azimuthal and radial mode indices. Similarly to quantum spin angular momentum (polarization) states, the family of first-order OAM states can be represented on a Poincaré-sphere analog which is also known as the first-order orbital Poincaré sphere [12,13]. The sphere is spanned by three parameters (associated with three different OAM states of light) that have properties identical to those of the three polarization Stokes parameters and which we name the orbital param-

eters. In this Letter we demonstrate the generation of squeezing in the orbital parameter of OAM modes using a spatially nondegenerate optical parametric oscillator and a specially tailored local oscillator. In addition, we measure quadrature entanglement between the two first-order LG modes thereby demonstrating a new type of entanglement from nondegenerate optical parametric oscillators (OPOs). The demonstration of similar spatially nondegenerate OPOs has recently been independently reported in Ref. [14].

The quantum polarization degree of freedom (or spin angular momentum) has been extensively explored and characterized in the Schwinger representation in terms of the quantum Stokes operators both in the two-dimensional and the infinite-dimensional Hilbert space, where the Stokes parameter eigenvalues are either discrete or continuous [2–4]. The Stokes operators are decomposed into field operators for orthogonal polarization modes and completely represent the quantum dynamics of the polarization of light. Likewise, we define the Stokes operator analogs—the orbital operators—for the first-order OAM modes as

$$\begin{aligned}\hat{O}_1 &= \hat{A}_{HG_{10}}^\dagger \hat{A}_{HG_{10}} - \hat{A}_{HG_{01}}^\dagger \hat{A}_{HG_{01}} \\ \hat{O}_2 &= \hat{A}_{HG_{10(45^\circ)}}^\dagger \hat{A}_{HG_{10(45^\circ)}} - \hat{A}_{HG_{10(135^\circ)}}^\dagger \hat{A}_{HG_{10(135^\circ)}} \\ \hat{O}_3 &= \hat{A}_{LG_0^1}^\dagger \hat{A}_{LG_0^1} - \hat{A}_{LG_0^{-1}}^\dagger \hat{A}_{LG_0^{-1}},\end{aligned}\quad (1)$$

where \hat{A}^\dagger and \hat{A} are the creation and annihilation operators for the various spatial first-order modes given by the indices, and illustrated on the sphere in Fig. 1. As clearly seen from these definitions, \hat{O}_1 , \hat{O}_2 , and \hat{O}_3 represent the difference in the photon number between the two modes HG_{10} and HG_{01} , $HG_{10(45^\circ)}$ and $HG_{10(135^\circ)}$, and LG_0^{+1} and LG_0^{-1} , respectively. Similarly to the polarization Stokes operators, these orbital operators completely represent the dynamics of the first-order spatial states, and thus these operators follow the same algebra as the Stokes operators,




FIG. 1 (color online). Orbital Poincaré sphere of the first-order OAM modes. Points on the sphere are associated with a superposition of OAM modes [13]: $\psi_{\phi, \varphi} = \cos(\phi/2)LG_0^1 + \exp(i\varphi)\sin(\phi/2)LG_0^{-1}$, with azimuthal and axial angles ϕ and φ .

namely, the SU(2) algebra [15]. Likewise, the commutation relations are $[\hat{O}_k, \hat{O}_l] = i\hat{O}_m$, where $k, l, m \in \{1, 2, 3\}$ of cyclic permutation.

Spatially multimode nonclassical CV states have previously been generated in pulsed optical parametric amplification [16], in atomic vapor [8], in a confocal optical parametric oscillator [17], and in linear interference between different single modes [18–20]. Here we use another approach that previously has been used for the generation of single-spatial-mode squeezed states, namely, a mode-stable (and nonconfocal) optical parametric oscillator [21]. By employing a type I phase-matched nonlinear crystal in a mode-stable optical cavity, the polarization, the frequency, and the spatial degree of freedom are usually degenerate, which leads to single mode quadrature squeezing as demonstrated by various groups. In all these experiments, the cavity supported only the Gaussian zeroth order LG mode. However, by changing the cavity resonance frequency it is possible to generate the two first-order modes simultaneously due to their frequency degeneracy (stemming from their identical Gouy phase shifts). This means that the down-converted signal and idler photons are produced in two distinct orthogonal spatial modes (the OAM modes, LG_0^{+1} and LG_0^{-1}), and thus create quadrature entanglement between these two modes similarly to the production of entanglement between polarization modes [22] or between frequency modes [23] in polarization or frequency non-degenerate OPOs. The nondegeneracy of the first-order LG modes therefore adds a new member to the family of nondegenerate OPOs capable of producing entanglement. This was also discussed in Ref. [24] for an OPO above the oscillation threshold. In the following, we experimentally demonstrate the generation of quadrature entangled LG modes and show that this can be used to produce squeezing in the first-order orbital parameters. Note that a thorough analysis of the transfer of OAM from the pump to the down-converted fields was carried out in Ref. [25].

Our experimental setup is depicted in Fig. 2 and, in addition to the laser source, it consists of two mode-cleaning cavities [for green and for infrared (IR)], an OPO, a HG-LG mode converter, and a homodyne detection (HD) scheme. The laser source (Diabolo from Innolight) delivers 400 mW of IR (1064 nm) light and 650 mW of green (532 nm) light. The OPO is composed of a bow-tie-shaped cavity in which a $1 \times 2 \times 10$ mm³ type I periodically poled potassium titanyl phosphate (KTP) crystal (Raicol, Inc.) is placed in the smallest beam waist. Our cavity consists of two curved mirrors of 25 mm radius of curvature and two plane mirrors. Three of the mirrors are highly reflective at 1064 nm, $R > 99.95\%$, while the output coupler has a transmittance of $T = 8\%$. At the wavelength of the pump beam (532 nm), the transmittance of the mirrors is larger than 95%. Besides the pump beam, the OPO cavity is seeded with a very dim HG_{10} beam at 1064 nm. The spatial profile of this beam has been tailored in the mode-cleaning cavity, and to enhance the transmission through the cavity, the Gaussian beam from the laser passes first through a phase-flip plate (PP) which produces a relative phase flip of π between the two halves of the Gaussian beam (and thus mimics the HG_{10} mode) [18]. Seeding the cavity with a HG_{10} mode has a twofold purpose: one is to enable an active cavity lock at the frequency of the HG_{10} (which coincides with the frequencies of the HG_{01} , LG_0^{+1} , and LG_0^{-1} modes), and




FIG. 2 (color online). (a) Schematics of the experimental setup to generate amplitude squeezing. (b) Mode converter (MC) for the generation of the local oscillator. The interference process in the MC is illustrated and some of the experimentally obtained interference patterns are shown. Balanced beam splitter (BBS) (50/50 BS); polarizing beam splitter (PBS); half-wave plate (HWP) ($\lambda/2$); quarter-wave plate (QWP) ($\lambda/4$); piezoelectrical element for controlling phases (PZT); phase plate (PP). The prism rotates the HG modes by 45° .

the other one is to ensure the generation of squeezing of the HG_{10} with a small coherent excitation (less than 1 mW). Although the ideal spatial profile of the pump beam for down-conversion efficiency optimization is a superposition of $\text{HG}_{00} + \text{HG}_{20/02}$ mode for the $\text{HG}_{10/01}$ [19], we have chosen to use a Gaussian mode for simplicity reasons, and the resulting decrease in efficiency is overcome by using a more intense beam. The relative phase between the pump and seed is locked to deamplification of the seed beam, thus generating amplitude squeezing.

To prove the existence of quadrature entanglement between the two LG modes, we measure the quadrature quantum noise of the spatial modes in a rotated basis composed of the first-order HG modes, HG_{10} and HG_{01} : By performing a simple basis transformation from the LG modes to HG modes, it is easy to show that $\hat{X}_{\text{HG}_{10}} = (\hat{X}_{\text{LG}_0^{-1}} + \hat{X}_{\text{LG}_0^{+1}})/\sqrt{2}$ and $\hat{X}_{\text{HG}_{01}} = (\hat{P}_{\text{LG}_0^{-1}} - \hat{P}_{\text{LG}_0^{+1}})/\sqrt{2}$, where \hat{X} and \hat{P} are the amplitude and phase quadratures of the modes denoted by the lower indices. According to the criterion of Duan *et al.* [26] and Simon [27], CV entanglement can be witnessed if

$$V(\hat{X}_{\text{LG}_0^{+1}} + \hat{X}_{\text{LG}_0^{-1}}) + V(\hat{P}_{\text{LG}_0^{+1}} - \hat{P}_{\text{LG}_0^{-1}}) < 2, \quad (2)$$

where $V(\cdot \cdot \cdot)$ is the variance. Using the transformation, the criterion reduces to $V(\hat{X}_{\text{HG}_{10}}) + V(\hat{X}_{\text{HG}_{01}}) < 2$, and thus by measuring the amplitude quadrature variances of the two HG modes, entanglement between the OAM modes can be witnessed.

The quadrature variances of the HG modes are analyzed using balanced HD with a spatially tailored local oscillator (LO) mode, which is either a HG_{10} or a HG_{01} mode depending on which signal mode is measured. We produce the HG_{10} LO using the mode-cleaning cavity as described above, and the HG_{01} LO is generated by converting the HG_{10} mode using a prism. The outcome of the HD is fed into a spectrum analyzer which is set to display the noise power (corresponding to the second moment) of the signal beam at 5.5 MHz with a bandwidth of 300 kHz (and 300 Hz averaging). We first calibrate the quantum noise limit (QNL), the result of which is illustrated in Fig. 3 by the horizontal lines. Next we measure the noise traces for the HG_{10} and HG_{01} modes while the phases of the LOs are scanned; see Fig. 3. All data are normalized to the QNL. By fitting the measured data to a theoretical squeezing curve, we find amplitude quadrature squeezing of -1.6 ± 0.2 dB and -1.4 ± 0.2 dB for the HG_{10} and HG_{01} mode, respectively (corresponding to the minima of the curves). Inserting these values into the entanglement criterion we find $V(\hat{X}_{\text{HG}_{10}}) + V(\hat{X}_{\text{HG}_{01}}) = 1.42 \pm 0.01 < 2$.

The measured squeezing values are degraded by the various inefficiencies of our setup. We estimate this efficiency to be $\eta_{\text{total}} = \eta_{\text{cav}} \eta_{\text{prop}} \eta_{\text{det}} \eta_{\text{hd}}$, where $\eta_{\text{prop}} = 0.97 \pm 0.02$ is the measured propagation efficiency, $\eta_{\text{det}} = 0.90 \pm 0.05$ is the measured photodiode (Epitaxx ETX500) efficiency, $\eta_{\text{cav}} = 0.94$ is the estimated cavity escape effi-




FIG. 3 (color online). Experimental squeezing traces for the (a) HG_{01} and (b) HG_{10} modes, where the relative phase between the LO and the squeezed beam is scanned. We measured -1.6 ± 0.2 dB of squeezing and $+4.0 \pm 0.2$ dB of antisqueezing for the HG_{10} mode, -1.4 ± 0.2 dB of squeezing and $+4.1 \pm 0.2$ dB of antisqueezing for the HG_{01} mode. (c) Illustration of the uncertainty volume on the orbital Poincaré sphere and associated with the orbital parameters. The relative elongation of the volume is based on the measured values (however, its size relative to the sphere is not in scale). The projection on the O_2 - O_3 plane is also shown.

ciency, and $\eta_{\text{HD}} = 0.96 \pm 0.02$ is the measured spatial overlap efficiency in the homodyne detector. The total estimated detection efficiencies for our experiment are therefore $\eta_{\text{total}} = 0.79 \pm 0.04$. From these efficiencies we can infer the following squeezing values: -2.2 ± 0.2 dB and -1.9 ± 0.2 dB for the HG_{10} and HG_{01} modes, respectively, and the entanglement criterion is $V(\hat{X}_{\text{HG}_{10}}) + V(\hat{X}_{\text{HG}_{01}}) = 1.25 \pm 0.01 < 2$. We thus have proven that the spatially nondegenerate OPO produces quadrature entanglement between the first-order OAM modes.

We now proceed by characterizing the variances of the orbital parameters that define the position of the state on the orbital Poincaré sphere. First we note that the orbital parameters can be linearized for our setup by decomposing the field operators into a part representing the coherent excitation and a part representing the quantum noise, $\hat{A}_{\text{HG}_{01}} = \langle \hat{A}_{\text{HG}_{01}} \rangle + \delta \hat{A}_{\text{HG}_{01}}$ and $\hat{A}_{\text{HG}_{10}} = \langle \hat{A}_{\text{HG}_{10}} \rangle + \delta \hat{A}_{\text{HG}_{10}}$, and noting that $\langle \hat{A}_{\text{HG}_{01}} \rangle = 0$ and $\langle \hat{A}_{\text{HG}_{10}} \rangle \neq 0$:

$$\begin{aligned} \delta \hat{O}_1 &= \langle \hat{A}_{\text{HG}_{\text{HG}_{10}}} \rangle^2 \delta \hat{X}_{\text{HG}_{10}} & \delta \hat{O}_2 &= \langle \hat{A}_{\text{HG}_{\text{HG}_{10}}} \rangle^2 \delta \hat{X}_{\text{HG}_{01}} \\ \delta \hat{O}_3 &= \langle \hat{A}_{\text{HG}_{\text{HG}_{10}}} \rangle^2 \delta \hat{P}_{\text{HG}_{01}}, \end{aligned} \quad (3)$$

where we have used the definitions $\delta \hat{X} = \delta \hat{A} + \delta \hat{A}^\dagger$ and $\delta \hat{P} = -i(\delta \hat{A} - \delta \hat{A}^\dagger)$. A similar simplification of the quantum OAM states addressing other regimes has been independently formulated in Ref. [28]. We see that in the regime where linearization is valid, the orbital operators




FIG. 4 (color online). (a) Time trace for the amplitude quadrature for spatial modes along a ring on the orbital Poincaré sphere spanned by \hat{O}_2 and \hat{O}_3 . (b) Amplitude quadrature variances for the various spatial modes on the ring. Power measurements of the mean values have been carried out simultaneously to establish the phase reference.

are linear functions of the amplitude and phase quadratures of different spatial modes. Therefore, the precision in determining first-order spatial modes depends on the quadrature noise of the light modes. For example, for a bright excitation in the HG_{10} mode, as in our case, its determination on the sphere is given by the noise in the orthogonal orbital plane spanned by \hat{O}_2 and \hat{O}_3 corresponding to the quadrature noise of the HG_{01} mode [see Eq. (3)]. The quadrature variances of this mode is given in Fig. 3(a). Moreover, the variance of \hat{O}_1 is given by the variance of the amplitude quadrature of the HG_{10} mode given by the minimum squeezing value in Fig. 3(b). Based on these measurements we may define a cigar-shaped uncertainty volume on the orbital Poincaré sphere associated with first-order OAM state as illustrated in Fig. 3(c).

Full tomographic reconstruction of the first-order OAM state is also possible by measuring all possible projections on the sphere using spatially tailored local oscillator modes. Some modes of the LO are made by interfering equally intense HG_{10} and HG_{01} modes with a continuous varying relative phase shift, as shown in Fig. 2(b). By matching these LO modes with the OAM modes, we measure the amplitude quadratures of the modes along a ring on the orbital sphere, thus mapping out the quadrature noise of a whole family of different OAM states. The results of the amplitude quadrature measurements are shown in Fig. 4(a) for a 150 kHz broad signal at 5.5 MHz. We also compute the variance of these data as depicted in Fig. 4(b).

To conclude, we have generated a new quantum state of light composed of quadrature entangled LG modes. For the generation we used an OPO operating in a new regime where all field parameters are degenerate except for its spatial degree of freedom for which it is twofold degenerate. The produced OAM states from the OPO are mapped on an orbital Poincaré sphere which is similar to the standard Poincaré sphere for polarization. We generate a classically bright HG_{10} mode described on this sphere, but its exact location is determined by the noise of the orthogonal HG_{01} dark mode. As this dark state was quadrature squeezed from the OPO, we have demonstrated squeezing in the orbital parameters defining the position of the mode on the orbital Poincaré sphere. In addition we have measured the noise of the bright orbital parameter and thereby reconstructed the uncertainty volume of the first-order OAM state.

We acknowledge financial support from the EU (COMPAS) and the Danish Research Council (FTP). M.L. acknowledge support from the Alexander von Humboldt Foundation.

- [1] L. Allen *et al.*, Phys. Rev. A **45**, 8185 (1992).
- [2] A. S. Chirkin *et al.*, Quantum Electron. **23**, 870 (1993).
- [3] N. Korolkova *et al.*, Phys. Rev. A **65**, 052306 (2002).
- [4] W. P. Bowen *et al.*, Phys. Rev. Lett. **88**, 093601 (2002).
- [5] N. K. Langford *et al.*, Phys. Rev. Lett. **93**, 053601 (2004).
- [6] S. P. Walborn *et al.*, Phys. Rev. A **71**, 053812 (2005).
- [7] A. Mair *et al.*, Nature (London) **412**, 313 (2001).
- [8] V. Boyer *et al.*, Phys. Rev. Lett. **100**, 143601 (2008).
- [9] V. Boyer *et al.*, Science **321**, 544 (2008).
- [10] R. Inoue, N. Kanai, T. Yonehara, Y. Miyamoto, M. Koashi, and M. Kosuma, Phys. Rev. A **74**, 053809 (2006).
- [11] D. V. Vasilyev, I. V. Sokolov, and E. S. Polzik, Phys. Rev. A **77**, 020302 (2008).
- [12] M. J. Padgett and J. Courtial, Opt. Lett. **24**, 430 (1999).
- [13] G. F. Calvo, Opt. Lett. **30**, 1207 (2005).
- [14] J. Janousek *et al.*, arXiv:0812.4686.
- [15] G. S. Agarwal, J. Opt. Soc. Am. A **16**, 2914 (1999).
- [16] S.-K. Choi *et al.*, Phys. Rev. Lett. **83**, 1938 (1999).
- [17] M. Martinelli *et al.*, Phys. Rev. A **67**, 023808 (2003).
- [18] N. Treps *et al.*, Phys. Rev. Lett., **88**, 203601 (2002).
- [19] M. Lassen *et al.*, Phys. Rev. Lett. **98**, 083602 (2007).
- [20] K. Wagner *et al.*, Science **321**, 541 (2008).
- [21] L. A. Wu *et al.*, Phys. Rev. Lett. **57**, 2520 (1986).
- [22] Z. Y. Ou, S. F. Pereira, H. J. Kimble, and K. C. Peng, Phys. Rev. Lett. **68**, 3663 (1992).
- [23] A. S. Villar *et al.*, Phys. Rev. Lett. **95**, 243603 (2005).
- [24] C. Navarrete-Benlloch *et al.*, Phys. Rev. Lett. **100**, 203601 (2008).
- [25] M. Martinelli *et al.*, Phys. Rev. A **70**, 013812 (2004).
- [26] L. M. Duan *et al.*, Phys. Rev. Lett. **84**, 2722 (2000).
- [27] R. Simon, Phys. Rev. Lett. **84**, 2726 (2000).
- [28] M. T. L. Hsu *et al.*, arXiv:0901.4813 [Phys. Rev. A (to be published)].