Technical University of Denmark



Inferring the location and scale of mixing between habitat areas of lesser sandeel through information from the fishery

Jensen, Henrik; Rindorf, Anna; Wright, Peter J.; Mosegaard, Henrik

Published in: ICES Journal of Marine Science

Link to article, DOI: 10.1093/icesjms/fsq154

Publication date: 2011

Document Version Peer reviewed version

Link back to DTU Orbit

Citation (APA):

Jensen, H., Ríndorf, A., Wright, P. J., & Mosegaard, H. (2011). Inferring the location and scale of mixing between habitat areas of lesser sandeel through information from the fishery. ICES Journal of Marine Science, 68(1), 43-51. DOI: 10.1093/icesjms/fsq154

DTU Library Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Inferring the location and scale of mixing between habitat areas of

2 lesser sandeel through information from the fishery

3 Henrik Jensen¹, Anna Rindorf^{1,*}, Peter J. Wright² and Henrik Mosegaard¹

4 Abstract

5 Sandeels are small pelagic fish which play an important role in the diet of a range of 6 natural predators. Due to their low occurrence in traditional survey gear, little is known 7 about their large scale distribution or the degree of mixing between habitat areas. We 8 used detailed information collected directly from the fishery to map fishing grounds and assumed these to reflect the foraging habitat of sandeel. Length distributions from 9 10 individual hauls were used to assess differences in length distributions as a function of the distance between samples. Sandeel foraging habitat covered approximately 5% of 11 12 the total area of the North Sea. Mixing between neighbouring fishing grounds was too low to eliminate differences in length distributions at distances between grounds down 13 to 5 km. Within fishing grounds, mixing was sufficient to eliminate differences in 14 length distributions at scales less than 28 km but insufficient at greater distances. The 15 16 lack of mixing between grounds may result in large differences in sandeel abundance 17 among adjacent fishing grounds. Further, high abundance at one end of a large fishing ground is not necessarily indicative of high abundance at the other end. 18

19 Key words

20 Fishery cooperation, habitat mapping, sandeel, scale of mixing

21 Addresses

¹Technical University of Denmark, National Institute of Aquatic Resources,

23 Charlottenlund Slot, DK2920 Charlottenlund, Denmark.

² Marine Laboratory, Marine Scotland Science, 375 Victoria Road, Aberdeen, AB11

25 9DB, Scotland, U.K.

^{*}Corresponding author. Tel. +45 33 96 33 78. Email: ar@aqua.dtu.dk

27 Introduction

28 Sandeels (Ammodytes spp.) are small semi pelagic fish with a worldwide distribution (Smith and Heemstra, 1986). They usually constitute a high proportion of the fish 29 30 biomass in the regions where they occur (see e.g. Reay, 1970), and are an important 31 prey species for numerous fish, seabirds and mammals (Daan, 1989; Furness, 1990; Wanless et al., 1998). In addition, they are the target of a large scale industrial fishery in 32 the North Sea (ICES, 2008a). This has led to concerns as to whether fisheries pose a 33 threat to top predators through their reduction of the food supply (Macer, 1966; 34 35 Monaghan, 1992; Wright, 1996; Wanless et al., 1998; Engelhard et al., 2008). Most 36 sandeel species inhabit shallow, turbulent sandy areas, located on depths between 20 and 70 m where the content of the finest particles silt and clay is low (Macer, 1966; 37 Reay, 1970; Wright et al., 2000). Because of the limited availability of such substrate 38 (Wright et al., 1998), the distribution of post-settled sandeels is highly patchy (Macer, 39 40 1966; Wright et al., 2000; Freeman et al., 2004; Holland et al., 2005). Post-settled sandeels are rarely found further than 15 km away from known habitat (Wright, 1996; 41 42 Engelhard *et al.*, 2008) and the maximum distance travelled by tagged individuals displaced from grounds was 64 km (Gauld, 1990). This lack of large scale dispersal 43

44 combined with a limited larval exchange between areas (Proctor *et al.*, 1999;

Christensen *et al.*, 2009) means that local aggregations may be vulnerable to depletion
by the fishery and increases the risk of adverse effects on local predators even if the
North Sea stock is inside biologically safe limits. Unfortunately, the lack of knowledge
on the location of the majority of these aggregations in the North Sea and the population
dynamics within them has hindered studies of local depletion except in a few local areas
(Wright, 1996; Rindorf *et al.*, 2000; Daunt *et al.*, 2008; Engelhard *et al.*, 2008).

There are several reasons for the limited knowledge of sandeel distribution and 51 52 population dynamics. Firstly, sandeel bury into the sediment when not feeding in the water column or when approached by predators foraging near the seabed (Winslade, 53 1974a,b,c; Girsa and Danilov, 1976; Pearson et al., 1984; Pinto et al., 1984). This 54 burying behaviour makes their accessibility to sampling in the water column highly 55 variable. Due to this variability and the patchy distribution of habitat, none of the 56 57 regular North Sea acoustic or trawl surveys provide reliable means of mapping sandeel 58 distribution, although both approaches have been used to investigate density in a few areas of the North Sea (Greenstreet et al. 2006, ICES 2008b, Johnsen et al. 2009). 59 60 Secondly, although the requirement for specific habitats is likely to limit sandeel movement, little direct information exists on the extent of horizontal movements of 61 62 post-settled sandeels within or between habitat areas. The few mark-recapture experiments that have been conducted were restricted to small regions because of the 63 64 difficulty of marking a representative number of this abundant species. Furthermore, the 65 recapture location could not be determined since the species is not sorted onboard fishing boats (Gauld, 1990). Consequently, the distribution of habitat areas must be 66

derived from non-conventional methods and movements of this species throughout alarge region such as the North Sea must be assessed from indirect methods.

69 Movements between habitat areas may be detected by differences between length 70 distributions. Local differences in recruitment, growth or mortality will result in 71 differences between areas unless fish mix between different areas. If differences arise 72 from variation in recruitment between areas, subsequent mixing would lead to a 73 decrease in the difference between length distributions over the season. Alternatively, 74 increasing differences in length distributions over the season could result from variation 75 in growth or mortality and lack of mixing between adjacent areas during the season. This increase should be greater the longer the distance between the habitat areas if 76 growth and mortality differences increase with distance. Alternatively, if mixing occurs 77 at small scales while growth and mortality varies independently of distance, differences 78 79 between length distributions should increase at small distances but vary independently 80 of distance at large scales. Based on the extent of identified sandeel aggregations 81 (Freeman et al., 2004; Holland et al., 2005) and the apparent limited movements of settled sandeels it is possible that mixing could be limited both within as well as among 82 83 habitat areas.

The determination of length distributions in local aggregations repeatedly over a longer period of time is a costly exercise if scientific vessels are required to sample the fish. However, the sandeel fishery could potentially provide crucial information at comparatively low costs. The fishery targets foraging sandeel near areas where the sandeels bury and the fishing grounds should therefore provide information on the distribution of sandeel habitat areas (Jensen 2001). Recognizing that information on sandeel habitat areas was crucial to the validity of the scientific advice given, a

collaboration between the Danish Fishermen's Association and the Danish Institute for
Fisheries Research, now Danish Technical University, was started in 1999 in order to
provide the information needed to improve the understanding of North Sea sandeel
population dynamics. Under this collaboration, data were collected by Danish sandeel
skippers directly from the fishing vessels.

The objective of this study was to use the detailed data collected in the fishery to produce a map of the foraging habitat of sandeel and then examine the extent of mixing between and within foraging habitats. Foraging habitats are defined here as areas with potential large densities of non-buried sandeel as sandeel bury when not feeding (Winslade, 1974a). Four predictions were made of the relationship between the difference in length distributions and distance between samples in time and space and the extent of mixing:

If the difference between length distributions increases with distance *within* a
 fishing ground, the degree of mixing between subareas of the fishing ground
 must be limited.

106 2) If the difference between length distributions increases with distance *between*107 fishing grounds, the degree of mixing between fishing grounds must be limited.

3) If the difference between length distributions *decreases* over the season, initial
 differences caused by differences in e.g. recruitment slowly disappear through
 mixing of fish.

4) If the difference between length distributions *increases* over the season, initial
differences are enhanced through changes in growth, mortality or emergence
behaviour and lack of mixing.

114 In both 1) and 2), the distance at which length distributions become significantly

different indicates the distance at which the rate of mixing becomes too low to

116 compensate for differences in recruitment, growth and mortality.

117 Methods

118 Sandeel foraging habitat

119 Three types of information were combined to derive the spatial distribution of foraging habitat: global positioning system (GPS) loggings from individual ships, vessel 120 121 montiroring system (VMS) data and evaluation of maps by fishermen. GPS based 122 computer systems are used by the fishermen together with information from sea charts and past experience to map and log fishing grounds and fishing activities. The maps 123 124 contain data on longitude and latitude, and, in some cases, names of the fishing grounds. 125 This detailed information was used to map about 10% of the grounds. To supplement this data, information about catches of sandeel in individual trawl hauls for which the 126 127 latitude and longitude of hauling were known were used to locate another 50% of the grounds. The remaining 40% of the fishing grounds were obtained from VMS data from 128 Danish vessels fishing sandeel. These data are logged continually while the vessel is at 129 130 sea. This means that the time not spend fishing for sandeel is included and must subsequently be excluded through filtering. These filtering routines were based on 131 distance between succeeding sample points/positions and additional information 132 133 collected from Danish fishermen on the physical characteristics (i.e. depth and seabed type etc.) of each fishing ground and the time of the year the ground is usually fished. 134 From the data collected, the final map of the sandeel fishing grounds was produced 135

using ArcMap and the Spatial Analyst extension (ESRI ArcGIS)(see supplementary

material S1). Each fishing ground is defined as a polygon, adjusted manually 137 138 considering raw data, information about individual grounds, the location of fishing tracks and information about topography. During this process the highest weight was 139 given to the information in the raw data. Fishermen from different harbours evaluated 140 141 the map of fishing grounds, after which it was modified according to the guidelines given by the fishermen. The evaluation by the fishermen resulted in the inclusion of 142 143 additional grounds (from additional navigation data) and deletion of non sandeel grounds. The map was subject to several such evaluations before being finalized. 144

145 Length distributions of sandeels

146 From 1999 to 2008, Skippers from 55 sandeel fishing vessels have collected detailed 147 haul information on 2774 individual trawl hauls. Each fishing vessel records 148 information about the exact location and time of shooting and hauling of the trawl, the 149 name of the fishing ground, and an estimate of the total weight of the catch in each individual trawl haul. Further, a sample of between 0.5 and 1 kg fish is collected from 150 151 each haul and frozen on board. In the laboratory, the sampled fish are thawed, the sandeel are sorted by species, and total length of all fish is measured to nearest half cm 152 153 below. Samples where less than 50 fish were measured were excluded due to the low 154 accuracy of length distributions based on small samples. Further, as the objective was to 155 compare length distributions between sites, hauls with more than 50 km between start and end positions and hauls with a distance greater than 25% of the length of the fishing 156 157 ground were excluded.

To derive an estimate of the variation in length distributions within and between
fishing grounds while accounting for the unbalanced sampling of weeks and vessels,
length distributions were analyzed using generalized linear models of continuation-ratio

logits (Rindorf and Lewy, 2001; Marques *et al.*, 2005). This method allows statistical
testing of the effect of both continuous and discrete variables. Further, by utilizing the
smoothness of length distributions as a function of length this method provides more
accurate estimates of length distributions than traditional methods (Rindorf and Lewy,
2001). The observations analyzed were distributions of observed lengths in half cm for
each of the hauls sampled. They were modelled by fitting fifth degree polynomials to
the continuation-ratio-logits (Rindorf and Lewy, 2001).

All models were fitted using the SAS[®] GENMOD procedure (SAS[®] version 9.1 for

169 Windows; SAS[®] Institute Inc. 2004). The dispersion parameter was estimated by

170 Pearson's χ^2 statistic divided by the degrees of freedom. Only length groups with at

171 least one sandeel on average in a sample were included in tests.

172 Within fishing ground variation

To examine whether length distributions differed within fishing grounds, fishing 173 174 grounds where more than 20 samples were taken in a 2-week period in a given year 175 were selected. This resulted in seven combinations of fishing ground and time. These fishing grounds were then divided into four parts. North-south oriented fishing grounds 176 177 were divided according to latitude whereas east-west oriented grounds were divided 178 according to longitude. Individual trawl hauls were then attributed to each of these subareas based on the midpoints between the shooting and hauling positions. Of the 179 180 fishing grounds selected, Scooter Plads had hauls all originating in the same quarter of 181 the fishing ground and for this fishing ground, this quarter of the fishing ground was 182 divided into two.

The statistical difference between subareas of a fishing ground was estimated using 183 F-tests to compare the deviance of a common length distribution to that of separate 184 length distributions in different subareas (Rindorf and Lewy, 2001), still using fifth 185 degree polynomials to model length distributions. Fishing grounds were analyzed 186 187 separately to assure that the values obtained were independent. From the F-test, a value of F and a probability of this value at the given number of degrees of freedom are 188 189 obtained. If the number of degrees of freedom is unaltered, a larger value of F signifies 190 greater difference between samples.

191 Between fishing ground variation

192 To examine whether length distributions differed between fishing grounds, grounds 193 where more than five samples were taken in a 2-week period were selected. The 194 distances between all hauls were computed and for each pair of fishing grounds in a 2-195 week period, the hauls were grouped in bins of 10 km-distance between hauls. Haul pairs from the same fishing ground or hauls more than 75 km apart were excluded. Two 196 197 models were then fitted for each combination of fishing grounds and distance using the method described in the previous section: one modelled a common length distribution 198 199 and the other modelled two length distributions, one for each fishing ground. From the 200 residual deviances, the F-value of assuming a common length distribution was estimated along with the probability of this value. This resulted in a list (cleaned for doublets) of 201 name of fishing ground 1, name of fishing ground 2, distance-group (10 km groups), F-202 203 values and the probability of all samples being derived from a common length 204 distribution.

205 Trends over time

To examine whether the difference between length distributions generally decreased or increased over time, the correlations between F of the between fishing ground comparison and week of the year were estimated for each distance-group. Only groups where three or more 2-week-periods were sampled in the main fishing season (week 14 to 22) were included.

211 **Results**

212 Sandeel foraging habitat

A total 217 individual habitat areas were identified with a total area of 33 566 km² equivalent to approximately 5% of the area of the North Sea and Skagerrak (Figure 1). The habitat areas vary greatly in size as the area ranges from 1 km² to 4023 km² (Figure 2). Given the average estimated biomass of sandeel in the North Sea (ICES 2008a), this leads to a density within habitat areas of on average 58 ton per km². With an average weight of sandeel of 7 g (ICES 2008a), this corresponds to eight sandeels per m² habitat.

220 Within fishing ground variation

Length distributions of sandeels from 187 hauls taken at five fishing grounds were used (Figure 1). Of these fishing grounds, the number of samples taken exceeded 20 in two consecutive 2-week periods on two of them. It is clear from visual inspections of the length distributions that the variation between fishing grounds is much larger than the within fishing ground variation (Figure 3). The length distributions differed significantly between subareas of the fishing ground on two fishing grounds (a total of three cases, Table 1, Figure 4) and the difference increased significantly with length of

the fishing ground (correlation between F and distance between outer subarea 228 229 midpoints=0.86, P=0.0135). The regression line exceeded an F-value of two (roughly 230 equal to a difference significant at the 5% level when the number of degrees of freedom is large) at a minimum distance between subareas of 28 km (Figure 5). At smaller 231 232 distances, the length distributions were not significantly different and hence mixing most likely occurred at these scales. As the distance refers to the distance between 233 234 fishing ground subarea centres and subareas are defined as quarters of fishing grounds, 235 this corresponds to a total mixing of fish at fishing grounds of lengths less than 112 km.

236 Between fishing ground variation

237 The length distributions differed significantly between fishing grounds in a total of 238 55 cases (90%). Length distributions of sandeels from 1038 hauls taken at 11 fishing 239 grounds were used, resulting in a total of 61 combinations of fishing ground pair and 10 240 km distance group (see Supplementary material S2). The F-value of the difference between length distributions increased significantly with distance between fishing 241 242 grounds (correlation between F and distance group=0.30, P=0.0184)(Figure 6a) and the regression line exceeded an F-value of two at a minimum distance between fishing 243 244 grounds of 5 km. The difference between length distributions was substantially higher 245 between fishing grounds than within fishing grounds even at the same geographical distance (Figure 6b). Whereas within-fishing ground comparisons did not reveal 246 significant differences at distances below 40 km (Figure 5), all of the between-fishing 247 248 ground comparisons made at distances less than 20 km (total of three) and all but one 249 made at distances of less than 30 km (total of 10) were significantly different (Figure 7).

250 Trends over time

251 None of the correlations between F and week were significant at the 5% level when 252 performed separately for each distance group and hence differences did not increase or 253 decrease significantly over time at a given distance. However, the correlation between F 254 and week within a distance group increased significantly with distance (Pearson correlation coefficient=0.93, P=0.0211)(Figure 8). This indicates that there was no 255 256 tendency for nearby grounds to become more similar over time (correlation between F and time was zero) while distant grounds became more different over time (F between 257 258 distant grounds increased over time).

259 **Discussion**

260 This study utilized the first North Sea wide map of lesser sandeel distribution to evaluate dynamics over a complex mosaic of grounds differing widely in size and 261 proximity to each other. The mixing of sandeel between these grounds was insufficient 262 to eliminate significant differences in length distributions at spatial scales down to 5 km. 263 264 Within grounds, the mixing is greater and at fishing grounds with a length smaller than 265 112 km, no difference in length distributions was found. However, beyond this distance, 266 the mixing was again insufficient to eliminate differences in length distributions. This lead to increased differences between fishing grounds late in the season as differences in 267 268 growth and mortality enhanced early season differences. The lack of mixing between 269 fishing grounds potentially increases the risk of adverse effects of the fishery on local 270 predators even if the population at a larger scale is inside biologically safe limits.

The validity of the map as an indicator of sandeel foraging habitat depends on whether all habitat areas are known and fished by the fishery and whether fishing is restricted to foraging areas. Comparison with smaller scale topographical and benthic

274 mapping suggests that the fishery tends to concentrate in areas where sandeel forage, 275 which often coincide with the edge of fishing grounds (Jensen, 2001; Holland et al., 276 2005; Mackinson and van der Kooij, 2006; Engelhard et al., 2008). Given the close proximity between foraging habitat and preferred substrate, fishing distribution should 277 278 provide a good proxy for areas of sandeel habitat (Wright et al., 2000; Van der Kooij et al., 2008). However, vessels only fish in areas with clear tows and a high catch rate and 279 280 avoid shallow waters as well as areas within the territorial limits of countries other than 281 Denmark. Hence, rugged and coastal habitat may be underrepresented in the map. The 282 majority of the catch from coastal habitats is, however, likely to consist of the more coastal sandeel species Ammodytes tobianus (Jensen et al. 2004). 283

284 We are not aware of any similar investigations where mixing has been deduced from comparisons of length distributions. There may be several reasons for this. Firstly, 285 286 comparisons of length distributions is not straight forward when the number of fish 287 measured is limited (Rindorf and Lewy, 2001). Secondly, a difference in length 288 distributions is not equivocal as this may be caused either by lack of mixing and highly 289 variable growth and mortality rates or by size-dependent migration where fish migrate 290 to specific areas when they reach a certain size. Similarly, identical length distributions may be a result of identical recruitment, growth and mortality patterns combined with a 291 292 lack of mixing. The lack of difference in length distribution seen within small fishing 293 grounds is therefore not necessarily a result of mixing but could be a result of 294 homogeneous recruitment pattern combined with low differences in growth and 295 mortality rates. However, as one subarea of the fishing ground did not consistently hold larger fish in the fishing ground sampled in two consecutive 2-week periods (Figure 4), 296 it seems unlikely that size dependent migration is the cause of the difference in length 297

distributions between fishing ground subareas. Furthermore, it seems unlikely that such 298 299 size dependent migration should be larger between fishing grounds than within fishing 300 grounds which would be required to explain the larger differences between grounds. While size-dependent migration may thus occur, it is insufficient to explain the large 301 302 differences between fishing grounds. This is consistent with the very high recapture 303 rates of tagged sandeels recorded on fished grounds (Kunzlik et al., 1986). The recorded 304 differences between length distributions reveal not only lack of extensive movement but 305 also large differences in local recruitment, growth or mortality and possibly also in 306 burying behaviour. Burying behaviour affects the length distribution if the proportion of time spent buried varies between different lengths of sandeel. If this behaviour 307 308 furthermore varies between fishing grounds, the effect can be increased differences over the season. Significant regional differences in length distribution, growth and maturity 309 310 have also been found in other studies (Wright, 1996; Bergstad et al., 2001; Boulcott et al., 2007; Johnsen et al., 2009). The present study provides further explanation for this 311 312 regional variability in dynamics.

In the extreme, local depletion may occur in one end of a large ground while there 313 314 are high sandeel densities in the other end or at one of two neighbouring fishing grounds. This could be caused by fishing if the fishery remains on the low abundance 315 316 end of the ground even when density is higher in the other end, a behaviour which 317 would however not appear to maximise profit. The limited exchange between nearby 318 areas potentially increases the effectiveness of closing local areas to fishing as the 319 sandeel inside the area will not simply 'spill-over' into adjacent fishing grounds and be caught there. However, the relationship between exchange and differences in local 320 densities were not investigated and hence a 'spill-over' effect could potentially occur if 321

322 there were large density differences between two adjacent areas. In this case, lower 323 competition for food in the low-density area may give sandeel energetic reasons to 324 switch area. Furthermore, the present analysis deals only with mixing in the feeding 325 season. Mature sandeel emerge from the sandbanks in December and January to spawn 326 (Gauld and Hutcheon, 1990; Boulcott et al., 2007), and whether mixing takes place during this event is unknown. However, as egg and early larval distributions closely 327 328 match known overwintering habitat, this appears unlikely (Wright and Bailey, 1996; 329 Munk et al., 2002). After hatching, larvae may drift considerable distances (Proctor et 330 al., 1998; Christensen et al., 2009) and the horizontal movements of pre-settled sandeel (Wright, 1996) most likely leads to exchange of recruits between fishing grounds. 331 332 Hence, closing a particular fishing ground will protect sandeel during the fishing season and may contribute to recruitment on other fishing grounds but is unlikely to assure the 333 334 presence of sandeel in the closed fishing ground in coming years.

In conclusion, the distribution of sandeel in the North Sea is highly patchy and there is limited exchange between even close fishing grounds during the fishing season. Some mixing occurs within fishing grounds but the mixing between grounds appears to be very low. Management through closed fishing grounds or fishing ground subareas therefore has the potential to protect local aggregations of sandeel during the fishing season and may result in increased recruitment to nearby fishing grounds, but such a protection is unlikely to assure continuous recruitment to the particular fishing ground.

342 Acknowledgements

We are deeply grateful to the fishing skippers who collected the data and placed them at our disposal, to the Danish Fishing association that supported their work, to Palle Brogaard, who improved the cooperation between the industry and scientists and

to Stina B. S. Hansen and the laboratory technicians who worked up the thousands of

347 length samples. The work of A. Rindorf and H. Mosegaard was supported by the Danish

348 national project 'ETOMTOBIS'. Georg Engelhard and an anonymous reviewer

349 provided valuable comments on an earlier version.

350 Supplementary material

- 351 The following supplementary material is available at *ICESJMS* online: Material S1
- shows the sandeel map and procedures for its derivation whereas Table S2 lists results
- of the comparison of length distributions between fishing grounds.

354 **References**

- Bergstad, O. A., Hoines, A. S. and Kruger-Johnsen, E. M. 2001. Spawning time, age
- and size at maturity, and fecundity of sandeel, *Ammodytes marinus*, in the north-eastern
- North Sea and in unfished coastal waters off Norway. Aquatic Living Resources, 14:

358 293-301.

- 359 Boulcott P., Wright P. J., Gibb F., Jensen H. and Gibb I. 2007. Regional variation in the
- maturation of sandeels in the North Sea. ICES Journal of Marine Science, 64: 369-376.
- 361 Christensen A., Mosegaard H. and Jensen H. 2009. Spatially resolved fish population
- analysis for designing of MPAs: influence on inside and neighbouring habitats. ICES
- 363 Journal of Marine Science, 66: 56-63.
- 364 Daan, N. 1989. Data base report of the stomach sampling project 1981. ICES
- 365 Cooperative Research Report No. 164.
- 366 Daunt, F., Wanless, S., Greenstreet, S. P. R., Jensen, H., Hamer, K. C. and Harris, M. P.
- 367 2008. The impact of the sandeel fishery closure on seabird food consumption,

- distribution, and productivity in the northwestern North Sea. Canadian Journal ofFisheries and Aquatic Sciences, 65: 362-381.
- 370 Engelhard, G. H., Van der Kooij, J., Bell, E. D., Pinnegar, J. K., Blanchard, J. L.,
- 371 Mackinson, S. and Righton, D. A. 2008. Fishing mortality versus natural predation on
- diurnally migrating sandeels *Ammodytes marinus*. Marine Ecology Progress Series, 369:

373 213-227.

- 374 Freeman, S., Mackinson, S. and Flatt, R. 2004. Diel patterns in the habitat utilisation of
- 375 sandeels revealed using integrated acoustic surveys. Journal of Experimental Marine
- Biology and Ecology, 305: 141-154.
- Furness, R. W. 1990. A preliminary assessment of the quantities of Shetland sandeels
 taken by seabirds, seals, predatory fish and the industrial fishery in 1981–83. Ibis, 132:
 205–217.
- Gauld, J. A. 1990. Movements of lesser sandeels (*Ammodytes marinus* Raitt) tagged in
 the northwestern North Sea. Journal du Conseil International pour l'Exploration de la
 Mer., 46: 229-231.
- Gauld, J. A. and Hutcheon, J. R. 1990. Spawning and fecundity in the lesser sandeel, *Ammodytes marinus* Raitt, in the north-western North Sea. Journal of Fish Biology, 36:
- **385** 611-613.
- 386 Girsa, I. I., and Danilov, A. N. 1976. Defensive behaviour in sand lance, Ammodytes
- *hexapterus* of the White Sea. Journal of Ichthyology, 16: 862-865.
- 388 Greenstreet, S. P. R., Armstrong, E., Mosegaard, H., Jensen, H., Gibb, I. M., Fraser, H.
- 389 M., Scott, B. E., Holland, G. J. and Sharples, J. 2006. Variation in the abundance of

- 390 sandeels Ammodytes marinus off southeast Scotland: an evaluation of area-closure
- 391 fisheries management and stock abundance assessment methods. ICES Journal of
- 392 Marine Science, 63: 1530-1550.
- Holland, G. J., Greenstreet, S. P. R., Gibb, I. M., Fraser, H. M., and Robertson, M. R.
- 394 2005. Identifying sandeel Ammodytes marinus sediment habitat preferences in the
- marine environment. Marine Ecology Progress Series, 303: 269–282.
- 396 ICES 2008a. Report of the Working Group on the Assessment of the Demersal Stocks
- in the North Sea and Skagerrak. ICES C.M. 2008/ACOM:09. 921 pp.
- 398 ICES 2008b. Report of the Ad Hoc Group on Sandeel (AGSAN) ICES C.M.
- 399 2008/ACOM:59. 68 pp.
- 400 Jensen, H. 2001. Settlement dynamics in the lesser sandeel Ammodytes marinus in the
- 401 North Sea. Ph.D. dissertation, University of Aberdeen, 230 pp.
- 402 Jensen, H., Kristensen, P. S., and Hoffmann, E. 2004. Sandeels in the wind farm area at
- 403 Horns Reef. Danish Institute for Fisheries Research, Depart of Marine Fisheries,
- 404 Charlottenlund Castle, 2920 Charlottenlund. Report to ELSAM, August 2004, 45 pp.
- 405 Johnsen, E., Pedersen, R., and Ona, E. 2009. Size-dependent frequency response of
- 406 sandeel schools. ICES Journal of Marine Science, 66: 1100–1105.
- 407 Kunzlik, P. A., Gauld, J. A. and Hutcheon, J. R. 1986. Preliminary results of the
- 408 Scottish sandeel tagging project, ICES CM 1986/G:7, pp. 8.

- 409 Macer, C.T. 1966. Sand eels (Ammodytidae) in the south -western North Sea; their
- 410 biology and fishery. Fishery Investigations. Series 2: Sea Fisheries. Ministry of
- 411 Agriculture, Fisheries and Food, London., 24: 1-55.
- 412 Mackinson, S. and Van der Kooij, J. 2006. Perceptions of fish distribution, abundance
- 413 and behaviour: Observations revealed by alternative survey strategies made by scientific
- and fishing vessels. Fisheries Research, 81: 306-315.
- 415 Marques, V., Chaves, C., Morais, A., Cardador, F. And Stratoudaskis, Y. 2005.
- 416 Distribution and abundance of snipefish (*Macroraphosus* spp.) off Portugal (1998-
- 417 2003). Scientia Marina, 69: 563-576.
- 418 Monaghan, P., 1992. Seabirds and sandeels: the conflict between exploitation and
- 419 conservation in the northern North Sea. Biodiversity and Conservation, 1: 98-111.
- 420 Munk, P., Wright, P. J. and Pihl, N. J. 2002. Distribution of the Early Larval Stages of
- 421 Cod, Plaice and Lesser Sandeel across Haline Fronts in the North Sea. Estuarine,
- 422 Coastal and Shelf Science, 55: 139-149.
- 423 Pearson, W. H., Woodruff, D. L., and Sugarman, P. C. 1984. The burrowing behavior of
- 424 sand lance, Ammodytes americanus: Effects of oil-contaminated sediment. Marine
- 425 Environmental Research, 83: 193-204.
- 426 Pinto, J. M., Pearson, W. H., and Anderson J. H. 1984. Sediment preferences and oil
- 427 contamination in the Pacific sand lance *Ammodytes hexapterus*. Marine Biology, 83:
 428 193-204.
- 429 Proctor R., Wright P. J. and Everitt A. 1998. Modelling the transport of larval sandeels
- 430 on the north-west European shelf. Fisheries Oceanography, 7: 347-354.

- 431 Reay P.J. 1970. Synopsis of biological data on North Atlantic sandeels of the genus
 432 *Ammodytes*. FAO Fisheries Synopsis, No. 82.
- 433 Rindorf, A., and Lewy P. 2001. Analyses of length and age distributions using
- 434 continuation-ratio logits. Canadian Journal of Fisheries and Aquatic Sciences, 58: 1141-

435 1152.

- 436 Rindorf, A., Wanless, S. and Harris, M. P. 2000. Effects of changes in sandeel
- 437 availability on the reproductive output of seabirds. Marine Ecology Progress Series,

438 202: 241-252.

- 439 SAS Institute Inc. 2004. What's New in SAS/STAT 9 and 9.1. SAS OnlineDoc® 9.1.3.
- 440 Cary, NC: SAS Institute Inc.
- Smith, M. M., and Heemstra, P. C. 1986. (eds) Smiths' sea fishes. Springer-Verlag,
 New York. 1047 pp.
- 443 Van der Kooij, J., Scott, B. E. and Mackinson, S. 2008. The effects of environmental
- 444 factors on daytime sandeel distribution and abundance on the Dogger Bank. Journal of
- 445 Sea Research, 60: 201-209.
- 446 Wanless, S., Harris, M.P. and Greenstreet, S.P.R. 1998. Summer sandeel consumption
- 447 by seabirds breeding in the Firth of Forth, south-east Scotland. ICES Journal of Marine
- 448 Science, 55: 1141-1151.
- 449 Winslade, P. 1974a. Behavioural studies on the lesser sandeel Ammodytes marinus
- 450 (Raitt) I. The effect of food availability on activity and the role of olfaction in food
- 451 detection. Journal of Fish Biology, 6: 565-576.

- 452 Winslade, P. 1974b. Behavioural studies on the lesser sandeel Ammodytes marinus
- 453 (Raitt) II. The effect of light intensity on activity. Journal of Fish Biology, 6: 577-586.
- 454 Winslade, P. 1974c. Behavioural studies on the lesser sandeel Ammodytes marinus
- 455 (Raitt) III. The effect of temperature on activity and the environmental control of the
- 456 annual cycle of activity. Journal of Fish Biology, 6: 587-599.
- 457 Wright P. J., Jensen H. and Tuck I. 2000. The influence of sediment type on the
- distribution of the lesser sandeel, *Ammodytes marinus*. Journal of Sea Research, 44:
 243-256.
- 460 Wright, P. J. 1996. Is there a conflict between sandeel fisheries and seabirds? *In*:,
- 461 Aquatic predators and their prey, pp. 154-165. Ed. by S. P. R. Greenstreet and M. L.
- 462 Tasker. Fishing News Books, Blackwell Science, Oxford. 191 pp.
- 463 Wright, P. J. and Bailey, M. C. 1996. Timing of hatching in *Ammodytes marinus* from
- 464 Shetland waters and its significance to early growth and survivorship. Marine Biology,465 126: 143-152.
- 466 Wright, P. J., Pedersen, S. A., Donald, L., Anderson, C., Lewy, P. and Proctor, R. 1998.
- 467 The influence of physical factors on the distribution of lesser sandeel, *Ammodytes*
- 468 *marinus* and its relevance to fishing pressure in the North Sea, ICES CM 1998/AA:3. 18
- 469 pp.

Table 1. Comparison of length distributions between subareas within fishing grounds.
Length indicates maximum distance between midpoints of subareas sampled. Total
deviance and df indicates the deviance from a common length distribution and the
degrees of freedom of this common length distribution. Subarea deviance indicates the
reduction in deviance obtained by modelling the length distribution in each subarea
separately and the associated loss of degrees of freedom.

Fishing ground	Length	Total		Subarea			
	(km)	deviance	df	deviance	df	F	P(F)
Scooter Plads	6	830.8	329	22.7	6	1.498	0.1780
Berwick Bank	13	1 191.7	281	46.8	7	1.576	0.1422
Berwick Bank	13	805.3	257	10.9	6	0.580	0.7464
Inner Shoal	25	1 166.3	468	27.2	6	1.819	0.0936
Sorel	41	806.4	307	98.7	10	3.758	0.0001
Elbow Spit	121	1 205.7	337	260.5	15	4.854	< 0.0001
Elbow Spit	121	1 635.8	341	336	19	3.686	< 0.0001

479 **Captions**

480 Fig. 1. Sandeel habitat areas (areas with potential high density of non-buried sandeel)481 and named fishing grounds referred to in the text.

- 482 Fig. 2. Frequency distribution of area (in km^2) of individual sandeel habitat areas.
- 483 Lower axis indicates lower limit of 100 km²-intervals..
- 484 Fig. 3. Length distributions at fishing grounds where more than 20 hauls were taken in a

485 2-week period. a: Scooter Plads (maximum distance between subarea midpoints 6 km),

b and c: Berwick bank (maximum distance between subarea midpoints 13 km) week 22

and 24, respectively, d: Inner Shoal (maximum distance between subarea midpoints 25

488 km), e: Sorel (maximum distance between subarea midpoints 41 km), f and g: Elbow

489 Spit (maximum distance between subarea midpoints 121 km) week 16 and 18,

490 respectively. Colours indicated subarea of fishing ground.

491 Fig. 4. Length distributions in different subareas. a: Sorel, b: Elbow Spit, week 16, c:

492 Elbow spit week 18. Solid black lines: subarea immediately west of centre, hatched

493 black lines: subarea close to the western end of the fishing ground, solid grey lines:

494 subarea immediately east of centre, hatched grey lines: subarea close to the eastern end

495 of the fishing ground.

Fig. 5. F-value of comparison of common and subarea-specific length distributions as a
function of the distance between sampled fishing ground subareas. Line is a regression
line.

499 Fig. 6. F-value of comparison of common and fishing-ground-specific length

500 distributions as a function of the distance between sampled fishing grounds (a) and a

501 comparison of within and between fishing ground results (b). Line is a regression line.

- 502 Filled symbols, solid line: between fishing ground comparisons. open symbols, hatched
- 503 line: within fishing ground comparisons. Note that the values on the y-axis differs
- 504 between figures as b shows only the lower half of a.
- 505 Fig.7. Proportion of analyses showing significant differences as a function of distance.
- 506 Filled symbols, solid line: between fishing grounds, open symbols, hatched line: within
- 507 fishing grounds.
- 508 Fig. 8. Correlation between week and F for given distance. Positive values indicate a
- 509 general rise in F over the season, negative a decrease. Line indicates regression line.

510 Fig. 1.





514 Fig. 2.



516 Fig. 3.



























Fig. 8. Correlation between week and F for given distance. Positive values indicate ageneral rise in F over the season, negative a decrease. Line indicates regression line.



Deriving the map of sandeel habitats (S1)

From the data collected (Figure S1), the final map of the sandeel fishing grounds was produced using the following 5-step procedure:

1) A point shapefile with the data was created from a textfile using ArcMap (ESRI ArcGIS).

2) A kernel density analysis (Silverman, 1986) was applied on the shapefile using a quadratic kernel function to fit a smoothly tapered surface to each point (search radius: 0.015 decimal degrees, output cell size: 0.006 decimal degrees) and produce a grid using the Spatial Analyst extension with ArcGIS.

From this grid, a new grid including only cells with a density of points greater than 50
 000 was produced. This grid was afterwards converted to a shapefile so that each fishing ground is defined as a polygon.

4) The polygons were adjusted manually considering raw data, information about individual grounds, the location of fishing tracks and information about topography. During this process the highest weight was given to the information in the raw data.

Lastly, fishermen from different harbours evaluated the map of fishing grounds, after which it was modified according to the guidelines given by the fishermen. The evaluation by the fishermen resulted in the inclusion of additional grounds (from additional navigation data) and deletion of non sandeel grounds. The map was subject to several such evaluations before the final map was accepted (Figure S2).

References

Silverman, B.W. 1986. Density Estimation for Statistics and Data Analysis. New York: Chapman and Hall, 175 pp.



Fig. S1. Map of raw data from various GPS based navigation systems.



Fig. S2. Sandeel habitat areas.

Table S2. Results of the comparison of length distributions between fishing grounds. Total deviance and df indicates the deviance from a common length distribution and the degrees of freedom of this common length distribution. Ground specific deviance indicates the reduction in deviance obtained by modelling the length-distribution in each area separately and the associated loss of degrees of freedom. Distance groups are 10 km groups denoted by the lowest distance in the group (e.g. 0 equals the group 0-10 km).

Total

		Distance						
Ground 1	Ground 2	group	deviance	df	deviance	df	F	P(F)
N.W. Rough	Stendysse/Lissis Revle	0	416	159	52	5	4.11	0.0012
N.W. Rough	Stendysse/Lissis Revle	10	433	174	53	5	4.35	0.0007
N.W. Rough	Stendysse/Lissis Revle	10	1 150	179	249	6	6.67	< 0.0001
N.W. Rough	Southernmost Rough	20	430	153	43	6	2.64	0.0159
N.W. Rough	Southernmost Rough	20	995	239	84	6	3.45	0.0022
N.W. Rough	Southernmost Rough	20	1 112	196	81	6	2.45	0.0234
N.W. Rough	Stendysse/Lissis Revle	20	474	160	120	5	8.39	< 0.0001
N.W. Rough	Stendysse/Lissis Revle	20	892	228	36	6	1.57	0.1524
Rute 18	N.W. Rough	20	278	129	42	5	4.08	0.0014
Scooter Plads	Wee Bankie - S.W.	20	1 243	382	284	6	14.79	< 0.0001
N.W. Rough	Southernmost Rough	30	482	168	56	6	3.36	0.0030
N.W. Rough	Southernmost Rough	30	1 291	259	63	6	2.15	0.0455
N.W. Rough	Southernmost Rough	30	640	186	36	7	1.56	0.1430
N.W. Rough	Stendysse/Lissis Revle	30	736	178	48	6	2.01	0.0614
Rute 18	N.W. Rough	30	522	179	74	5	5.25	0.0001
Sorel	N.W. Rough	30	1 723	255	1 141	6	28.81	< 0.0001
Sorel	N.W. Rough	30	791	164	30	5	1.29	0.2664
Sorel	Stenkanten v. Sorel	30	513	204	118	6	8.06	< 0.0001

Ground specific

Middle Rough	Sorel	40	679	177	101	5	5.40	0.0001
N.W. Rough	Southernmost Rough	40	557	157	148	6	7.20	< 0.0001
N.W. Rough	Southernmost Rough	40	425	153	47	6	2.91	0.0086
N.W. Rough	Southernmost Rough	40	1 447	270	534	6	16.96	< 0.0001
N.W. Rough	Southernmost Rough	40	1 937	388	64	6	2.17	0.0432
N.W. Rough	Southernmost Rough	40	1 313	324	272	7	9.81	< 0.0001
Rute 18	Stendysse/Lissis Revle	40	617	159	157	5	8.36	< 0.0001
Sorel	Lisborgs Revle	40	663	209	185	6	9.99	< 0.0001
Sorel	N.W. Rough	40	2 098	313	1 344	6	34.07	< 0.0001
Sorel	N.W. Rough	40	849	197	191	6	7.62	< 0.0001
Sorel	N.W. Rough	40	985	248	47	5	2.40	0.0356
Sorel	Stenkanten v. Sorel	40	554	218	135	6	9.08	< 0.0001
Middle Rough	Sorel	50	1 148	300	234	5	12.42	< 0.0001
N.W. Rough	S.W. Spit	50	434	140	63	6	3.56	0.0019
N.W. Rough	Southernmost Rough	50	580	158	143	6	6.74	< 0.0001
N.W. Rough	Southernmost Rough	50	425	138	34	6	1.94	0.0736
N.W. Rough	Southernmost Rough	50	1 591	285	424	6	12.93	< 0.0001
N.W. Rough	Southernmost Rough	50	1 762	341	72	6	2.36	0.0284
N.W. Rough	Southernmost Rough	50	2 090	394	575	6	18.35	< 0.0001
N.W. Rough	Southernmost Rough	50	1 374	236	280	6	8.21	< 0.0001
Rute 18	Southernmost Rough	50	618	168	97	5	5.45	0.0001
Sorel	Lisborgs Revle	50	663	209	185	6	9.99	< 0.0001
Sorel	N.W. Rough	50	2 873	441	1 652	6	42.85	< 0.0001
Sorel	N.W. Rough	50	813	197	138	6	5.73	< 0.0001
Sorel	N.W. Rough	50	820	192	32	5	1.53	0.1788
Sorel	Stenkanten v. Sorel	50	382	154	71	6	4.94	0.0001
Middle Rough	Sorel	60	1 110	299	231	5	12.64	< 0.0001
N.W. Rough	Southernmost Rough	60	468	130	118	6	5.71	< 0.0001
N.W. Rough	Southernmost Rough	60	1 447	270	534	6	16.96	< 0.0001
N.W. Rough	Southernmost Rough	60	1 584	291	110	6	3.44	0.0022

N.W. Rough	Southernmost Rough	60	1 996	380	664	6	21.39	< 0.0001
N.W. Rough	Southernmost Rough	60	1 374	236	280	6	8.21	< 0.0001
Rute 18	Southernmost Rough	60	618	168	97	5	5.45	0.0001
Sorel	Lisborgs Revle	60	527	153	131	5	7.85	< 0.0001
Sorel	N.W. Rough	60	2 819	441	1 619	6	42.78	< 0.0001
Sorel	N.W. Rough	60	948	232	222	6	9.29	< 0.0001
Sorel	N.W. Rough	60	1 181	293	91	5	4.61	0.0004
Middle Rough	Sorel	70	872	230	273	5	14.71	< 0.0001
N.W. Rough	Southernmost Rough	70	943	139	493	5	15.05	< 0.0001
N.W. Rough	Southernmost Rough	70	986	188	157	6	5.15	< 0.0001
Sorel	N.W. Rough	70	2 975	469	1 720	6	45.76	< 0.0001
Sorel	N.W. Rough	70	990	244	252	6	10.60	< 0.0001
Sorel	N.W. Rough	70	1 181	293	91	5	4.61	0.0004