

Technical University of Denmark



Changes in distribution and lengths of *Mnemiopsis leidyi* in the central Baltic Sea between fall and spring

Storr-Paulsen, Marie; Huwer, Bastian

Published in:
Aquatic Invasions

Link to article, DOI:
[10.3391/ai.2008.3.4.9](https://doi.org/10.3391/ai.2008.3.4.9)

Publication date:
2008

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Storr-Paulsen, M., & Huwer, B. (2008). Changes in distribution and lengths of *Mnemiopsis leidyi* in the central Baltic Sea between fall and spring. *Aquatic Invasions*, 3(4), 429-434. DOI: 10.3391/ai.2008.3.4.9

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Changes in distribution and lengths of *Mnemiopsis leidyi* in the central Baltic Sea between fall and spring

Marie Storr-Paulsen and Bastian Huwer

DTU Aqua, National Institute of Aquatic Resources, Technical University of Denmark, Charlottenlund Castle, DK-2920

Charlottenlund, Denmark

E-mails: mstp@aqu.dtu.dk or bhu@aqu.dtu.dk

Received 14 October 2008; accepted in revised form 26 November 2008; published online 18 December 2008

Abstract

In March 2008, we conducted a survey to investigate the distribution and abundance of the invasive ctenophore *Mnemiopsis leidyi* in the Bornholm Basin after the winter period. Compared to the situation in November 2007 the centre of the distribution had shifted towards the deeper parts of the Basin where temperatures were higher than at the surface. Furthermore, we found a decrease in average size from $18.6 \text{ mm} \pm 7.6 \text{ SD}$ in November to an average of $10.5 \text{ mm} \pm 4.9 \text{ SD}$ in March, which may indicate the emergence of a new generation of *M. leidyi* between the two sampling dates.

Key words: *Mnemiopsis leidyi*, Ctenophora, invasive species, Baltic Sea, Bornholm Basin

Since the first observations of *Mnemiopsis leidyi* A. Agassiz 1865 in Northern Europe (Faasse and Bayha 2006; Hansson 2006; Javidpour et al. 2006; Oliveira 2007), several studies have described occurrences of this invasive lobate ctenophore in the Baltic, spanning from Kattegat to the Bothnian Sea (Haslob et al. 2007; Janas and Zgrundo 2007; Kube et al. 2007; Lehtiniemi et al. 2007; Tendal et al. 2007; Javidpour et al. 2008; Viitasalo et al. 2008). In November 2007, Huwer et al. (2008) investigated the horizontal and vertical distribution of *Mnemiopsis leidyi* in the central Baltic Sea. The studied area covered the Bornholm Basin, an important spawning ground for cod and sprat (Köster et al. 2005). Four months later, in March 2008, we conducted a survey in the same area to investigate if *M. leidyi* had survived the winter and how the distribution and abundances of this invasive species had changed during wintertime.

M. leidyi were sampled during a cruise with the Danish research vessel 'DANA' from March 1st to 17th 2008, on a station grid covering the Bornholm Basin. Additional sampling was conducted on a transect through the Arkona Basin consisting of 6 stations (Figure 1). To assess abundances and the horizontal distribution of *M. leidyi*, hauls with a Bongo net were taken on a total of 75 stations. The gear was equipped with two nets of different mesh size ($335 \mu\text{m}$ and $500 \mu\text{m}$) and with flowmeters to measure the volume of filtered water. Double oblique hauls from the surface to 2 meter above the seafloor were conducted at a towing speed of 3 knots.

Upon retrieval of the gear, specimens of *M. leidyi* were immediately sorted from the samples, counted and the body oral-aboral length of each individual was measured to the nearest 1.0 mm with a sliding caliper. All samples were analysed within 30 minutes after capture. Counts of




Figure 1. *M. leidyi*. Horizontal distribution and abundances (ind. m⁻²) sampled with a Bongo net in March 2008. Station numbers are identical with the numbers in Annex 1

M. leidyi were standardised to 1 m² surface area by accounting for the filtered water volume and the maximum depth of the tow.

A total of 208 and 261 specimens of *M. leidyi* were caught in the 335 µm and 500 µm net, respectively. There was no significant difference ($P = 0.634$) between the size ranges of specimens caught with the two different mesh sizes, as average lengths were $10.2 \text{ mm} \pm 4.6 \text{ SD}$ and $10.7 \text{ mm} \pm 5.1 \text{ SD}$ for the 335 µm and 500 µm net, respectively. Accordingly, samples from the two nets have been pooled in the further analysis.

For a comparison of length distributions of *M. leidyi* between fall and spring, length data from November 2007 were used. The sampling procedure for these data is described in Huwer et al. (2008).

Horizontal distribution and abundance

Low temperatures have been found to be a limiting factor for the distribution and growth of *M. leidyi* (Kremer 1994; Mutlu 1999; Purcell et al. 2001; Shiganova et al. 2001; Purcell and Decker 2005). Therefore, we were interested in

investigating the abundance and distribution of *M. leidyi* in the central Baltic after the winter period.

Huwer et al. (2008) found *M. leidyi* to be present over large areas of the central Baltic Sea in November 2007. In the present study in March 2008 the species was again distributed over large parts of the area. However, in comparison to the situation in fall 2007, the centre of the distribution had shifted to the deeper central parts of the Bornholm Basin (Figure 1). During the present study in spring 2008, no ctenophores were found on the 18 stations with depths <50 meters. The abundances of *M. leidyi* increased with increasing depth and almost 90% of the specimens were registered on stations with depths >70 meter (Table 1).

A possible reason for this change may be colder winter temperatures above the halocline in the Bornholm Basin. Figure 2 shows temperature and salinity profiles from a station in the central Bornholm Basin (Station 23 in Figure 1 and Annex 1; 55.292°N, 15.750°E) in November 2007 and March 2008. The situation in March showed a well mixed surface layer with a

temperature of 4°C, and higher temperatures of 6 to 8°C in the bottom layer below 50 m depth. In November no such distinct temperature difference was detected and the surface water was much warmer with temperatures of about 9.5°C. In contrast to temperature, salinity did not seem to have an influence on the change in distribution in the present study. The salinity profiles showed little differences between fall and spring due to the permanent halocline in the Bornholm Basin.

Kube et al. (2007) have shown that *M. leidyi* was able to survive the winter in the Baltic Sea. Even though abundances at a sampling station in the Bornholm Basin were generally low, they found a decrease in abundances from February to May.

Table 1. Abundances of *M. leidyi* at different depth strata in March 2008

Depth	Stations (n)	<i>M. leidyi</i> (n)	<i>M. leidyi</i> (%)	Average <i>M. leidyi</i> m ⁻²
< 50 m	18	0	0%	0
50-59 m	12	18	3.8%	0.12±0.36
60-69 m	16	41	8.7%	0.19±0.44
70-79 m	17	178	38.0%	0.68±1.19
> 80 m	12	232	49.4%	1.37±0.98

Length distribution

Specimens of *M. leidyi* were significantly (P<0.001) larger in November 2007 than in March 2008. *M. leidyi* between 6-59 mm (average 18.6 mm ± 7.6 SD) and 3-27 mm (average 10.5 mm ± 4.9 SD) were caught in November 2007 and March 2008, respectively (Figure 3). Javidpour et al. (2008) investigated seasonal changes and population dynamics of *M. leidyi* in Kiel Fjord in the western Baltic Sea. Abundance peaks were observed in late summer-early autumn, which is in good correspondence to population dynamics in native habitats of *M. leidyi* (Kremer 1994). Javidpour et al. (2008) further describe seasonal shifts in the dominant size classes and conclude that peak reproduction in Kiel Fjord takes place in late summer and fall. However, their finding of a higher larvae: adult ratio from October 2006 to January 2007 may also indicate some reproduction in the winter period. In the present study, occurrence of smaller individuals in spring could be interpreted as the outcome of a new generation during fall or winter. This may indicate that successful




Figure 2. Temperature and salinity profiles in the central Bornholm Basin (station 23 in Figure 1 and Annex 1) in November 2007 and March 2008

reproduction and overwintering also occurs in the Bornholm Basin. However, because ctenophores can hardly be aged and are known to shrink at low food abundances (Reeve et al. 1989, Purcell et al. 2001), smaller size ranges in March could be due to starvation during winter-time. Another possible explanation for the shift in the length distribution may be passive transport of smaller individuals into the Bornholm Basin. A drift modeling study has shown that such a transport, e.g. from areas in the western Baltic, could be possible within few months (Postel et al. 2007).




Figure 3. *M. leidyi*. Relative size distribution of ctenophores sampled in March 2008 (dark grey bars, n=469) and November 2007 (light grey bars, n=364; from Huwer et al. 2008)

The maximum abundance of *M. leidy* in the Bornholm Basin in November 2007 was 8.92 ind. m⁻², with an average level of 1.58 ± 2.12 ind. m⁻² (Huwer et al. 2008). In the present study four months later, abundances were lower with a maximum abundance of 4.83 ind. m⁻², and an average level of 0.43 ± 0.86 ind. m⁻².

However, even though abundances had decreased from fall to spring, *M. leidy* has survived the second winter after its introduction to the central Baltic.

Conclusions

It can be concluded that *M. leidy* was still present in the Bornholm Basin after the winter 2007/2008 and that the size distribution had shifted to smaller specimens, which may indicate the emergence of a new generation. Furthermore, there has been a shift in the horizontal distribution towards deeper parts of the Basin from fall to spring. Accordingly, *M. leidy* may migrate from the colder surface layer to the warmer deep water during winter, while expanding its distribution to warmer upper layers and coastal areas during summer and fall. A similar temperature dependent vertical distribution was reported by Javidpour et al. (2008) for the *M. leidy* population in Kiel Fjord. Other possible reasons for the changing distribution might be passive advection or differences in food availability. Thus, the present study provides further support for the hypothesis that *M. leidy* uses the deep layers of the Bornholm Basin as overwintering refuge (Kube et al. 2007; Huwer et al. 2008).

However, to obtain a better understanding of *M. leidy* distribution and population dynamics in the central Baltic, a temporally explicit monitoring is highly advisable. *M. leidy* is known to prey on fish eggs and larvae (Cowan and Houde 1993; Purcell and Arai 2001; Purcell et al. 2001) and to compete with fish for zooplankton prey. Therefore, the seasonal development of this invasive predatory species receives special interest in the light of spatio-temporal interactions with ichthyoplankton in the Bornholm Basin (Haslob et al. 2007; Huwer et al. 2008). This area is at present the most important spawning ground for Eastern Baltic cod, a fish stock already suffering from unfavorable environmental conditions and heavy exploitation (Köster et al. 2005).

Acknowledgements

This study has been carried out with support of the National Institute of Aquatic Resources at the Technical University of Denmark and the Commission of the European Communities, specifically SSP8-2004-513670, PROTECT. We do appreciate all the extra work conducted by the crew and scientific staff on 'DANA' during the survey especially the help of Thomas Warnar.

References

- Claussen DL, Hopper RA, Sanker AM (2000) The effects of temperature, body size, and hydration state on the terrestrial locomotion of the crayfish *Orconectes rusticus*. *Journal of Crustacean Biology* 20: 218-223
- Cowan JH Jr., Houde ED (1993) Relative predation potentials of scyphomedusae, ctenophores and planktivorous fish on ichthyoplankton in Chesapeake Bay. *Marine Ecology Progress Series* 95: 55-65
- Faasse MA, Bayha KM (2006) The ctenophore *Mnemiopsis leidy* A. Agassiz 1865 in coastal waters of the Netherlands: an unrecognized invasion? *Aquatic Invasions* 1: 270-277
- Hansson HG (2006) Ctenophores of the Baltic and adjacent Seas – the invader *Mnemiopsis* is here! *Aquatic Invasions* 1: 295-298
- Haslob H, Clemmesen C, Schaber M, Hinrichsen H-H, Schmidt J, Voss R, Kraus G, Köster FW (2007) Invading *Mnemiopsis leidy* as a potential threat to Baltic fish. *Marine Ecology Progress Series* 349: 303-306
- Huwer B, Storr-Paulsen M, Riisgård H-U, Haslob H (2008) Abundance, horizontal and vertical distribution of the invasive ctenophore *Mnemiopsis leidy* in the central Baltic Sea, November 2007. *Aquatic Invasions* 3: 113-124
- Janas U, Zgrundo A (2007) First record of *Mnemiopsis leidy* A. Agassiz, 1865 in the Gulf of Gdańsk (southern Baltic Sea). *Aquatic Invasions* 2: 450-454
- Javidpour J, Sommer U, Shiganova, T (2006) First record of *Mnemiopsis leidy* A. Agassiz 1865 in the Baltic Sea. *Aquatic Invasions* 1: 299-302
- Javidpour J, Molinero J C, Peschutter J, Sommer U, Shiganova T (2008) Seasonal changes and population dynamics of the ctenophore *Mnemiopsis leidy* after its first year of invasion in the Kiel Fjord, Western Baltic Sea. *Biological Invasions*. DOI 10.1007/s10530-008-9300-8
- Kremer P (1994) Patterns of abundance for *Mnemiopsis* in US coastal waters: A comparative overview. *ICES Journal of Marine Science* 51: 347-354
- Kube S, Postel L, Honnef C, Augustin C (2007) *Mnemiopsis leidy* in the Baltic Sea – distribution and overwintering between autumn 2006 and spring 2007. *Aquatic Invasions* 2: 137-145
- Köster FW, Möllmann C, Hinrichsen H-H, Wieland K, Tomkiewicz J, Kraus G, Voss R, Makarchouk A, MacKenzie BR, St. John MA, Schnack D, Rohlf N, Linkowski T, Beyer JE (2005) Baltic cod recruitment - the impact of climate variability on key processes. *ICES Journal of Marine Science* 62: 1408-1425

Mnemiopsis leidyi in the central Baltic

- Lehtiniemi M, Pääkkönen J-P, Flinkman J, Gorokhova E, Karjalainen M, Viitasalo S, Björk H (2007) Distribution and abundance of the American comb jelly (*Mnemiopsis leidyi*) – A rapid invasion to the northern Baltic Sea during 2007. *Aquatic Invasions* 2: 272-276
- Mutlu E (1999) Distribution and abundance of ctenophores and their zooplankton food in the Black Sea. II. *Mnemiopsis leidyi*. *Marine Biology* 135: 603-614
- Oliveira OMP (2007) The presence of the ctenophore *Mnemiopsis leidyi* in the Oslofjorden and considerations on the initial invasion pathways to the North and Baltic Seas. *Aquatic Invasions* 3: 185-189
- Postel L, Kube S, Hammer C, Zimmermann C, Sommer U, Javidpour J, Clemmesen C, Boersma M (2007) Die Invasion der räuberischen Rippenqualle *Mnemiopsis leidyi* in der Ostsee. Final Report, November 2007. IOW - Leibniz Institute for Baltic Sea Research, Warnemünde
- Purcell JE, Arai MN (2001) Interactions of pelagic cnidarians and ctenophores with fish: A review. *Hydrobiologia* 451: 27–44
- Purcell JE, Shiganova TA, Decker MB, Houde ED (2001) The ctenophore *Mnemiopsis* in native and exotic habitats: U.S. estuaries versus the Black Sea basin. *Hydrobiologia* 451: 145-176
- Purcell JE, Decker MB (2005) Effects of climate on relative predation by scyphomedusae and ctenophores on copepods in Chesapeake Bay during 1987–2000. *Limnology and Oceanography* 50: 376-387
- Reeve MR, MA Syms, Kremer P (1989) Growth dynamics of a ctenophore (*Mnemiopsis*) in relation to variable food supply. I. Carbon biomass, feeding, egg production, growth and assimilation efficiency. *Journal of Plankton Research* 11: 535-552
- Shiganova TA, Mirzoyan ZA, Studenikina EA, Volovik SP, Siokou-Frangou I, Zervoudaki S, Christou ED, Skirta AY, Dumont HJ (2001) Population development of the invader ctenophore *Mnemiopsis leidyi*, in the Black Sea and in other seas of the Mediterranean basin. *Marine Biology* 139: 431-445
- Tendal O, Jensen KR, Riisgård HU (2007) Invasive ctenophore *Mnemiopsis leidyi* widely distributed in Danish waters. *Aquatic Invasions* 2: 272-277
- Viitasalo S, Lehtiniemi M, Katajisto T (2008) The invasive ctenophore *Mnemiopsis leidyi* overwinters in high abundances in the subarctic Baltic Sea. *Journal of Plankton Research* 30: 1431-1436

Annex 1

Station number, position, sampling date, depth and abundance of *Mnemiopsis leidyi* caught with 335 µm and 500 µm Bongo nets, respectively. The station numbers refer to Figure 1

Station	Geographic coordinates		Date	Sampling depth (m)	Abundance	Abundance
	Latitude, °N	Longitude, °E			(ind. m ⁻²) 335 µm	(ind. m ⁻²) 500 µm
A1	54.817	13.083	16.03.2008	0- 35	0.00	0.00
A2	55.067	13.333	16.03.2008	0- 41	0.00	0.00
A3	54.983	13.667	16.03.2008	0- 45	0.00	0.00
A4	55.033	13.967	16.03.2008	0- 46	0.00	0.00
A5	55.183	14.133	16.03.2008	0- 45	0.00	0.00
A6	55.250	14.300	16.03.2008	0- 45	0.00	0.00
1	55.458	14.750	13.03.2008	0- 69	0.12	0.12
2	55.625	14.750	06.03.2008	0- 66	0.00	0.00
02a	55.792	14.750	06.03.2008	0- 47	0.00	0.00
3	55.625	15.000	13.03.2008	0- 75	0.33	0.11
03a	55.792	15.000	06.03.2008	0- 37	0.00	0.00
4	55.625	15.250	12.03.2008	0- 76	0.50	1.49
04a	55.792	15.250	06.03.2008	0- 51	0.00	0.00
5	55.625	15.500	12.03.2008	0- 67	0.13	0.13
05a	55.792	15.500	06.03.2008	0- 59	0.00	0.00
6	55.625	15.750	12.03.2008	0- 70	0.42	0.00
06a	55.792	15.750	06.03.2008	0- 42	0.00	0.00
7	55.625	16.000	08.03.2008	0- 73	0.83	0.69
8	55.792	16.000	07.03.2008	0- 61	0.00	0.00
9	55.792	16.250	07.03.2008	0- 60	0.00	0.00
10	55.625	16.250	08.03.2008	0- 72	0.43	0.57
11	55.792	16.500	07.03.2008	0- 56	0.00	0.00
11a	55.792	16.750	07.03.2008	0- 47	0.00	0.00
11b	55.792	17.000	07.03.2008	0- 37	0.00	0.00
12	55.625	16.500	07.03.2008	0- 62	0.00	0.00
12a	55.625	16.750	07.03.2008	0- 44	0.00	0.00
12b	55.625	17.000	07.03.2008	0- 26	0.00	0.00
13	55.458	16.500	05.03.2008	0- 59	0.00	0.00

Annex 1 (continued)

Station	Geographic coordinates		Date	Sampling depth (m)	Abundance	Abundance
	Latitude, °N	Longitude, °E			(ind. m ⁻²) 335 µm	(ind. m ⁻²) 500 µm
14	55.458	16.250	05.03.2008	0- 73	0.56	0.00
15	55.458	16.000	05.03.2008	0- 84	2.14	2.32
16	55.458	15.750	05.03.2008	0- 87	0.77	0.62
17	55.458	15.500	05.03.2008	0- 84	1.48	0.49
18	55.458	15.250	12.03.2008	0- 87	1.54	0.56
19	55.458	15.000	12.03.2008	0- 76	0.15	0.00
20	55.292	15.000	13.03.2008	0- 71	0.00	0.00
21	55.292	15.283	13.03.2008	0- 77	0.00	0.00
22	55.292	15.500	13.03.2008	0- 93	1.59	0.87
23	55.292	15.750	14.03.2008	0- 95	1.21	1.91
24	55.292	16.000	14.03.2008	0- 90	0.43	0.98
25	55.292	16.250	14.03.2008	0- 74	4.01	5.66
26	55.292	16.500	10.03.2008	0- 63	0.00	0.00
27	55.125	16.500	10.03.2008	0- 52	0.00	0.00
28	55.125	16.250	10.03.2008	0- 81	2.19	5.21
29	55.125	16.000	10.03.2008	0- 90	2.07	2.81
30	55.125	15.750	12.03.2008	0- 90	0.48	0.85
31	55.125	15.500	12.03.2008	0- 67	0.15	0.29
32	55.125	15.250	12.03.2008	0- 58	0.00	0.00
33	54.958	15.250	09.03.2008	0- 45	0.00	0.00
34	54.958	15.500	09.03.2008	0- 75	0.39	0.26
35	54.958	15.750	08.03.2008	0- 80	0.36	1.62
36	54.958	16.000	08.03.2008	0- 74	2.42	1.78
37	54.958	16.250	08.03.2008	0- 53	0.00	0.00
38	54.792	16.000	08.03.2008	0- 53	1.25	1.25
38a	54.792	16.250	08.03.2008	0- 50	0.00	0.20
38b	54.625	16.000	08.03.2008	0- 47	0.00	0.00
39	54.792	15.750	08.03.2008	0- 75	0.19	0.93
40	54.792	15.500	09.03.2008	0- 79	0.17	0.52
41	54.792	15.250	09.03.2008	0- 68	0.00	0.13
42	54.792	15.000	09.03.2008	0- 58	0.00	0.00
42a	54.792	14.750	09.03.2008	0- 46	0.00	0.00
43	54.625	15.250	10.03.2008	0- 58	0.16	0.00
43a	54.625	15.000	10.03.2008	0- 32	0.00	0.00
43b	54.625	14.750	09.03.2008	0- 47	0.00	0.00
44	54.625	15.500	09.03.2008	0- 65	1.44	2.02
45	54.625	15.750	09.03.2008	0- 60	0.50	0.84
46	55.230	16.800	04.03.2008	0- 75	0.34	0.34
47	55.250	17.080	04.03.2008	0- 80	0.00	0.36
48	55.233	17.367	04.03.2008	0- 66	0.00	0.33
49	55.250	17.583	05.03.2008	0- 63	0.00	0.00
50	55.233	17.917	05.03.2008	0- 58	0.00	0.00
51	55.483	17.917	05.03.2008	0- 66	0.00	0.00
52	55.417	17.583	04.03.2008	0- 76	0.00	0.00
53	55.350	17.367	04.03.2008	0- 70	0.00	0.00
54	55.350	17.083	04.03.2008	0- 66	0.00	0.00
55	55.350	16.800	04.03.2008	0- 63	0.00	0.00