

Technical University of Denmark



Engineering the spatial organization of metabolic pathways

Buron, Line Due; Maury, Jerome; Bach, Lars Stougaard; Nielsen, Jens; Mortensen, Uffe Hasbro

Publication date:
2011

[Link back to DTU Orbit](#)

Citation (APA):

Albertsen, L., Maury, J., Bach, L. S., Nielsen, J., & Mortensen, U. H. (2011). Engineering the spatial organization of metabolic pathways. Abstract from 6th Danish Conference on Biotechnology and Molecular Biology, Vejle, Denmark.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Poster abstract for DBC6: Cell Factories and synthetic biology, Munkebjerg Vejle. 26-27. Maj 2011.

Authors: Line Albertsen, Jerome Maury, Lars Bach, Jens Nielsen, Uffe Mortensen

Center for Microbial Biotechnology, DTU Systems Biology Building 227, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark

Engineering the spatial organization of metabolic pathways

One of the goals of metabolic engineering is to optimize the production of valuable metabolites in cell factories. In this context, modulating the gene expression and activity of enzymes are tools that have been extensively used. Another approach that is gaining interest is the engineering of the spatial organization of biosynthetic pathways. Several natural systems for ensuring optimal spatial arrangement of biosynthetic enzymes exist. Sequentially acting enzymes can for example be positioned in close proximity by attachment to cellular structures, up-concentration in membrane enclosed organelles or assembly into large complexes. The vision is that by positioning sequentially acting enzymes in close proximity, the cell can accelerate reaction rates and thereby prevent loss of intermediates through diffusion, degradation or competing pathways. The production of valuable metabolites in cell factories does however often depend on both heterologous and host enzymes. In this case, no spatial coordination of the biosynthetic enzymes can be expected to be in place. Presumably this contributes to the low productivity regularly observed for heterologous pathways. In one test case, we investigated whether a heterologous pathway could be optimized by positioning two sequentially acting enzymes in close proximity. More specifically, we fused a sesquiterpene synthase of plant origin to a natural yeast enzyme and expressed it in the well-characterised cell factory *Saccharomyces cerevisiae*. Successfully, the sesquiterpene production was increased two-fold when the enzymes were fused

compared to when they were expressed from the same promoters as free enzymes. Moreover, the strategy could be used in combination with other traditional metabolic engineering strategies to increase the production of a desired product, as enzyme fusion combined with down-regulation of a competing pathway and up-regulation of a selected pathway enzyme resulted in a five-fold higher sesquiterpene production. This simple test case demonstrates that engineering of the spatial organization of pathways has great potential for diverting flux towards a desired product.