

Technical University of Denmark



Fumonisin from *Aspergillus niger* in grapes and derived products

Mogensen, Jesper Mølgaard; Knudsen, Peter Boldsen; Larsen, Thomas Ostenfeld; Frisvad, Jens Christian; Thrane, Ulf; Nielsen, Kristian Fog

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Mogensen, J. M., Knudsen, P. B., Larsen, T. O., Frisvad, J. C., Thrane, U., & Nielsen, K. F. (2011). Fumonisin from *Aspergillus niger* in grapes and derived products. Poster session presented at The MycoRed Africa Conference, Cape Town, South Africa, .

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Fumonisin from *Aspergillus niger* in grapes and derived products

Jesper Mølgaard Mogensen, Peter Boldsen Knudsen, Thomas Ostenfeld Larsen, Jens Christian Frisvad, Ulf Thrane and Kristian Fog Nielsen

Introduction

Black *Aspergilli* are present on grape clusters from early in the season with increasing frequency during the growth stages of the grapes. Of the various *Aspergillus* species, *A. niger* is by far the most commonly found on grapes and are shown in one study to occur on more than 80% of samples of grapes and derived products. Although *A. niger* is the predominant species, *A. carbonarius* is the most problematic because it consistently produces high amounts of ochratoxin A while only 0-40% of *A. niger* strains produce this toxin, which is the main mycotoxin-related health concern in grape-derived products. The discovery of a fumonisin B₂, B₄ and B₆ production in *Aspergillus niger*, raises concerns about the presence of these mycotoxins in grapes as well as derived products.



Worst case

The potential fumonisin production by *A. niger* on grapes and raisins was determined by growth experiments on either commodity for 7 days at 25°C:

Grapes: The production of fumonisin B₂ on grapes varied almost 50 fold from 0.2 to 8 mg/kg while the B₄ varied from 0.01 to 1 mg/kg.

Raisins: Fumonisin B₂ and fumonisin B₄ in raisins with increasing water activity were produced in the range of 229-6476 and 27-356 µg/kg. Raisins with a decreasing water activity had a fumonisin B₂ concentration of 5-784 µg/kg and fumonisin B₄ of 12-672 µg/kg.

Cat-ion exchange purification and LC-MS/MS (2 transitions per compound)

Cat-ion exchange purification and LC-MS/MS (2 transitions per compound)

Analysis of wine

A total of 77 wine samples from 13 countries were analyzed. 18 wine samples (23%) were positive for fumonisin B₂ and contained 1-25 µg/L. Of the 18 positive samples, 16 were red wine, 1 was white wine, and 1 was port wine. As ochratoxin A FB₂ is more frequently found in red wine (28% positive) compared to white wine (7% positive). This was also confirmed by a later study (Logrieco et al 2010)

Analysis of retail raisins

A total of 21 raisin brands collected in Denmark, Germany and The Netherlands were analyzed. In 10 brands (48%), Fumonisin B₂ and B₄ were detected at 1.3-13 and 0.26-1.3 µg/kg, respectively. Large package variations were observed with up to 3-fold differences between four packages of the same brand, indicating a non-homogeneous infection level, which may be due to contaminated raisins.

Conclusion

- Fumonisin are frequently present in grape and derived products. This indicate that *A. niger* is apparently a commonly contaminant of grapes in the fields.
- Although frequently detected the amount of fumonisin is significantly below the regulatory limit set for similar food types (maize)
- The low levels found is presumably due to efficient removal of damaged grapes, initiated after problems with ochratoxin A in grapes and derived products were reported in the late 1990s. This lead to very strict regulations (EC472/2002), including a maximum allowance of 10 µg/kg ochratoxin A in dried vine fruits and 2 µg/kg in wine.

References

- J.M. Mogensen, T.O. Larsen, K.F. Nielsen. Widespread occurrence of the mycotoxin Fumonisin B₂ in wine. Journal of Agricultural and Food Chemistry. 2010, 58:4583-4587
 J.M. Mogensen, J.C. Frisvad, U. Thrane, K.F. Nielsen. Production of fumonisin B₂ and B₄ by *Aspergillus niger* on grapes and raisins. Journal of Agricultural and Food Chemistry. 2010, 58:954-958
 P. B. Knudsen, J. M. Mogensen, T. O. Larsen and K. F. Nielsen. Occurrence of fumonisins B₂ and B₄ in retail raisins. Journal of Agricultural and Food Chemistry. 2011, 59:772-776
 Logrieco, A.; Ferracane, R.; Visconti, A.; Ritieni, A. Natural occurrence of fumonisin B₂ in red wine from Italy. Food Additives and Contaminants Part A. 2010, 27, 1136-1141
 Logrieco, A.; Ferracane, R.; Haidukowsky, M.; Cozzi, G.; Visconti, A.; Ritieni, A. Fumonisin B₂ production by *Aspergillus niger* from grapes and natural occurrence in must. Food Additives and Contaminants Part A. 2009, 26, 1495-1500.