

Technical University of Denmark



## Vertical small scale variations of sorption and mineralization of three herbicides in subsurface limestone and sandy aquifer

Janniche, Gry Sander; Mouvet, C.; Albrechtsen, Hans-Jørgen

*Published in:*  
Journal of Contaminant Hydrology

*Link to article, DOI:*  
[10.1016/j.jconhyd.2011.01.002](https://doi.org/10.1016/j.jconhyd.2011.01.002)

*Publication date:*  
2011

[Link back to DTU Orbit](#)

*Citation (APA):*  
Janniche, G. S., Mouvet, C., & Albrechtsen, H-J. (2011). Vertical small scale variations of sorption and mineralization of three herbicides in subsurface limestone and sandy aquifer. *Journal of Contaminant Hydrology*, 123(3-4), 167-177. DOI: 10.1016/j.jconhyd.2011.01.002

**DTU Library**  
Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

**VERTICAL SMALL SCALE VARIATIONS OF SORPTION  
AND MINERALIZATION OF THREE HERBICIDES IN  
SUBSURFACE LIMESTONE AND SANDY AQUIFER**

G. S. Janniche<sup>a\*</sup>, C. Mouvet<sup>b</sup> and H.-J. Albrechtsen<sup>a</sup>

<sup>a</sup>DTU Environment, Department of Environmental Engineering, Technical University of Denmark, Miljoevej,

Building 113, DK-2800 Kgs. Lyngby, Denmark

<sup>b</sup> BRGM, 3 avenue Claude Guillemin, BP 6009, 45060 Orléans Cedex 2, France

\*Corresponding author

Telephone: (+45) 45 25 14 59

Fax: (+45) 45 93 28 50

Email: gsf@env.dtu.dk

## Abstract

Vertical variation in sorption and mineralization potential of mecoprop (MCP), isoproturon and acetochlor was investigated at low concentrations ( $\mu\text{g}$ -range) at the cm-scale in unsaturated sub-surface limestone samples and saturated sandy aquifer samples from an agricultural catchment in Brévilles, France. From two intact core drills, four heterogenic limestone sections were collected from 4.50-26.40 m below surface (mbs) and divided into 12 sub-samples of 8-25 cm length, and one sandy aquifer section from 19.20-19.53 m depth divided into 7 sub-samples of 4-5 cm length. In the sandy aquifer section acetochlor and isoproturon sorption increased substantially with depth; in average 78 % (acetochlor) and 61 % (isoproturon) per 5 cm. Also the number of acetochlor and isoproturon degraders (most-probable-number) was higher in the bottom half of the aquifer section ( $93 \rightarrow 16\,000$  /g) than in the upper half (4-71 /g). One 50 cm long limestone section with a distinct shift in color showed a clear shift in mineralization, number of degraders and sorption: In the two brown, uppermost samples, up to 31% mecoprop and up to 9% isoproturon was mineralized during 231 days, the numbers of mecoprop and isoproturon degraders were 1 300 to  $>16\,000$  /g, and the sorption of both isoproturon and acetochlor was more than three times higher, compared to the two deeper, grayish samples just below where mineralization ( $\leq 4$  %) and numbers of degraders (1-520 /g) were low for all three herbicides. In both unsaturated limestone and sandy aquifer, variations and even distinct shifts in both mineralization, number of specific degraders and sorption were seen within just 4-15 cm of vertical distance. A simple conceptual model of herbicides leaching to groundwater through a 10 m unsaturated limestone was established, and calculations showed that a 30 cm active layer with the measured sorption and mineralization values hardly impacted the fate of the investigated herbicides, whereas a total thickness of layers of 1 m would substantially increase natural attenuation.

50 **Keywords:** Isoproturon; mecoprop (MCP); acetochlor; degradation; pesticide; fate.

## 51 **1 Introduction**

52 Pesticides and their degradation products are often detected in surface water and  
53 groundwater (Battaglin et al., 2000; Kolpin et al., 1997; Thorling, 2009) even above the EU  
54 groundwater guideline of 0.1µg/L (European Parliament and Council, 2006) and they are in  
55 this way posing a substantial risk to drinking water resources. This diffuse pesticide pollution  
56 has become one of the major environmental challenges for using groundwater as drinking  
57 water, and to predict the fate of pollutants in groundwater the entire catchment has to be  
58 considered (Turner et al., 2006).

59 Knowledge on sorption and degradation is essential for evaluating risks of pesticides  
60 contamination of groundwater since these two processes are the major drivers in natural  
61 attenuation. Degradation of pesticides is often considered to decrease with depth (Fomsgaard,  
62 1995) but degradation can be faster in sub-soil than in top-soil (Mills et al., 2001). Sorption  
63 may also decrease with depth because of decreasing organic matter content which on the  
64 other hand may increase the bioavailability (Bending and Rodriguez-Cruz, 2007). Spatial  
65 variability in herbicide degradation and sorption has mainly been studied in topsoil or the  
66 upper meter of agricultural fields (e.g. Bending et al., 2001, 2003; Charnay et al., 2005;  
67 Jacques et al., 1999; Fredslund et al., 2008; Vinther et al., 2008; Walker et al., 2001, 2002)  
68 and even at the millimeter scale (Gonod et al., 2003).

69 Although several studies have focused on pesticide fate in subsurface materials and  
70 aquifers only few have studied the vertical variability in these environments (Albrechtsen et  
71 al., 2001; Charnay et al., 2005; Coquet et al., 2004; Johnson et al., 2000; Larsen et al., 2000;  
72 Wood et al., 2002). Even fewer have considered chalk or limestone, probably because of the  
73 complexity and the difficulties in sampling such geological settings. Vertical variation in  
74 limestone of atrazine mineralization was investigated in seven samples collected from 4.6-

75 10.0 mbs (Charnay et al, 2005), and small-scale vertical variation in isoproturon degradation  
76 was investigated in chalk (four 5-cm core sections from 4.42-4-88 mbs) (Johnson et al.,  
77 2000), but since degradation was negligible in both studies, their contribution to insight in the  
78 spatial variation was limited. In limestone (Charnay) the sorption of atrazine, isoproturon and  
79 metamitron varied vertically only to a limited degree in 10 out of 11 samples (3-12 mbs), but  
80 one sample had significantly higher  $K_d$ -values which could not be explained by the content of  
81 clay or organic matter (Coquet et al., 2004). Hence, only vertical variation in sorption but not  
82 in degradation of herbicides in subsurface carbonate rock was reported.

83         If sediment layers with increased mineralization could be identified it would be a step  
84 towards better understanding the interaction between sediment and biodegradation, which  
85 may improve the predictions and modeling of herbicides fate. Such sediment layers may in  
86 fact be “ecotones” where the microbial activity is high, found at the interface between  
87 different lithological layers (Goldscheider et al., 2006).

88         Spatial variability at the cm-scale in natural mineralization potential and sorption  
89 capacity of herbicides has to our knowledge not been studied before in deep subsurface  
90 samples (below 5 mbs), and especially not in limestone. The aim of this current work was to  
91 investigate i) if there is a vertical small scale spatial variability in sorption and in  
92 mineralization potential of the herbicides mecoprop, isoproturon and acetochlor in sediments  
93 from subsurface limestone and sandy aquifer; ii) if the variations could be related to a simple  
94 geochemical parameter; and iii) at which such a variation occurs and becomes important for  
95 the whole system.

## 96         **2 Materials and Methods**

### 97         **2.1 Sediment sampling and characterization**

98         Sediment was collected as intact cores in 2005 (Janniche et al., *submitted*), from a 4.5  
99 km<sup>2</sup> agricultural catchment with diffuse pesticide pollution. The unsaturated zone consists of

100 lutetian heterogeneous limestone, and the sandy aquifer is unconfined and aerobic with the  
101 groundwater table between 13.3 and 41.9 m below surface (mbs) (Morvan et al., 2006). Five  
102 sections were collected from two cores (Pz14 and Pz17c); four from the unsaturated  
103 limestone (sections A to D) and one from the saturated sand (section E). The sections were  
104 sampled from 4.50 to 26.40 mbs (Table 2). The sections were further divided vertically in 4  
105 to 25 cm long subsamples. The four limestone sections were selected based on a visual  
106 inspection: three with rather consistent color and texture over depth, and one section with a  
107 distinct shift in color that divided the section in a brownish upper part and a grayish lower  
108 part. The sand section had a color gradient over the section length; ranging from brown in  
109 the upper samples to dark grey especially in the two bottom samples.

110 Water content of the sediment samples was measured as weight loss after 24 h at 105  
111 °C. Sediment  $pH_w$  was measured by electrode in 10 g sediment mixed with 10 mL deionized  
112 (Bigham et al., 1996). Total inorganic carbon (TIC) was calculated as the difference between  
113 total carbon content (TC) and Total organic (TOC). TOC and TC were measured based on the  
114 method by Heron et al. (1997) by a total elemental carbon analyzer (LECO CS-225) in  
115 triplicates (0.2000 g) where carbonates were removed by adding 2 mL 6 % sulphurous acid  
116 ( $H_2SO_3$ ) six times to ensure that no bubble formation as released  $CO_2$  could be observed.  
117 Grain size (only for unconsolidated samples) was characterized by sieving (sieves of 0.063-  
118 2.0 mm) and, for the particles below 63  $\mu m$ , by particle distribution (Sedigraph 5100,  
119 Micromeritics Gemini). The specific surface area was measured by Multipoint  $N_2$ -BET  
120 analysis (Micromeritics, Gemini III 2375 surface area analyzer), after being outgassed  
121 (Micromeritics, FlowPrep 060 Degasser) for 4h at 70 °C.

## 122 **2.2 Herbicides**

123 Isoproturon and acetochlor are hydrophobic, non-polar organic herbicides whereas  
124 mecoprop is a hydrophilic phenoxy acid with a  $pK_a$  value of 3.78 and at near-neutral pH it is  
125 therefore negatively charged (Table 1). The three herbicides have all been used in the field

126 site to a wide extent. The experiments were carried out with [Ring-<sup>14</sup>C]-pesticides (Table 1).  
127 In the MPN also unlabelled acetochlor with 99.4 % purity was used.

128 **Table 1**

### 129 **2.3 Sorption investigations**

130 Sorption was investigated by a batch equilibrium technique based on the OECD  
131 guideline 106 (1993) at initial pesticide concentrations of 1 and 50 µg/L. Stock solutions of  
132 <sup>14</sup>C-labeled pesticide was prepared with <sup>14</sup>C-labelled pesticides in sterile filtered MilliQ water  
133 (Clausen et al., 2001). The sediment samples were crushed (only limestone), freeze-dried and  
134 sieved (<2 mm) (Madsen et al., 2000). The investigations were carried out in glass tubes (10  
135 mL) with Teflon caps at 10°C in the dark and previous tests excluded adsorption on those  
136 materials (Madsen et al., 2000). The water from the Brévilles Spring in the catchment was  
137 used as equilibrium liquid in the investigations as representative for the groundwater after  
138 treatment with activated carbon to remove pesticides and sterile filtering. Triplicate 5 ±  
139 0.0005 g of sediment was pre-equilibrated with 4 mL sterile filtered (0.2 µm Minisart,  
140 Sartorius) water for 20 h, then 1 mL pesticide solution was added resulting in an initial  
141 concentration of 1 or 50 µg/L (these low concentrations were chosen to mimic  
142 environmentally relevant concentrations), and each test tube was placed in a vertical rotator.  
143 Based on preliminary investigations (Mouvet et al., 2004) 168 h (for mecoprop and  
144 acetochlor) or 336 h (for isoproturon) was sufficient to reach equilibrium, and after this  
145 equilibrations time the suspension was centrifuged and 1 mL of supernatant collected for  
146 quantification of <sup>14</sup>C-activity (Clausen et al., 2001). Reference samples were prepared in  
147 triplicate without sediment but otherwise handled identically to the other samples. The  
148 amount of adsorbed pesticide was calculated as the difference between the <sup>14</sup>C-activity in the  
149 supernatant solution at equilibrium and the <sup>14</sup>C-activity in the reference solution (detection  
150 limit of 1 %). To check if the herbicides were degraded during the sorption equilibration 1  
151 mL of supernatant was transferred to a "double-vial" system; 20 mL polyethylene

152 scintillation vial containing a 6 mL scintillation vial with 1 mL 2.5 M NaOH to trap  $^{14}\text{CO}_2$   
153 after acidification of the subsample (0.1 mL 37%). The inner-vial was removed after 48  
154 hours, added scintillation cocktail and the  $^{14}\text{C}$ -activity in e.g. the  $^{14}\text{CO}_2$ -fraction was  
155 quantified.

## 156 **2.4 Mineralization**

157 Mineralization was investigated in laboratory batch incubations set up under unsaturated  
158 or saturated conditions depending on the origin of each sample (Janniche et al., *submitted*)  
159 and incubated aerobically in the dark at 10°C for up to 251 days. Autoclaved (20 min at 1.5  
160 bars and 120°C three times with one-day intervals) control incubations and duplicates were  
161 set up for selected sediments.

162 Unsaturated sediment was set up with 40 g (wet weight) sediment (limestone samples  
163 were “gently” crushed prior to a size that could pass the bottleneck with a diameter of 3 cm)  
164 in a 100 mL DURAN<sup>®</sup> bottle with Teflon inlayer caps and spiked with 0.5 mL of pesticide  
165 stock solution to reach initial concentrations of 1 µg/kg. The produced  $^{14}\text{CO}_2$  was trapped in a  
166 test tube placed inside each bottle with 2 mL 0.5 M NaOH. When sampled, the sodium  
167 hydroxide was replaced and mixed with 10 mL scintillation fluid and the  $^{14}\text{C}$ -activity  
168 quantified. The bottles were left open for 5 minutes in a laminar flow bench to replenish  
169 oxygen. Mineralization was calculated as the cumulated  $^{14}\text{CO}_2$  evolution over time as a  
170 percentage of initial  $^{14}\text{C}$ -pesticide e.g.  $^{14}\text{CO}_2/^{14}\text{C}_0$  (%).

171 Aquifer sediment was set-up with 40 g (wet weight) sediment and 60 mL groundwater  
172 in a 118 mL glass serum bottle, sealed with a 1 cm thick butyl rubber stopper and crimp caps.  
173 1 mL stock solution of  $^{14}\text{C}$  labeled pesticide was added through the stopper resulting in an  
174 initial concentration in the water phase of approximately 1 µg/L of each pesticide. As for the  
175 sorption investigations, the groundwater was collected from the Brévilles spring, treated with  
176 activated carbon and sterile filtered. When sampling, 2 mL filtered (0.2 µm PTFE-filter;  
177 Advantec/MFS 13HP) subsamples were acidified to strip off (from the water phase)  $^{14}\text{CO}_2$



178 that then was captured by NaOH in a "double-vial" system as described earlier. Pesticide  
179 content over time was calculated as the fraction of initial  $^{14}\text{C}$ -pesticide corrected for pesticide  
180 removed during sampling. Mineralization over time was calculated as  $^{14}\text{CO}_2$  evolution  
181 (corrected for loss to headspace) as a percentage of the corrected initial  $^{14}\text{C}$ -pesticide e.g.  
182  $^{14}\text{CO}_2/^{14}\text{C}_0$  (%).

## 183 **2.5 Enumeration of specific pesticide degraders**

184 The specific pesticide degraders was enumerated by a modified (Janniche et al.,  
185 *submitted*) most-probable-number method ( $^{14}\text{C}$ -MPN) (Lehmicke et al., 1979). For each  
186 pesticide, sediment samples were mixed with pyrophosphate-buffer (1.2 mM  $\text{Na}_4\text{P}_2\text{O}_7$ , pH =  
187 7.0) and 10-fold serial dilutions ( $10^0 - 10^{-4}$ ) were prepared from which five replicates were  
188 made, resulting in 25 MPN-tubes to achieve 1 enumeration. The dilutions were added a  
189 complex mineral medium (ISO 7827, 1984) with pH = 6.8 and 25  $\mu\text{g/L}$   $^{14}\text{C}$ -mecoprop or  $^{14}\text{C}$ -  
190 isoproturon, or 10  $\mu\text{g/L}$   $^{14}\text{C}$ -acetochlor and 20  $\mu\text{g/L}$  acetochlor (due to late receipt of the  
191 acetochlor isotope the MPN-assay was set up with unlabeled acetochlor, and 3 month later  
192 was isotope added). The MPN-tubes were incubated 11-23 months in the dark at 15 °C and  
193 the  $^{14}\text{CO}_2$  evolved during incubation was trapped and measured by liquid scintillation  
194 counting. Controls were set up with sterile pyrophosphate-buffer without sediment  
195 suspension.

196 The MPN values and 95 % confides intervals were calculated by the computer software  
197 MPN calculator (Curiale, 2000) based on the number of positive scored tubes in each of the  
198 five dilutions. A positive score means that the MPN-tube contained bacteria capable of  
199 mineralizing the specific herbicide. The MPN-tubes were scored positive if the trapped  $^{14}\text{CO}_2$   
200 was greater than the average of 10 sterile controls plus three times the standard deviation  
201 (Lehmicke et al. 1979). The MPNs containing mecoprop were scored as positive if the  
202 radioactivity in the basetrap was >2.7% of the initially added amount of labeled; the values  
203 for isoproturon was 1.2% and 2.0% for acetochlor.

## 204 **3 Results and Discussion**

### 205 **3.1 Sediment characterization**

206 Core sections A-D were taken from the heterogeneous unsaturated limestone and  
207 section E from the sandy aquifer (Table 2). The heterogeneous limestone samples had  
208 relatively low values of total inorganic carbon (TIC): 17-69 %  $\text{CaCO}_3$  with the lowest values  
209 found in the samples with marls. Though high TIC content may interfere with the total  
210 organic carbon (TOC) measuring, the results have high reproducibility, and are equivalent to  
211 reports for similar settings, e.g. 0.1-2.9 % TOC in limestone and 0-5.9 % in marlstone  
212 (Coquet et al., 2004), who also reported  $\text{CaCO}_3$  content of 69-88 % in limestone and 69-74 %  
213 in marlstone. The heterogeneous nature of our limestone may explain the relatively low TIC  
214 content compared to samples of pure limestone.

215 The limestone sections from Pz14 (A and C) had in general higher contents of water,  
216 TOC and TIC but a lower specific surface area than those from Pz17c (B and D). Besides  
217 differences observed between sections, differences were also observed at cm scale within a  
218 given section. Thus, limestone section D had a clear visual color shift and a clear shift in  
219 specific surface area with much higher specific surface area in the two uppermost samples  
220 (D1 and D2) compared to the two bottom samples (D3 and D4). Also the TIC content shifted  
221 with a lower TIC content in D1 and D2 compared to D3 and D4 expressing more pure  
222 limestone in D3-D4 than in D1-D2. Besides, TOC was much lower in the first sample, D1,  
223 than the other samples D2-D4 within this section.

224 Aquifer section (E) from Pz17c had high specific surface area but low TOC (Table 2).  
225 The color gradually changed across the whole section from brownish to grayish and TIC  
226 slightly increased with depth; samples E1 and E2 had approximately half the TIC content of  
227 E3-E7.

228 In summary, the five core sections represented two major geological settings and  
229 different limestone settings; the limestone cores varied in textures, but the geology of the

230 individual sections was very similar at the macro geological scale. Section D and E had clear  
231 small-scale variations at the micro scale in the sediment characterization parameters.

232 **Table 2**

233

### 234 **3.2 Sorption**

235 The sorption investigations included two concentrations (1 and 50  $\mu\text{g/L}$ ) and linear  
236 distribution coefficients ( $K_d$ ) were calculated. The different concentrations only to a limited  
237 degree affected the  $K_d$ -values (Fig. 1), and only noteworthy for isoproturon in the aquifer  
238 samples E3, E4 and E5, where sorption at the high concentration surprisingly was clearly  
239 higher (0.78-1.69 L/kg) than at the low concentration (0.52-0.87 L/kg). However, all  
240 isoproturon  $K_d$  values from section E followed the same trend of increased sorption over  
241 depth (Fig. 1).

242 **Fig. 1**

243 In limestone samples, sorption was in general low for all three herbicides and so was the  
244 standard deviation reflecting high reproducibility (Fig. 1 A-D). In all limestone sections,  
245 isoproturon sorption was at or below the detection limit of 0.010 L/kg except in section D  
246 ( $K_d$ : 0.03-0.25 L/kg). These findings are within the range reported for isoproturon sorption to  
247 limestone and chalk: <0.01-0.55 L/kg (Besien et al., 2000; Coquet et al., 2004; Johnson et al.,  
248 1998; Madsen et al., 2000; Mouvet et al., 2004). The  $K_d$  values for acetochlor in the  
249 limestone was <0.01-0.34 L/kg; literature  $K_d$  values for acetochlor, in the range of 0.48-7.11  
250 L/kg (Hiller et al., 2009; Lengyel and Foldenyi, 2003; Ma et al., 2004), are predominantly  
251 reported for topsoil which explains their substantially higher  $K_d$  values. The mecoprop  
252 sorption to limestone was also low ( $\leq 0.15$  L/kg) and in the range (0-0.17 L/kg) reported in  
253 the literature for chalk and limestone (Clausen et al., 2001; Kristensen et al., 2001; Madsen et  
254 al., 2000).

255           The sorption of isoproturon and acetochlor to limestone was highest in D1 and D2  
256 with higher specific surface area, and lower TOC than in the other limestone samples (Table  
257 2). This indicated that isoproturon and acetochlor sorption in heterogenic limestone to a  
258 higher degree was controlled by mineralogy than by organic carbon content. Isoproturon  
259 sorbs to kaolinite, but not to quartz, calcite, alumina (Clausen et al., 2001) or iron oxides  
260 (Clausen and Fabricius, 2001), and then sorption is strongly related to the content of organic  
261 matter and clay (Coquet et al., 2004; Madsen et al., 2000). This supports that the increased  
262 isoproturon sorption was caused by the increased clay content within the limestone marls.  
263 Sorption of acetochlor is likely to be controlled by the same parameters, but comparable  
264 literature data are limited for acetochlor sorption in subsurface sediments, low in TOC.  
265 Sorption of mecoprop was not increased in the same sections as isoproturon and acetochlor  
266 because the sorption mechanisms of the anionic mecoprop differ from the nonionic herbicides  
267 isoproturon and acetochlor; mecoprop sorbs to positive sites on the mineral surface (Clausen  
268 et al., 2001).

269           In sandy aquifer samples sorption was in general substantial for acetochlor (0.25-5.12  
270 L/kg) and isoproturon (<0.01-1.69 L/kg) except in E1 (Fig. 1), whereas mecoprop sorption  
271 was very low ( $\leq 0.12$  L/kg) in the two investigated samples (E2 and E3). The sorption could  
272 not be explained solely by variations in the investigated geochemical parameters (clay  
273 content, organic and inorganic carbon content). The grayish color observed in the bottom two  
274 subsamples (E6 and E7) of the aquifer section indicated reduced conditions which may,  
275 however, explain the increased sorption capacity over depth because reduced organic matter  
276 induce a more hydrophobic surface whereto nonionic compounds better sorbs (Clausen et al.,  
277 2004).  $K_d$  values of 0-0.4 L/kg have been reported for mecoprop sorption onto sand (Madsen  
278 et al., 2000; Pedersen, 2000; Tuxen et al., 2000), and our finding is within this range.  $K_d$   
279 values for isoproturon are reported to be 0.007-1 L/kg in sand (Coquet et al., 2004; Madsen et

280 al., 2000; Pedersen, 2000; Tuxen et al., 2000); our results are in close agreement with these  
281 values.

### 282 **3.3 Mineralization potential**

283 Mecoprop, isoproturon and acetochlor degraders were detected in many samples (Fig. 2)  
284 which indicated a degradation potential of these compounds in the catchment. Furthermore,  
285 this suggested that the catchment was polluted with all three herbicides since another study  
286 only detected phenoxy acid (mecoprop, dichlorprop and 2,4-D) degraders in the part of the  
287 aquifer exposed to phenoxy acids (<40 µg/L) but not in the upstream pristine aquifer (de  
288 Liphay et al., 2003). The number of mecoprop degraders found in our investigation was at  
289 same level as observed by de Liphay et al. (2003) but the number of isoproturon degraders  
290 was higher than reported by Bending & Rodriguez-Cruz (2007).

#### 291 **Fig. 2**

292 In all the 12 limestone samples mineralization of acetochlor was below 2%. Isoproturon  
293 was mineralized (up to 9%) in D1 and D2; in the remaining samples <1% was mineralized.  
294 19-31 % mecoprop was mineralized in D1 and D2 and 4 % in D3 and C4 (average of two  
295 replicates with 6 and 2 %). The other samples exhibited mecoprop mineralization below 1 %.  
296 The number of degraders was in general high in section D, but also in sample C3 the number  
297 of isoproturon degraders was extremely high (>16 000 /g). Some samples had high number of  
298 degraders without subsequent high mineralization (e.g. D1 (acetochlor), C3 (isoproturon)).

299 In the sandy aquifer (E), mecoprop was also mineralized substantially; 10-22% was  
300 recovered as <sup>14</sup>CO<sub>2</sub> with only 49-790 /g mecoprop degraders. The mineralization of both  
301 isoproturon (1-5%) and acetochlor (3-6%) was markedly lower than for mecoprop, despite  
302 higher numbers of isoproturon (27-2 200 /g) and acetochlor degraders (4-16 000 /g).

### 303 **3.4 Small scale variability**

304 The 19 samples from the 5 sections fell into 3 different cases: 1) In unsaturated  
305 limestone sections A, B and C sorption, mineralization and MPN were in general limited for

306 all three studied herbicides (one exception was mecoprop sorption which was  $\leq 0.01$  L/kg in  
307 sample A1 but 10-fold higher (0.11-0.15 L/kg) in A2 – the sample just below ). 2)  
308 Unsaturated limestone section D clearly showed small scale variations with a clear shift in  
309 sorption, mineralization and MPN. 3) Sandy aquifer section E had evident small scale  
310 variation in sorption but without a clear shift; it was rather a gradient, but also with variation  
311 in especially MPN.

### 312 **3.4.1 Unsaturated limestone from section D**

313 Sorption of isoproturon and acetochlor, and mineralization of mecoprop and acetochlor  
314 as well as number of mecoprop and acetochlor degraders were increased in the upper part of  
315 section D (Fig. 1 and 2). A shift in color divided the section in two: the two brownish  
316 uppermost samples (D1 and D2) with high sorption (isoproturon: 0.17-0.25 L/kg and  
317 acetochlor: 0.22-0.34 L/kg), mineralization (mecoprop: 19-31 % and isoproturon: 9 %), and  
318 number of herbicide degraders (mecoprop: 1300-1400 /g and isoproturon: 4300->16000 /g)  
319 and on the other hand the grayish, lower samples (D3 and D4) with lower sorption  
320 (isoproturon: 0.03-0.07 L/kg and acetochlor: <0.01-0.11 L/kg), mineralization (mecoprop:  
321 0.5-4 % and isoproturon: 0.2-0.5 %), and number of herbicide degraders (mecoprop: 1-6 /g  
322 and isoproturon: 3-520 /g). Already after 14 days of incubation the mineralization in D1 and  
323 D2 was noticeable higher than in D3 and D4 (Fig. 3). Furthermore, the  $^{14}\text{CO}_2$  evolution in D4  
324 was as low as in the autoclaved incubations with sediment from D1-D4. Hence, a clear small  
325 scale shift in sorption and mineralization potential within only 25 cm was evident in the  
326 limestone.

327 **Fig. 3**

### 328 **3.4.2 Sandy aquifer section E**

329 In the sandy aquifer samples sorption was low in the two uppermost samples, and  
330 increased gradually with depth. Acetochlor sorption increased in average 78 % pr every 5 cm

331 in the sandy aquifer section (E). Also the isoproturon sorption increased gradually with depth  
332 (in average 61 % pr 5 cm) in the sandy aquifer (E) and hence the small scale variance in  
333 acetochlor and isoproturon sorption within these 33 cm was substantial. Despite the  
334 acetochlor mineralization in the sandy aquifer was rather constant, the number of acetochlor  
335 degraders increased dramatically in the three bottom samples; from 4-9/g in E1-E4 to 120-16  
336 000/g in E5-E7. Also the number of isoproturon degraders was higher in the bottom samples  
337 (93-2 200/g in E4-E7) than in the top samples (27-71/g in E1-E3), and mineralization was  
338 faster in E1-E4 (2-5 %) than in E5-E7 (<1 %). For mecoprop however, the middle part (E3-  
339 E5) had the highest number of degraders; 490-790/g compared to 49-93/g, and mineralization  
340 was also substantially faster in E5 (22 %) compared to the rest of the aquifer (10-15 %).  
341 Mineralization of acetochlor and isoproturon in the saturated sandy aquifer samples (Fig. 2)  
342 could very well be negatively affected by the increased sorption with depth, leaving less  
343 herbicide available for mineralization, and could explain the lack of faster mineralization,  
344 despite very high number of degraders in some samples. This inverse relationship between  
345 sorption strength and mineralization was reported for MCPA (Jensen et al., 2004). This could  
346 explain why isoproturon was mineralized (1.7-4.7 %) in the uppermost samples (E2-E4) but  
347 not in the bottom samples (<1 % in E5-E7). However, acetochlor which sorbed stronger did  
348 not follow that trend.

349 Conclusively, sorption and mineralization potential varied substantially within just a  
350 few cm in both limestone and sand. These small scale variations were found in sections with  
351 shifts in color, but could not be strongly related to the investigated geochemical parameters.

### 352 **3.5 Importance of an active limestone layer on groundwater** 353 **contamination**

354 The investigated four limestone sections revealed an overall picture of very low  
355 herbicide sorption and mineralization, except in the upper two 15 cm samples of section D,  
356 where both sorption and mineralization potential was increased. This section contained marls

357 (Table 2) and upper Lutetian marly layers are ubiquitous in the catchment, and may reduce  
358 the flow rate and subsequent solute transport (Gutierrez and Baran, 2009). This marly layer  
359 with increased water content may be an environment favoring microbial activity.

360 To evaluate the relative impact on mecoprop, isoproturon and acetochlor of such an  
361 active layer with respect to sorption and mineralization in the unsaturated limestone a simple  
362 conceptual model was set up. Although both matrix flow and fracture flow may occur in  
363 heterogeneous unsaturated limestone matrix flow has in several studies shown to be  
364 predominant in deep groundwater sites and at steady-state flow (Haria et al., 2003; Mathias et  
365 al., 2005; Wellings, 1984). Slow transfer velocities of 0.5 to 1 m/yr have earlier been  
366 determined for our catchment based on profiles of water contents, tritium activities, and  
367 nitrate concentrations (Gutierrez and Baran, 2009). We therefore assumed that the dominant  
368 flow occurred in matrix in the calculations of the conceptual model consisting of a 10 m  
369 unsaturated heterogeneous limestone profile with an active layer of a certain thickness ( $x_a$ ).  
370 Sorption and mineralization was confined to this active layer and hence microbial activity is  
371 assumed in the matrix. Such an active layer with sorption will cause retardation ( $R$ ) of the  
372 herbicides which was calculated assuming matrix flow as:

$$373 \quad R = \frac{x_w}{x_s} = \frac{v_w}{v_s} = \frac{\rho_b}{\varepsilon} K_d + 1$$

374 where  $v_s$  is the solute velocity [m/y],  $\rho_b$  is the bulk density [kg/L] calculated as  $\rho_b = (1 - \varepsilon)\rho$ ,  $\varepsilon$   
375 is the porosity [-], and  $K_d$  is the linear sorption coefficient [L/kg]. The delay in break through  
376 due to sorption of the herbicides in the active layer was calculated as:

$$377 \quad Delay = \frac{x_a}{v_w} (R - 1)$$

378 Mineralization of herbicide will reduce the risk of groundwater contamination in  
379 accordance with the observed mineralization patterns (Fig. 3) and the removal of herbicides  
380 due to mineralization was calculated based on first order mineralization kinetics as:



381 
$$removal = \left( 1 - \exp\left( k \frac{x_a R}{v_w} \right) \right) \times 100\%$$

382 where k is the first order mineralization constant.  $K_d$ -values and k-values are mean values as  
383 observed for D1 and D2 (Fig. 1 and 2). Unfortunately, density ( $\rho$ ) and porosity ( $\epsilon$ ) were not  
384 measured specifically for D1 and D2 due to lack of sample material but in 7 other samples  
385 from the catchment;  $\rho$  ranged from 2.69 to 2.73 kg/L and  $\epsilon$  from 0.08 to 0.37 (Janniche et al.,  
386 *submitted*). We have used the mean value of 0.18 for  $\epsilon$  and 2.72 kg/L for  $\rho$  in this simple  
387 conceptual model calculation which results in a  $\rho_b$  of 2.2 kg/L. If a lower porosity ( $\epsilon$ ) and  
388 hence a higher bulk density ( $\rho_b$ ) was applied the retardation would have increased, and vice  
389 versa.

390 Three scenarios were set up: 1) a reference scenario with no active layer, 2) an active  
391 layer of 30 cm thickness, and 3) an active layer of 1 m thickness (Fig. 4). In the reference  
392 scenario the herbicides are transported at the same velocity as water ( $v_w$ ), and with a water  
393 velocity of 1 m/yr it will take 10 years before breakthrough of the herbicides through the 10  
394 m long profile. Presence of a 30 cm thick layer (scenario 2) will only to a very limited degree  
395 cause delay. It would take only 10.2 year for mecoprop to reach the groundwater (bottom of  
396 profile), 10.7 year for isoproturon and 11.0 year for acetochlor (Fig. 4). However, the  
397 mineralization in the layer would remove approximately 24 % mecoprop, 10 % isoproturon  
398 and 3 % acetochlor. The heterogeneous limestone could contain several active layers or one  
399 thicker layer. This may be realistic since the limestone settings in the field site are more than  
400 30 m thick in 61 % and less than 10 m thick in 7 % of the surface of the catchment (Morvan  
401 et al., 2006). Assuming the summed extent of active layers to be 1 m over the whole 10 m  
402 profile (scenario 3), then the breakthrough times would be 10.6 years (mecoprop), 12.4 years  
403 (isoproturon) and 13.3 years (acetochlor) respectively, and the proportion of mineralized  
404 herbicide would then become 59 % mecoprop, 31 % isoproturon and only 9 % acetochlor  
405 (Fig. 4).

406 **Fig. 4**

407 In conclusion, if only one active layer of 30 cm thickness is present pr 10 m the  
408 retardation due to sorption will be negligible for all three herbicides and the removal will  
409 negligible for acetochlor but not for isoproturon and mecoprop, whereas the effect of both  
410 retardation and removal will be notable for all three herbicides if the layer is 3 times thicker.  
411 Considering the thickness of unsaturated limestone in the investigated field site is up to 42 m  
412 thick it is not unlikely that the summed extend of an active layer will be greater than the  
413 observed 30 cm. Hence the importance of such active layers in the unsaturated limestone  
414 depends on their magnitude and consequently a very detailed knowledge of the limestone  
415 characteristics of the site is paramount.

## 416 **4 Conclusions**

417 Within few centimeters small scale vertical variability was present beyond measuring  
418 and sampling uncertainties showing clear shifts in sorption and microbial activity in both  
419 heterogeneous unsaturated limestone and sandy aquifer. None of the investigated parameters  
420 (TOC; TIC; specific surface area; sand, silt and clay content; water content;  $\text{pH}_w$ )  
421 individually could explain the variations. These variations followed a color shift or a color  
422 gradient in the core section.

423 The significance of these layers, or ecotones, with increased sorption and microbial  
424 mineralization depends on the specific herbicide and the extent of these layers, and it is  
425 therefore important to determine this. Such a detailed knowledge will improve the predictions  
426 of herbicide fate in the catchment if the geology is well described and included in reactive  
427 geochemical models.

## 428 **Acknowledgement**

429 The European Union FP6 Integrated Project AquaTerra (Project no. GOCE 505428)  
430 under the thematic priority: sustainable development, global change and ecosystems funded

431 this study together with the Technical University of Denmark (DTU). <sup>14</sup>C-labelled acetochlor  
432 was kindly provided by Dow Agroscience. The authors thank Ellinor Lindberg and Mona  
433 Refstrup for laboratory assistance, and Torben Dolin for the graphic assistance. Also thank to  
434 Bolette Nygaard and BRGM who took active part during drilling at Brévilles, and Alexis  
435 Gutierrez and Nicole Baran for always being helpful.

## 436 **References**

- 437 Albrechtsen, H.-J., Mills, M.S., Aamand, J., Bjerg, P.L., 2001. Degradation of herbicides in  
438 shallow Danish aquifers: an integrated laboratory and field study. *Pest Manag. Sci.*  
439 *57*, 341-350.
- 440 Battaglin, W.A., Furlong, E.T., Burkhardt, M.R., Peter, C.J., 2000. Occurrence of  
441 sulfonyleurea, sulfonamide, imidazolinone, and other herbicides in rivers, reservoirs  
442 and ground water in the Midwestern United States, 1998. *Sci. Total Environ.* *248*,  
443 123-133.
- 444 Bending, G.D., Lincoln, S.D., Sorensen, S.R., Morgan, J.A.W., Aamand, J., Walker, A., 2003.  
445 In-field spatial variability in the degradation of the phenyl-urea herbicide isoproturon  
446 is the result of interactions between degradative *Sphingomonas* spp. and soil pH.  
447 *Appl. Environ. Microb.* *69*, 827-834.
- 448 Bending, G.D., Rodriguez-Cruz, M.S., 2007. Microbial aspects of the interaction between  
449 soil depth and biodegradation of the herbicide isoproturon. *Chemosphere.* *66*, 664-  
450 671.
- 451 Bending, G.D., Shaw, E., Walker, A., 2001. Spatial heterogeneity in the metabolism and  
452 dynamics of isoproturon degrading microbial communities in soil. *Biol. Fert. Soils.*  
453 *33*, 484-489.
- 454 Besien, T.J., Williams, R.J., Johnson, A.C., 2000. The transport and behaviour of isoproturon  
455 in unsaturated chalk cores. *J. Contam. Hydrol.* *43*, 91-110.
- 456 Bigham, J.M. (Editor-in-Chief) et al., 1996. *Methods of Soil Analysis. Part 3 Chemical*  
457 *Methods.* Soil Science Society of America, Inc. and American Society of Agronomy,  
458 Inc., Madison, Wisconsin, USA.
- 459 Charnay, M.P., Tuis, S., Coquet, Y., Barriuso, E., 2005. Spatial variability in C-14-herbicide  
460 degradation in surface and subsurface soils. *Pest Manag. Sci.* *61*, 845-855.
- 461 Clausen, L., Fabricius, I., 2001. Atrazine, isoproturon, mecoprop, 2,4-D, and bentazone  
462 adsorption onto iron oxides. *J. Environ. Qual.* *30*, 858-869.
- 463 Clausen, L., Fabricius, I., Madsen, L., 2001. Adsorption of pesticides onto quartz, calcite,  
464 kaolinite, and alpha-alumina. *J. Environ. Qual.* *30*, 846-857.

- 465 Clausen, L., Larsen, F., Albrechtsen, H.-J., 2004. Sorption of the herbicide dichlobenil and  
466 the metabolite 2,6-dichlorobenzamide on soils and aquifer sediments. *Environ. Sci.*  
467 *Technol.* 38, 4510-4518.
- 468 Coquet, Y., Ribiere, C., Vachier, P., 2004. Pesticide adsorption in the vadose zone: a case  
469 study on Eocene and Quaternary materials in Northern France. *Pest Manag. Sci.* 60,  
470 992-1000.
- 471 Curiale, 2000. MPN Calculator [Build 24], Software program.  
472 <http://members.ync.net/mcuriale/mpn/index.html>.
- 473 de Liphthay, J.R., Tuxen, N., Johnsen, K., Hansen, L.H., Albrechtsen, H.-J., Bjerg, P.L.,  
474 Aamand, J., 2003. In situ exposure to low herbicide concentrations affects microbial  
475 population composition and catabolic gene frequency in an aerobic shallow aquifer.  
476 *Appl. Environ. Microb.* 69, 461-467.
- 477 European Parliament and Council, 2006. Directive 2006/118/EC of the European Parliament  
478 and of the Council of 12 December 2006 on the protection of groundwater against  
479 pollution and deterioration.
- 480 Fomsgaard, I.S., 1995. Degradation of Pesticides in Subsurface Soils, Unsaturated Zone - A  
481 Review of Methods and Results. *Int. J. Environ. An. Ch.* 58, 231-245.
- 482 Fredslund, L., Vinther, F.P., Brinch, U.C., Elsgaard, L., Rosenberg, P., Jacobsen, C.S., 2008.  
483 Spatial variation in 2-methyl-4-chlorophenoxyacetic acid mineralization and sorption  
484 in a sandy soil at field. *J. Environ. Qual.* 37, 1918-1928.
- 485 Goldscheider, N., Hunkeler, D., Rossi, P., 2006. Review: Microbial biocenoses in pristine  
486 aquifers and an assessment of investigative methods. *Hydrogeol. J.* 14, 926-941.
- 487 Gonod, L.V., Chenu, C., Soulas, G., 2003. Spatial variability of 2,4-dichlorophenoxyacetic  
488 acid (2,4-D) mineralisation potential at a millimetre scale in soil. *Soil Biol. Biochem.*  
489 35, 373-382.
- 490 Gutierrez, A., Baran, N., 2009. Long-term transfer of diffuse pollution at catchment scale:  
491 Respective roles of soil, and the unsaturated and saturated zones (Brevilles, France). *J.*  
492 *Hydrol.* 369, 381-391.
- 493 Haria, A.H., Hodnett, M.G., Johnson, A.C., 2003. Mechanisms of groundwater recharge and  
494 pesticide penetration to a chalk aquifer in southern England. *J Hydrol*, 275, 122-137.
- 495 Heron, G., Barcelona, M.J., Andersen, M.L., Christensen, T.H., 1997. Determination of  
496 nonvolatile organic carbon in aquifer solids after carbonate removal by sulfurous acid.  
497 *Ground Water* 35, 6-11.
- 498 Hiller, E., Cernansky, S., Krascenits, Z., Milicka, J., 2009. Effect of soil and sediment  
499 composition on acetochlor sorption and desorption. *Environ. Sci. Pollut. R.* 16, 546-  
500 554.
- 501 ISO 7827, 1984. Water quality-evaluation in an aqueous medium of the "ultimate" aerobic  
502 biodegradability of organic compounds-methods by analysis of dissolved organic  
503 carbon (DOC). International Organization for Standardization.

- 504 Jacques, D., Mouvet C., Mohanty, B., Vereecken, H., Feyen, J., 1999. Spatial variability of  
505 atrazine sorption parameters and other soil properties in a podzoluvisol. *J. Contam.*  
506 *Hydrol.* 36, 31-52.
- 507 Janniche, G., Lindberg, E., Mouvet, C., Albrechtsen, H.-J. Distribution of isoproturon,  
508 mecoprop and acetochlor mineralization and mineralizers in deep unsaturated  
509 limestone and sandy aquifer. *Chemosphere submitted*.
- 510 Jensen, P.H., Hansen, H.C.B., Rasmussen, J., Jacobsen, O.S., 2004. Sorption-controlled  
511 degradation kinetics of MCPA in soil. *Environ. Sci. Technol.* 38, 6662-6668.
- 512 Johnson, A.C., Hughes, C.D., Williams, R.J., Chilton, P.J., 1998. Potential for aerobic  
513 isoproturon biodegradation and sorption in the unsaturated and saturated zones of a  
514 chalk aquifer. *J. Contam. Hydrol.* 30, 281-297.
- 515 Johnson, A.C., White, C., Bhardwaj, C.L., 2000. Potential for isoproturon, atrazine and  
516 mecoprop to be degraded within a chalk aquifer system. *J. Contam. Hydrol.* 44, 1-18.
- 517 Kolpin, D.W., Kalkhoff, S.J., Goolsby, D.A., Sneek-Fahrer, D.A., Thurman, E.M., 1997.  
518 Occurrence of selected herbicides and herbicide degradation products in Iowa's  
519 ground water, 1995. *Ground Water* 35, 679-688.
- 520 Kristensen, G.B., Johannesen, H., Aamand, J., 2001. Mineralization of aged atrazine and  
521 mecoprop in soil and aquifer chalk. *Chemosphere* 45, 927-934.
- 522 Larsen, L., Sorensen, S.R., Aamand, J., 2000. Mecoprop, isoproturon, and atrazine in and  
523 above a sandy aquifer: Vertical distribution of mineralization potential. *Environ. Sci.*  
524 *Technol.* 34, 2426-2430.
- 525 Lehmicke, L.G., Williams, R.T., Crawford, R.L., 1979. C-14-Most-Probable-Number  
526 Method for Enumeration of Active Heterotrophic Microorganisms in Natural-Waters.  
527 *Appl. Environ. Microb.* 38, 644-649.
- 528 Lengyel, Z. Foldenyi, R., 2003. Acetochlor as a soil pollutant. *Environ. Sci. Pollut. R.* 10, 13-  
529 18.
- 530 Ma, Q.L., Rahman, A., Holland, P.T., James, T.K., McNaughton, D.E., 2004. Field  
531 dissipation of acetochlor in two New Zealand soils at two application rates. *J.*  
532 *Environ. Qual.* 33, 930-938.
- 533 Madsen, L., Lindhardt, B., Rosenberg, P., Clausen, L., Fabricius, I., 2000. Pesticide sorption  
534 by low organic carbon sediments: a screening for seven herbicides. *J. Environ. Qual.*  
535 29, 1488-1500.
- 536 Mathias, S.A., Butler, A.P., McIntyre, N., Wheeler, H.S., 2005. The significance of flow in  
537 the matrix of the Chalk unsaturated zone. *J Hydrol*, 310, 62-77.
- 538 Mills, M.S., Hill, I.R., Newcombe, A.C., Simmons, N.D., Vaughan, P.C., Verity, A.A., 2001.  
539 Quantification of acetochlor degradation in the unsaturated zone using two novel in  
540 situ field techniques: comparisons with laboratory-generated data and implications for  
541 groundwater risk assessments. *Pest Manag. Sci.* 57, 351-359.

- 542 Morvan, X., Mouvet, C., Baran, N., Gutierrez, A., 2006. Pesticides in the groundwater of a  
543 spring draining a sandy aquifer: Temporal variability of concentrations and fluxes. *J.*  
544 *Contam. Hydrol.* 87, 176-190.
- 545 Mouvet, C., Albrechtsen, H.-J., Baran, N., Chen, T., Clausen, L., Darsy, C., Desbionne, S.,  
546 Douguet, J.M., Dubus, I.G., Fialkiewicz, W., Gutierrez, A., Haverkamp, R., Herbst,  
547 M., Howles, D., Jarvis, N.J., Jørgensen, P.R., Larsbo, M., Meiwirth, K., Mermoud, A.,  
548 Morvan, X., Normand, B., O'Connor, M., Ritsema, C., Roessle, S., Roulier, S.,  
549 Soutter, M., Stenemo, F., Thiery, D., Vachaud, G., Vereecken, H., Vischetti, C., 2004.  
550 PEGASE. Pesticides in European Groundwaters: detailed study of representative  
551 quifers and Simulation of possible Evolution scenarios. Dubus I.G. and C. Mouvet  
552 (Editors). Final report of the European project #EVK1-CT1990-00028. BRGM/RP-  
553 52897-FR, 358 p.
- 554 OECD, 1993. OECD guidelines for testing of chemicals.106 Adsorption/desorption Paris,  
555 France.
- 556 Pedersen, P. G., 2000. Pesticide degradability in groundwater: Importance of redox  
557 conditions. Ph.D.Thesis, 1-67. Department of Environmental Science and  
558 Engineering, Technical University of Denmark, Kgs. Lyngby, Denmark.
- 559 Thorling, L. (ed.), 2009. Grundvandsovervågning 2008 (Groundwater surveillance 2008).  
560 GEUS, Denmark. [in Danish]
- 561 Tomlin, C.D.S. (editor), 1997. The Pesticide Manual, Eleventh ed. British Crop Protection  
562 Council, UK.
- 563 Turner, J., Albrechtsen, H.-J., Bonell, M., Duguet, J.P., Harris, B., Meckenstock, R.,  
564 McGuire, K., Moussa, R., Peters, N., Richnow, H.H., Sherwood-Lollar, B.,  
565 Uhlenbrook, S., van Lanen, H., 2006. Future trends in transport and fate of diffuse  
566 contaminants in catchments, with special emphasis on stable isotope applications.  
567 *Hydrol. Process.* 20, 205-213.
- 568 Tuxen, N., Tuxsen, P.L., Ruge, K., Albrechtsen, H.-J., Bjerg, P.L., 2000. Fate of seven  
569 pesticides in an aerobic aquifer studied in column experiments. *Chemosphere* 41,  
570 1485-1494.
- 571 Vinther, F.P., Brinch, U.C., Elsgaard, L., Fredslund, L., Iversen, B.V., Torp, S., Jacobsen,  
572 C.S., 2008. Field-scale variation in microbial activity and soil properties in relation to  
573 mineralization and sorption of pesticides in a sandy soil. *J. Environ. Qual.* 37, 1710-  
574 1718.
- 575 Walker, A., Bromilow, R.H., Nicholls, P.H., Evans, A.A., Smith, V.J.R., 2002. Spatial  
576 variability in the degradation rates of isoproturon and chlorotoluron in a clay soil.  
577 *Weed Res.* 42, 39-44.
- 578 Walker, A., Jurado-Exposito, M., Bending, G.D., Smith, V.J.R., 2001. Spatial variability in  
579 the degradation rate of isoproturon in soil. *Environ. Pollut.* 111, 407-415.
- 580 Wellings, S.R., 1984. Recharge of the Upper Chalk Aquifer at A Site in Hampshire, England  
581 1. Water-Balance and Unsaturated Flow. *J Hydrol*, 69, 259-273.  
582

583 Wood, M., Issa, S., Albuquerque, M., Johnson, A.C., 2002. Spatial variability in herbicide  
584 degradation in the subsurface environment of a groundwater protection zone. *Pest*  
585 *Manag. Sci.* 58, 3-9.