



Effect of a dietary supplementation of glycerol and maslinic acid on the muscle proteome of gilthead seabream

Silva, Tomé S.; Matos, Elisabete; Cordeiro, Odete; Wulff, Tune; Dias, Jorge; Jessen, Flemming; Rodrigues, Pedro

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Silva, T. S., Matos, E., Cordeiro, O., Wulff, T., Dias, J., Jessen, F., & Rodrigues, P. (2011). Effect of a dietary supplementation of glycerol and maslinic acid on the muscle proteome of gilthead seabream. Poster session presented at Proteomic Forum, Berlin, Germany, .

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Effect of a dietary supplementation of glycerol and maslinic acid on the muscle proteome of gilthead seabream

Tomé S. Silva^{a,b,*}
 Elisabete Matos^a
 Odete Cordeiro^a
 Tune Wulff^b
 Jorge Dias^a
 Flemming Jessen^b
 Pedro Rodrigues^a

^aCCMAR
 Centro de Ciências do Mar do Algarve,
 Universidade do Algarve,
 Faro, Portugal.


^bDTU Food
 Fødevareinstituttet,
 Danmark Tekniske Universitet,
 Lyngby, Denmark.



INTRODUCTION

- ✓ The onset of **post-mortem muscle degradation** depends on the availability of energy stored as glycogen;
- ✓ **Degradation of muscle tissue** often impairs fish meat quality;
- ✓ **Glycerol** and **maslinic acid** are known to **modulate glycogen storage and mobilization**, respectively;
- ✓ The use of a **proteomic approach** can be beneficial, by providing **untargeted information** about the impact of a dietary supplementation of glycerol and maslinic acid on reared fish.


EXPERIMENTAL DESIGN




RESULTS

Detoxification processes

aldehyde dehydrogenase (ALDH6A1)




aldehyde dehydrogenase (ALDH7A1)




Energy homeostasis and signaling

phosphohistidine phosphatase




adenylate kinase




Oxidative stress


transferrin




transferrin



heat shock protein cognate (HSC70)



heat shock protein cognate (HSC71)



CONCLUSIONS

- ✓ Dietary supplementation of **glycerol** and **maslinic acid** has a **relatively low impact** on the soluble muscle proteome of *Sparus aurata*, **especially when combined** (synergistic effect);
- ✓ Affected pathways seem to be related mostly to **detoxification processes**, **energy homeostasis**, **signaling**, **cytoskeleton** and the cellular response to **oxidative stress**;
- ✓ **Correlation with non-proteomic time-dependent data** will allow a **full assessment of the potential** use of these substances as dietary supplements for fish feeds.