

Technical University of Denmark



The Effect of Fertilizer Applications on the Bio-Availability of the CD and ZN

Lorenz, S.; Hamon, R.; McGrath, S.P.; Holm, P.; Christensen, Thomas Højlund

Published in:

Proceedings of 4th Conference on Current Research Developments in Soil and Related Sciences

Publication date:

1993

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Lorenz, S., Hamon, R., McGrath, S. P., Holm, P., & Christensen, T. H. (1993). The Effect of Fertilizer Applications on the Bio-Availability of the CD and ZN. In Proceedings of 4th Conference on Current Research Developments in Soil and Related Sciences London: British Soil Science Society.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

THE EFFECT OF FERTILIZER APPLICATIONS ON THE BIO-AVAILABILITY OF CD AND ZN

S. Lorenz¹⁾, R. Hamon¹⁾, S.P. McGrath¹⁾, P. Holm²⁾, T. Christensen²⁾

¹⁾ Soil Science Department, AFRC Rothamsted Experimental Station, Harpenden, Herts AL5 2JQ, U.K.

²⁾ Department of Environmental Engineering, Technical University, 2800 Lyngby, DK

Concentrations of heavy metals in soil solution determine their uptake by plants. The effect of fertilizers have been largely neglected in former experiments on the bio-availability of heavy metals and their concentrations in soil solution. An experiment was conducted to study the changes over time of the concentrations of Cd and Zn in soil solution and relate these to their uptake by radish plants.

Radish was grown under controlled conditions in soil that had been contaminated with heavy metals due to the past application of sewage sludge. Fertilizer solution (NH_4NO_3 , KNO_3 , MgSO_4) was added daily at a constant rate according to the total requirements of the plants in order to avoid the precipitation of CaSO_4 at the bottom of the pots. The soil was kept at 60% of the water holding capacity and daily water use by the plants was recorded. Over a period of 45 days, soil solution was obtained at 5 day intervals by displacement with water from three replicate pots. The plants were then cut and analysed.

The total fertilizer added equalled the plants' requirements at the final harvest but exceeded them during most of the experiment. Excess fertilizer concentrations were closely related to large increases in the concentrations of major and heavy metal ions in soil solutions and to a decrease in soil pH. Concentrations in the plant leaves reflected the increased concentrations of Cd and Zn in solution.

In order to study effects the plants have on metal solubility and to obtain a soil solution that has not been altered by the addition of fertilizers, nutrients must be added daily according to the growing needs of the plants. An example is shown where ion concentrations in soil solution remained constant over time by adding fertilizer according to the plant transpiration rate.

It is concluded that studies on the solubility and availability of heavy metals where fertilizers have been added in bulk and their effects not considered should be interpreted with caution.