

Technical University of Denmark



## Radioactivity in the Risø District July-December 2010

Nielsen, Sven Poul; Andersson, Kasper Grann; Miller, Arne

*Publication date:*  
2011

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Nielsen, S. P., Andersson, K. G., & Miller, A. (2011). Radioactivity in the Risø District July-December 2010. Roskilde: Danmarks Tekniske Universitet, Risø Nationallaboratoriet for Bæredygtig Energi. (Denmark. Forskningscenter Risoe. Risoe-R; No. 1779(EN)).

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Radioactivity in the Risø District July-December 2010

Risø-R-Report

Sven P. Nielsen, Kasper G. Andersson and Arne Miller  
Risø-R-1779(EN)  
June 2011

Risø DTU  
National Laboratory for Sustainable Energy

---



**Author:** Sven P. Nielsen, Kasper G. Andersson and Arne Miller  
**Title:** Radioactivity in the Risø District July-December 2010  
**Division:** Radiation Research

**Risø-R-1779(EN)**  
**June 2011**

**Abstract (max. 2000 char.):** The environmental surveillance of the Risø environment was continued in July - December 2010. The mean concentrations in air were:  $0.34 \pm 0.26 \mu\text{Bq m}^{-3}$  of  $^{137}\text{Cs}$ ,  $2.58 \pm 1.09 \text{ mBq m}^{-3}$  of  $^7\text{Be}$  and  $0.23 \pm 0.19 \text{ mBq m}^{-3}$  of  $^{210}\text{Pb}$  ( $\pm 1$  S.D.;  $N = 26$ ). The depositions by precipitation at Risø in the second half of 2010 were:  $0.045 \text{ Bq m}^{-2}$  of  $^{137}\text{Cs}$ ,  $744 \text{ Bq m}^{-2}$  of  $^7\text{Be}$ ,  $47.0 \text{ Bq m}^{-2}$  of  $^{210}\text{Pb}$  and  $< 1.2 \text{ kBq m}^{-2}$  of  $^3\text{H}$ . The average background dose rate (TLD) at Risø (Zone I) was  $65 \text{ nSv h}^{-1}$  compared with  $58 \pm 7 \text{ nSv h}^{-1}$  ( $\pm 1$  S.D.;  $N = 4$ ) in the four zones around Risø.

**ISSN 0106-2840**  
**ISBN 978-87-550-3907-0**

**Contract no.:**

**Group's own reg. no.:**  
1400103-07

**Sponsorship:**

**Cover :**

**Pages: 24**  
**Tables: 14**  
**References:**

Information Service Department  
Risø National Laboratory for  
Sustainable Energy  
Technical University of Denmark  
P.O.Box 49  
DK-4000 Roskilde  
Denmark  
Telephone +45 46774005  
[bibl@risoe.dtu.dk](mailto:bibl@risoe.dtu.dk)  
Fax +45 46774013  
[www.risoe.dtu.dk](http://www.risoe.dtu.dk)

# Contents

Table 1.	Radionuclides in air	5
Table 2.1.	Radionuclides in precipitation	6
Table 2.2.	Radionuclides in precipitation	6
Table 2.3.	Tritium in precipitation	7
Table 2.4.	Tritium in precipitation	7
Table 3.1.	Radionuclides in sediment samples	8
Table 4.1.	Radionuclides in seawater	8
Table 4.2.	Tritium in seawater	8
Table 5.1.	Radionuclides in grass	9
Table 5.2.	Radionuclides in sea plants	10
Table 7.1.	Waste water	11
Table 8.1.	Background dose rates around the border of Risø (TLD)	12
Table 8.2.	Background dose rates around Risø (TLD)	13
Table 8.3.	Terrestrial dose rates at the Risø zones (NaI(Tl) detector)	14
Fig. 1.	Map of Risø	15
Fig. 1.1.	Caesium-137 in air	16
Fig. 1.2.	Beryllium-7 and lead-210 in air	16
Fig. 2.3.1	Tritium in precipitation (1 m <sup>2</sup> rain collector)	17
Fig. 2.3.2	Tritium in precipitation (10 m <sup>2</sup> rain collector)	17
Fig. 3.1	Caesium-137 in sediment samples	18
Fig. 4.1	Caesium-137 in seawater	19
Fig. 4.2	Tritium in seawater	19
Fig. 7.1	Total-beta radioactivity in waste water	20
Fig. 8.1.	Map of Risø with locations for TLD measurements	21
Fig. 8.2.	The environment of Risø	22



Table 1. Radionuclides in ground level air collected at Risø (cf. Figs. 1, 1.1 and 1.2), July - December 2010. (Unit:  $\mu\text{Bq m}^{-3}$ )

Date	$^7\text{Be}$	$^{137}\text{Cs}$	$^{210}\text{Pb}$
28-Jun-10 – 05-Jul-10	3131	0.224	170
05-Jul-10 – 12-Jul-10	3616	0.221	253
12-Jul-10 – 19-Jul-10	3848	0.124	207
19-Jul-10 – 26-Jul-10	3880	0.209	221
26-Jul-10 – 02-Aug-10	2407	0.105	145
02-Aug-10 – 09-Aug-10	3468	0.264	205
09-Aug-10 – 16-Aug-10	2983	0.132	220
16-Aug-10 – 24-Aug-10	2504	0.097	178
24-Aug-10 – 30-Aug-10	2317	0.083	88
30-Aug-10 – 06-Sep-10	3639	0.346	199
06-Sep-10 – 13-Sep-10	3999	0.702	422
13-Sep-10 – 20-Sep-10	2133	0.076	74
20-Sep-10 – 27-Sep-10	2720	0.309	318
27-Sep-10 – 04-Oct-10	2480	0.568	250
04-Oct-10 – 11-Oct-10	5255	0.926	917
11-Oct-10 – 18-Oct-10	1040	0.529	245
18-Oct-10 – 25-Oct-10	2452	0.178	101
25-Oct-10 – 01-Nov-10	2968	0.308	232
01-Nov-10 – 08-Nov-10	2455	0.237	95
08-Nov-10 – 15-Nov-10	1353	0.206	49
15-Nov-10 – 22-Nov-10	639	0.398	81
22-Nov-10 – 29-Nov-10	1120	0.260	94
29-Nov-10 – 06-Dec-10	2075	0.393	204
06-Dec-10 – 13-Dec-10	1210	0.230	55
13-Dec-10 – 20-Dec-10	1836	0.818	213
20-Dec-10 – 27-Dec-10	1598	0.980	602
Mean	2582	0.343	225
SD	1093	0.259	185

Table 2.1. Radionuclides in precipitation in the 10 m<sup>2</sup> rain collector at Risø (cf. Fig. 1), July - December 2010. (Unit: Bq m<sup>-3</sup>)

Month	<sup>7</sup> Be	<sup>137</sup> Cs	<sup>210</sup> Pb
July	2833	0.215	154
August	1544	0.088	106
September	1620	0.160	133
October	1630	0.063	100
November	1232	0.026	27
December	620	0.079	80

Table 2.2. Radionuclides in precipitation in the 10 m<sup>2</sup> rain collector at Risø (cf. Fig. 1), July - December 2010. (Unit: Bq m<sup>-2</sup>)

Month	Precipitation (m)	<sup>7</sup> Be	<sup>137</sup> Cs	<sup>210</sup> Pb
July	0.050	143	0.0108	7.8
August	0.188	291	0.0165	20.0
September	0.041	67	0.0066	5.5
October	0.064	104	0.0040	6.4
November	0.081	100	0.0021	2.2
December	0.063	39	0.0050	5.1
Sum	0.487	744	0.0450	47.0

Table 2.3. Tritium in precipitation collected at Risø (cf. Figs. 1, 2.3.1 and 2.3.2). July - December 2010. (Unit: kBq m<sup>-3</sup>)

Month	1 m <sup>2</sup> rain collector*	10 m <sup>2</sup> rain collector*
July	2.1	< 2.1
August	< 2.1	2.2
September	< 2.1	< 2.1
October	< 2.1	3.3
November	< 2.1	< 2.1
December	< 2.1	< 2.1
Double determinations*.		

Table 2.4. Tritium in precipitation collected at Risø (cf. Fig. 1). July - December 2010. (Unit: kBq m<sup>-2</sup>)

Month	Precipitation (m)	1 m <sup>2</sup> rain collector	10 m <sup>2</sup> rain collector
July	0.050	0.105	< 0.105
August	0.188	< 0.395	0.414
September	0.041	< 0.086	< 0.086
October	0.064	< 0.134	0.211
November	0.081	< 0.170	< 0.170
December	0.063	< 0.132	< 0.132
Sum	0.487	< 1.023	< 1.118



*Table 3.1. Radionuclides in sediment samples collected at Bolund in Roskilde Fjord.(cf. Fig. 3.1) July - December 2010. (Unit: Bq kg<sup>-1</sup> dry)*

Date	<sup>137</sup> Cs	K*
5 July	1.8	16.7
*Unit: g kg <sup>-1</sup> dry		

*Table 4.1. Radionuclides in seawater collected in Roskilde Fjord (cf. Fig. 4.1) July - December 2010. (Unit: Bq m<sup>-3</sup>)*

Date	<sup>137</sup> Cs
1 July	12.4

*Table 4.2. Tritium in seawater collected in Roskilde Fjord (Risø pier) (cf. Fig. 4.2) July - December 2010.*

Month	kBq m <sup>-3</sup>
July	< 2.1 *
August	< 2.1 *
September	< 2.1 *
October	< 2.1 *
November	2.3 *
December	< 2.1 *
* Double determinations	

Table 5.1. Radionuclides in grass (\* snow) collected at Risø (near the Waste Treatment Station (cf. Fig. 1)), July - December 2010. (\*\*Measured on bulked ash samples)

Week no. or month	Date	K (g kg <sup>-1</sup> fresh)	<sup>137</sup> Cs (Bq kg <sup>-1</sup> fresh)	<sup>137</sup> Cs (Bq m <sup>-2</sup> )
27	5 July	7.9	<0.6	
28	12 July	6.5	<0.6	
29	19 July	4.5	<0.5	
30	26 July	5.5	<0.6	
31	2 August	4.7	<0.5	
32	9 August	4.8	<0.5	
33	16 August	5.6	<0.6	
34	23 August	4.0	<0.4	
35	30 August	4.3	<0.4	
36	6 September	4.5	<0.5	
37	13 September	4.2	<0.5	
38	20 September	4.1	<0.5	
39	27 September	5.4	<0.5	
40	4 October	5.9	<0.6	
41	11 October	3.9	<0.4	
42	18 October	4.9	<0.4	
43	25 October	3.9	<0.4	
44	1 November	3.5	<0.4	
45	8 November	2.3	<0.5	
46	15 November	3.9	<0.5	
47	22 November	2.6	<0.3	
48	29 November*	-	<0.3	
49	6 December*	<0.2	<0.3	
50	13 December*	<0.1	<0.3	
51	20 December*	<0.2	<0.4	
52	27 December*	<0.1	<0.1	
**July		6.1	0.068	0.016
**August		5.4	0.118	0.057
**September		5.6	0.074	0.034
**October		4.8	0.027	0.012
**November		3.6	0.118	0.058
**December		-	-	-

*Table 5.2. Radionuclides in Fucus vesiculosus collected at Bolund in Roskilde Fjord. July - December 2010. (Unit: Bq kg<sup>-1</sup> dry)*

Date	<sup>137</sup> Cs	K*	% dry matter
5 July	4.3	35	20

\*Unit: g kg<sup>-1</sup> dry

Table 7.1. Waste water collected at Risø (cf. Fig. 1), July - December 2010.

Week number	eqv. mg KCl l <sup>-1</sup>	<sup>137</sup> Cs (Bq m <sup>-3</sup> )	<sup>131</sup> I (Bq m <sup>-3</sup> )	<sup>226</sup> Ra (Bq m <sup>-3</sup> )
27	139	163	<113	<214
28	115	178	<119	<220
29	125	106	<113	<212
30	97	<122	<116	<213
31	92	<119	<118	<217
32	93	<131	<121	<234
33	48	<123	<126	<243
34	43	<110	<110	<207
35	58	<116	<119	<219
36	111	<109	<111	<208
37	223	453	<121	<230
38	128	418	<123	<248
39	112	<115	<113	<212
40	143	184	<121	<222
41	99	<104	<115	<221
42	93	91	<64	337
43	76	<120	<117	428
44	66	<114	<114	<207
45	54	<122	<121	<230
46	48	<111	<115	<217
47	38	<109	<114	<213
48	35	<112	<109	<209
49	32	<109	<115	<213
50	28	<99	<106	<204
51	40	<116	<115	<219
52	56	<72	<135	<156
Mean	84.3			
SD	45.5			

*Table 8.1. Background dose rates around the border of Risø (cf. Fig. 8.1) measured with thermoluminescence dosimeters (TLD) in the period May 2010 – October 2010. (Results are normalized to nSv h<sup>-1</sup>)*

Location	nSv h <sup>-1</sup>
1	50
2	46
3	41
4	57
5	50
6	56
Mean	50

Table 8.2. Background dose rates around Risø (cf. Fig. 8.2 and Fig. 1) measured with thermoluminescence dosimeters (TLD) in the period May 2010 – October 2010. (Results are normalized to  $\text{nSv h}^{-1}$ )

Risø zone	Location	$\text{nSv h}^{-1}$
I	1	36
I	2	54
I	3	107
I	4	63
I	5	-
Mean		65
II	P1	69
II	P2	80
II	P3	79
II	P4	42
Mean		68
III	P1	58
III	P2	61
III	P3	61
Mean		60
IV	P1	42
IV	P2	45
IV	P3	53
IV	P4	60
IV	P5	47
IV	P6	46
IV	P7	59
Mean		50
V	P1	56
V	P2	52
V	P3	63
V	P4	41
V	P5	58
V	P6	45
V	P7	50
V	P8	63
V	P9	62
V	P10	60
Mean		55

Table 8.3. Terrestrial dose rates at the Risø zones (cf. Fig. 8.2 and Fig. 1) July - December 2010. Measured with a NaI(Tl) detector. (Unit: nSv h<sup>-1</sup>)

Risø zone	Location	July	October
I	P1	36	40
I	P2	47	48
I	P3	348	367
I	P4	41	42
I	P5	59	43
Mean		106	108
II	P1	39	42
II	P2	39	41
II	P3	36	38
II	P4	39	41
Mean		38	40
III	P1		46
III	P2		47
III	P3		41
Mean			45
IV	P1		38
IV	P2		48
IV	P3		43
IV	P4		45
IV	P5		40
IV	P6		43
IV	P7		43
Mean			43
V	P1		36
V	P2		48
V	P3		52
V	P4		41
V	P5		46
V	P6		45
V	P7		39
V	P7a		45
V	P8		41
V	P9		47
V	P10		35
Mean			43



*Fig. 1. Locations for measurements of gamma-background radiation Zone I and II (cf. Tables 8.2 and 8.3)*






Fig. 1.1. Caesium-137 in ground level air collected at Risø in July-December 2010. (Unit:  $\mu\text{Bq m}^{-3}$ )




Fig. 1.2. Beryllium-7 and lead-210 in ground level air collected at Risø in July-December 2010. (Unit:  $\mu\text{Bq m}^{-3}$ )




Fig. 2.3.1. Tritium in precipitation collected at Risø ( 1 m<sup>2</sup> rain collector ) 1980 - 2010. (Unit: kBq m<sup>-3</sup>; DL = detection limit )





Fig. 2.3.2. Tritium in precipitation collected at Risø ( 10 m<sup>2</sup> rain collector ) 1980 - 2010. (Unit: kBq m<sup>-3</sup>; DL = detection limit )



*Fig. 3.1. Caesium-137 in sediment samples collected at Bolund in Roskilde Fjord. 1980 – 2010. (Unit: Bq kg<sup>-1</sup> dry matter)*




Fig. 4.1. Caesium-137 in seawater collected in Roskilde Fjord 1980 - 2010. (Unit:  $\text{Bq m}^{-3}$ )





Fig. 4.2. Tritium in seawater collected in Roskilde Fjord 1980 - 2010. (Unit:  $\text{kBq m}^{-3}$ ; DL = detection limit)



*Fig. 7.1. Total-beta radioactivity in waste water collected at Risø 1994 - 2010.  
(Unit: eqv. mg KCl l<sup>-1</sup>)*




Fig. 8.1. Locations (1-6) for TLD measurements around the border of Risø (cf. Table 8.1).



*Fig. 8.2. Locations for measurements of background radiation around Risø in Zones III, IV and V.*





Risø DTU is the National Laboratory for Sustainable Energy. Our research focuses on development of energy technologies and systems with minimal effect on climate, and contributes to innovation, education and policy. Risø DTU has large experimental facilities and interdisciplinary research environments, and includes the national centre for nuclear technologies.

---

**Risø DTU**  
**National Laboratory for Sustainable Energy**  
**Technical University of Denmark**

Frederiksborgvej 399  
PO Box 49  
DK-4000 Roskilde  
Denmark  
Phone +45 4677 4677  
Fax +45 4677 5688

[www.risoe.dtu.dk](http://www.risoe.dtu.dk)