

Technical University of Denmark



Fumonisin in *Aspergillus niger*: Industrial and food aspects

Frisvad, Jens Christian; Nielsen, Kristian Fog; Mogensen, Jesper; Thrane, Ulf; Larsen, Thomas Ostenfeld

Publication date:
2011

Document Version
Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):
Frisvad, J. C., Nielsen, K. F., Mogensen, J., Thrane, U., & Larsen, T. O. (2011). Fumonisin in *Aspergillus niger*: Industrial and food aspects. Abstract from 26th Fungal Genetics Conference, Pacific Grove, CA, United States.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

26th Fungal Genetics Conference, Asilomar, California, USA, 14/3-20/3 2011

Poster (abstract):

Fumonisin in *Aspergillus niger*: Industrial and food aspects

Jens C. Frisvad, Kristian F. Nielsen, Jesper Mogensen, Ulf Thrane, Thomas O. Larsen, Center for Microbial Biotechnology, Department of Systems Biology, Technical University of Denmark, Søtofts Plads B. 221, DK-2800 Kgs. Lyngby, Denmark, e-mail: jcf@bio.dtu.dk

Introduction:

Fumonisin are toxic secondary metabolites from *Fusarium verticillioides* and other Fusaria, from *Tolypocladium* and *Aspergillus niger*^{1,2}. Being a generalist *Aspergillus niger* is the workhorse in a very large number of industrial applications, and is also a common contaminant in foods. Fumonisin production by *A. niger* is depending on temperature and water activity, but is produced mostly on substrates with high amounts of sugar or salt^{1,3,4}. We wanted to find out whether industrial strains could produce fumonisins in worst case scenarios and if fumonisin production was only a feature of few black aspergilli or of widespread occurrence.

Methods

The black aspergilli were grown on CYAS and YES media for 7 days at 25° C in darkness. Small agar plugs were extracted with 75% MeOH, filtered, and analyzed by gradient HPLC-MS.

Results and discussion:

All publications describing industrial applications of *A. niger* were scrutinized and approximately half of the strains used were not available for the scientific community. The other strains were available, and few of them proved to be *A. tubingensis*, *A. acidus*, *A. carbonarius* or *A. brasiliensis*. Among the real *A. niger* strains, nearly all strains ever used in biotechnology could produce fumonisins B₂, B₄ & B₆. The strains could be subdivided into two clades (representing *A. niger* and the "phylospecies" *A. awamori*), and there were fumonisin producers in both clades. Ochratoxin A was also produced by strains in both clades, but only of approximately 6% of the strains. None of the other species in the black Aspergilli produced fumonisins. One strain (NRRL 337), called the "food fungus", because it is used for single cell protein based on cheap growth substrates, produced both fumonisins and ochratoxin A. Industrial citric acid producers produced fumonisins in pure culture, so we tested whether they could produce fumonisins on citric acid production media in shake flasks, and they could indeed produce small amounts of fumonisins.

Conclusions:

Most strains of *Aspergillus niger* can produce fumonisins. In order to have entirely safe production conditions there are several possibilities:

- The gene clusters responsible for fumonisin and ochratoxin A production can be inactivated
- A non-toxicogenic strain of *A. niger* can be used for industrial applications
- A closely related species can be used industrially, fx *A. brasiliensis*, *A. vadensis*, *A. acidus*, or *A. tubingensis*, as they are not able to produce fumonisins or ochratoxins
- The fumonisin producing strains of *A. niger* can be grown under conditions, where fumonisin accumulation is not possible, but in that case a strict scheme for daily chemical control is necessary. This practice is used for *Fusarium venenatum*, used for single cell protein (Quorn). This fungus can produce trichothecenes under optimal conditions.

References:

- ¹ Frisvad, J.C., Smedsgaard, J., Samson, R.A., Larsen, T.O., and Thrane U. 2007. Fumonisin B₂ production by *Aspergillus niger*. *J. Agric. Food Chem.* **55**: 9727-9732.
- ² Nielsen, K.F., Mogensen, J.M., K Johansen, M., Larsen, T.O., Frisvad, J.C. 2009. Review of secondary metabolites and mycotoxins from the *Aspergillus niger* group. *Analytical and Bioanalytical Chemistry* **395**: 1225-1246.
- ³ Sørensen, L.M., Lametsch, R., Andersen, M.R., Nielsen, P.V. and Frisvad, J.C. 2009. Proteome analysis of *Aspergillus niger*: Lactate added in starch-containing medium can increase production of the mycotoxin fumonisin B₂ by modifying acetyl-CoA metabolism. *BMC Microbiology* **9**: 255
- ⁴ Mogensen, J.M., Nielsen, K.F., Frisvad, J.C., Samson, R.A. and Thrane, U. 2009. Effect of temperature and water activity on the production of fumonisin B₂ by *Aspergillus niger* and *Fusarium* species. *BMC Microbiology* **9**: 281