

Technical University of Denmark



## Assessment of pedometer-determined physical activity in Danish adults: the importance of non-ambulatory activities

Rothausen, Berit Worm; Gille, Maj-Britt; Bilstoft-Jensen, Anja Pia; Raustorp, Anders; Matthiessen, Jeppe

*Publication date:*  
2009

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Rothausen, B. W., Gille, M-B., Bilstoft-Jensen, A. P., Raustorp, A., & Matthiessen, J. (2009). Assessment of pedometer-determined physical activity in Danish adults: the importance of non-ambulatory activities. Poster session presented at Annual Conference of the International Society of Behavioral Nutrition and Physical Activity, Lisbon, Portugal, .

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Assessment of pedometer-determined physical activity in Danish adults: the importance of non-ambulatory activities

Rothausen BW<sup>1</sup>, Gille M-B<sup>1</sup>, Biltoft-Jensen A<sup>1</sup>, Raustorp A<sup>2</sup>, Matthiessen J<sup>1</sup>

<sup>1</sup>Department of Nutrition, National Food Institute, Technical University of Denmark, email: bewro@food.dtu.dk

<sup>2</sup>Department of Health and Behavioral Science, University of Kalmar, Sweden

## Purpose

To estimate mean values in steps/day for a representative sample of Danish adults (15-75 years) using two different conversion methods for non-ambulatory (non-step) activities.

## Methods

A simple random sample comprising 229 adults (52% men) from the Danish National Survey of Dietary Habits and Physical Activity 2007-08 wore a pedometer (Yamax SW-200 Tokyo, Japan) and recorded daily steps and non-ambulatory activities for seven consecutive days. Time spent on non-ambulatory activities was converted to step equivalents using 1) a simple conversion method (SCM) adding 200 step equivalents/min, and 2) a more complex conversion method (CCM) adding activity-specific step equivalents/min (assuming that 3 METs equals 100 steps/min) (Table 1). Very similar conversion methods have been suggested by Miller et al. 2006<sup>1</sup>. Moreover, data from a Danish pilot study (n=28) were used to adjust for double-counting during cycling. When adding steps for time spent cycling, the mean number of steps recorded during cycling found in the pilot study was subtracted from the additional steps. The addition of >10000 step equivalents/day was truncated to 10000 step equivalents/day to avoid overestimation.


**Table 1.** The selected non-ambulatory activities, their prevalence in the sample, and the steps/min rate derived from the CCM conversion method (associated MET levels were identified in the Compendium of Physical Activities<sup>2</sup>)

Activity	Participants (n=122)	Days	METs	Steps/min
Cycling, 16-19 km/h	98	306	6	200
Cycling, 16-19 km/h (adjusted for double-counting)				160
Cycling, 19-22 km/h	4	9	8	267
Cycling, 19-22 km/h (adjusted for double-counting)				227
Cycling stationary, vigorous	3	5	10.5	350
Cycling stationary, vigorous (adjusted for double-counting)				310
Weightlifting	18	38	6	200
Swimming laps	1	2	7	233
Swimming, leisurely	4	6	6	200
Swimming / water aerobics	4	4	4	133
Horseback riding*	4	4	4	133

\* Participants were instructed not to wear the pedometer during horseback riding

## Results

53% (45% of men, 62% of women) engaged in non-ambulatory activities during the registration period. The most frequently reported non-ambulatory activity was cycling, especially as transportation, which was reported by 39% (men: 31%, women: 48%) with a mean of 125 min/week (men: 107, women: 138 min/week). Our pilot study showed a mean recording of 40 steps/min during cycling. Calculated steps/day after adjusting for double-counting during cycling are shown in Figure 1 for each of the conversion methods.



**Figure 1.** Mean steps/day, raw and calculated using the SCM conversion method, adding 200 step equivalents/min, and the CCM conversion method, adding activity-specific step equivalents/min.

\* p<0.05, difference between men and women in mean steps/day

Truncation was carried out for 4-5% of all days for each conversion method. At group level the conversion methods added 1482-1494 step equivalents/day corresponding to a 17% increase. For men 1126-1138 step equivalents/day were added (13-14% increase) vs. 1875-1886 step equivalents/day for women (22% increase), thus significantly higher for women compared to men, p=0.03

## Conclusion

In populations like the Danish, where cycling and other non-ambulatory activities are popular, it will be relevant to account for these activities when assessing pedometer-determined physical activity. Adding activity-specific step equivalents/min, or simply adding 200 step equivalents/min, provides very similar results. Furthermore, when accounting for cycling an adjustment for double-counting should be considered.

<sup>1</sup> Miller R, Brown W J, Tudor-Locke C. But what about swimming and cycling? How to count non-ambulatory activity when using pedometers to assess physical activity. JPAH 2006;3:257-266.

<sup>2</sup> Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ, O'Brien WL, Bassett DR Jr, Schmitz KH, Emplaincourt PO, Jacobs DR Jr, Leon AS. Compendium of physical activities: an update of activity codes and MET intensities. Med Sci Sports Exerc. 2000;32:S498-504.

