

Technical University of Denmark



Ruthenium-catalyzed synthesis of amides from alcohols and amines: mechanistic study and application

Makarov, Ilya; Fristrup, Peter; Madsen, Robert

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Makarov, I., Fristrup, P., & Madsen, R. (2012). Ruthenium-catalyzed synthesis of amides from alcohols and amines: mechanistic study and application. Abstract from Challenges in Organic Chemistry and Chemical Biology (ISACS7), Edinburgh, United Kingdom.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Ruthenium-catalyzed synthesis of amides from alcohols and amines: mechanistic study and application

Ilya S. Makarov, Peter Fristrup, Robert Madsen*

Department of Chemistry, Technical University of Denmark, 2800 Lyngby, Denmark

The amide bond is one of the most important linkages in organic chemistry and constitutes the key functional group in peptides, polymers and many pharmaceuticals.^{1,2} Amides are often synthesized by coupling of carboxylic acids or its derivatives with amines. In spite of their efficiency these methods are not environmentally friendly since toxic reagents are used and a lot of waste is produced. Development of new synthetic methods which are both efficient and “green” continues to receive significant attraction. Recently in Madsen group a new method for amide synthesis has been discovered where alcohols and amines are coupled directly with the liberation of hydrogen gas.^{3,4} (Scheme 1).



Scheme 1. Ruthenium-catalyzed amide synthesis.

In the present work an experimental and theoretical mechanistic investigation of this reaction has been carried out. In the experimental part various kinetic aspects of the reaction were studied. Para-substituted benzyl alcohols were used as substrates to elucidate the electronic influence of the *para* substituent on the reactivity, *i.e.* a Hammett study. Deuterium-labeled substrates were employed to probe the kinetic isotope effect. By using DFT calculations with the M06 functional, energetic and structural parameters were determined for the intermediates and transition states in the proposed catalytic cycle. From the experimental and theoretical studies the rate determining step was established which is important for further optimization of the reaction.

References:

1. V. R. Pattabiraman, J. W. Bode, *Nature*, 2011, **480**, 471-479.
2. J. W. Bode, *Curr. Opin. Drug Discov. Dev.* 2006, **9**, 765-775
3. L. U. Nordstrøm, H. Vogt, R. Madsen, *J. Am. Chem. Soc.*, 2008, **130**, 17672-17673.
4. J. H. Dam, G. Osztrovszky, L. U. Nordstrøm, R. Madsen, *Chem. Eur. J.*, 2010, **16**, 6820-6827.