

Technical University of Denmark



Particle simulations of optical emissions in sprite streamers

Chanrion, Olivier; Neubert, Torsten; Stenbaek-Nielsen, H.

Published in:
Geophysical Research Abstracts

Publication date:
2008

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Chanrion, O. A., Neubert, T., & Stenbaek-Nielsen, H. (2008). Particle simulations of optical emissions in sprite streamers. *Geophysical Research Abstracts*, 10, EGU2008-A-08005.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Particle simulations of optical emissions in sprite streamers

O. Chanrion (1), T. Neubert (1) and H. Stenbaek-Nielsen (2)

(1) National Space Institute, Danish Technical University, Copenhagen, Denmark
(chanrion@space.dtu.dk) (2) Geophysical Institute, University of Fairbanks, Alaska, USA

The optical emission rates of sprites in the mesosphere contain important information on the discharge process. A precise understanding of the process is needed in order to estimate perturbations of sprites to the composition, density and temperature of the mesosphere. For this purpose, a particle code has been developed to study electric discharges in the atmosphere. The code simulates the development of discharges from seed ionization through electron avalanches into streamers. The code is based on the particle-in-cell (PIC) technique for updating the electron velocities and their positions and for calculating the self-consistent spacecharge fields. A Monte Carlo technique (MC) is used for simulating interactions with the ambient atmosphere including excitation and emission rates. The code is in 2D cylindrical coordinates, giving realistic spatial variations of the fields during the initial stages of a discharge presented here. The simulations of optical emission rates are compared to recent high-speed imaging observations of sprites, which allow for a simple and robust estimation of the instantaneous emission rate in streamer heads. The presentation will discuss electric fields, electron energetics and optical emission rates in the streamer head region.