Technical University of Denmark



Carp head-kidney leukocytes display different oxygen radical production patterns after DAMP and PAMP stimulation.

Jiménez, Natalia Ivonne Vera; Nielsen, Michael Engelbrecht

Publication date: 2012

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA):

Jiménez, N. I. V., & Nielsen, M. E. (2012). Carp head-kidney leukocytes display different oxygen radical production patterns after DAMP and PAMP stimulation.. Abstract from 12th Congress of the International Society of Developmental and Comparative Immunology (ISDCI), Fukuoka, Japan.

DTU Library Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Carp head-kidney leukocytes display different oxygen radical production patterns after DAMP and PAMP stimulation.

N.I. Vera-Jiménez¹, M.E. Nielsen¹

¹ Biological Quality Research Group, National Food Institute, Technical University of Denmark

In order to survive multicellular organisms have to be able to distinguish whether their cells are alive or death, they must detect intruders and infections, eradicate invading microorganisms and repair tissue damage. Mammalian studies have demonstrated that the immune-system capacity to distinguish between self and nonself is achieved through the recognition of pathogen-associated molecular patterns (PAMPs) in case of infection, and damage-associated molecular pattern (DAMPs) as self warning signals. It has been speculated that DAMPs and PAMPs are able to drive the nature of the immune-response leading towards pathogen eradication and/or tissue regeneration. The reactive oxygen species (ROS) have shown to play an important role during pathogen eradication and tissue repair. In fish however, the situation is not so clear, the immune-responses elicited by infection, including oxygen radical production have been relatively well described, although the involvement of the fish immune-system in tissue regeneration is not well characterized. The objective of this study is to test the ability of the fish immune-system to differentiate between PAMPs and DAMPs, and thus react to them by producing different oxygen radical profiles, corresponding to bacterial eradication and tissue regeneration respectively. Carp cultured head-kidney leukocytes were stimulated either with β -glucan (MacroGard[®]), or DAMPs obtained from a carp fibroblast cell culture. The kinetics of oxygen radical production were followed for 210 minutes using chemiluminescence. The pattern of oxygen radical production during β-glucan stimulation showed a fast and strong increase in oxygen radicals, in consistency with a pathogenic situation, where the organism targets the intruder for eradication. On the other hand, stimulation with carp fibroblast cellular components elicited a more gradual, steadier and lenient production of oxygen radicals, compatible with a tissue regeneration situation, where low doses of oxygen radicals modulate cellular proliferation and act as cellular messengers. The ability of carp head-kidney leukocytes of reacting differently to self and non-self warning signals is an important feature that opens possibilities for immune-manipulation of the tissue regeneration process in fish. Furthermore, attention is called to the significance of considering the kinetics of oxygen radical production during immune-related processes such as pathogen eradication and tissue regeneration.