

Technical University of Denmark



Trends in CO₂ exchange over a deciduous forest based on continuous eddy covariance measurements over 14 years

Pilegaard, Kim; Ibrom, Andreas; Dellwik, Ebba

Published in:
Geophysical Research Abstracts

Publication date:
2010

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Pilegaard, K., Ibrom, A., & Dellwik, E. (2010). Trends in CO₂ exchange over a deciduous forest based on continuous eddy covariance measurements over 14 years. *Geophysical Research Abstracts*, 12, EGU2010-9169.

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Trends in CO₂ exchange over a deciduous forest based on continuous eddy covariance measurements over 14 years

Kim Pilegaard (1), Andreas Ibrom (1), and Ebba Dellwik (2)

(1) Risø DTU, Biosystems Division, Roskilde, Denmark (kipi@risoe.dtu.dk, +45 4677 4101), (2) Risø DTU, Wind Energy Division, Roskilde, Denmark

The CO₂ exchange was measured by eddy covariance technique over a Danish mature beech forest (Sorø) continuously from June 1996 to December 2009. The results show a significant increase in annual net ecosystem exchange (NEE) of 30 g C m⁻² y⁻¹. Estimates of ecosystem respiration (RE) and gross ecosystem exchange (GEE) were derived from NEE. The trend in RE (-8 g C m⁻² y⁻¹) was not significant, but there was a significant increase in GEE (22 g C m⁻² y⁻¹). An analysis of the seasonal changes in NEE showed that the carbon uptake period (CUP), i.e. the period with net uptake increased significantly by 1.7 d y⁻¹, whereas the period with leaves on the trees (LP) stayed constant. Light response curves for NEE revealed that the assimilation capacity of the forest at maximum ambient light (F_∞) increased significantly by 0.7 μmol m⁻² s⁻¹ per year, whereas there was no significant trend in the apparent quantum yield.

Trends in the explaining variables were examined. No significant trends could be found in temperature, light or precipitation, whereas there was a significant increase in global atmospheric CO₂ concentration (1.9 ppmv y⁻¹).

Simulation of the NEE data from the Sorø forest by the model MAESTRA showed that the increase in CO₂ could only explain a small part of the increased uptake capacity of the forest. A much larger part could be explained by an increase in N-content of the leaves. Such an increase is not yet documented. In general the deposition of atmospheric N has decreased, but the forest is already N-saturated and the availability of N in the soil might have increased due to climate change. Other possibilities for explaining the observed trends are discussed.