

Technical University of Denmark



A simple alternative to the light-use efficiency approach that considers canopy photosynthesis saturation with light

Ibrom, Andreas; Oltchev, A.; Panferov, O.; Kreinlein, H.; June, T.; Rakkibu, G.; Gravenhorst, G.

Published in:
Geophysical Research Abstracts

Publication date:
2008

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Ibrom, A., Oltchev, A., Panferov, O., Kreinlein, H., June, T., Rakkibu, G., & Gravenhorst, G. (2008). A simple alternative to the light-use efficiency approach that considers canopy photosynthesis saturation with light. *Geophysical Research Abstracts*, 10, EGU2008-A-06950.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



A simple alternative to the light-use efficiency approach that considers canopy photosynthesis saturation with light

Ibrom A. (1,3) Oltchev, A. (2,3), Panferov, O. (3), Kreinlein, H. (3), June, T. (4,5), Rakkibu, G. (3), Gravenhorst, G. (3)

(1) Biosystems Department, Risø National Laboratory, Denmark, Technical University of Denmark (DTU), 4000 Roskilde Denmark, (andreas.ibrom@risoe.dk/ Fax: ++45 46773160 / Phone: ++45 46774177), (2) A.N. Severtsov Institute of Ecology and Evolution of RAS, Moscow, Russia (3) Institute of Bioclimatology, Germany, Georg-August-University, Göttingen, (4) SEAMEO BIOTROP, Bogor, Indonesia, (5) Institute Pertanian Bogor (IPB), Bogor, Indonesia

Abstract

We report eddy covariance CO₂ flux and meteorological observations from a tropical rain forest in Indonesia that clearly show saturation of canopy photosynthesis (P_g) at relatively moderate incident and absorbed photosynthetically active radiation (PAR) (Ibrom et al. 2008). CO₂ fluxes were measured with and open-path fast response IRGA (LI-7500, LI-COR, NE, USA) and a 3D ultrasonic anemometer (USA1, Metek, Elmshorn, Germany). PAR absorption was calculated with a one dimensional model (Oltchev et al. 2007) and complied with measured short wave albedo at the site, the Lore Lindu National Park, Central Sulawesi, Indonesia.

We investigated how this saturation propagated moving from half hourly to larger time scales and found that in the tropical climate, saturation was the dominant phenomenon of the photosynthetic light response both at daily and monthly time scales. From this we conclude that the fundamental assumption of the very often used light-use efficiency approach (Monteith 1972), i.e. that light use efficiency is a constant under non limiting water vapour saturation deficit (D), temperature and soil water availability,

does not apply at this site. We also revisited studies from other forests and climates and found indications for P_g saturation. These were however masked by combining the light-use efficiency approach with D reduction functions that are used to include certain reduction of ε at dry (and at the same time sunny) weather situations in the simulations.

Apart from the critical evaluation of the light-use efficiency approach we also suggest a similarly simple approach that is able to include both saturation and reduction of P_g at high D levels based on a simple Blackman response approach. This approach uses the light use efficiency approach only for non saturating condition and a constant maximum P_g (P_{max}) when absorbed PAR exceeds saturating levels. As D was found to reduce P_{max} , the PAR level at which P_g saturated became a well described function of D .

We advocate replacing the current version of the light-use efficiency approach by the combined Blackman response when simulating GPP from remote sensing products. Future research should focus to investigate biome characteristic relationships of the additional parameter, P_{max} , with climate and canopy structure.

Acknowledgements

This work was funded by the DFG and part of the collaborated research project “Stability of rain forest margin areas” STORMA, SFB 552.

References

- Ibrom, A., A. Oltchev, T. June, H. Kreilein, G. Rakkibu, T. Ross, O. Panferov and G. Gravenhorst. 2008. Variation in photosynthetic light-use efficiency in a mountainous tropical rain forest in Indonesia. *Tree Physiol.* accepted for publication
- Monteith, J.L. 1972. Solar radiation and production in tropical ecosystems. *J. Appl. Ecol.* 9
- Oltchev, A., A. Ibrom, T. Ross et al. 2007. A modelling approach for simulation of water and carbon dioxide exchange between multi-species tropical rain forest and the atmosphere. *Ecol. Modell.* doi:10.1016/j.ecolmodel.2007.10.021