

Technical University of Denmark



Marginal thinning in Northwest Greenland during 2002-2011

Khan, Shfaqat Abbas; Kjær, K. H.; Wahr, J. M.; Bevis, M.; Korsgaard, N.; Bjørk, A. A.; Kjeldsen, K. K.; Timm, L. M.; Dam, T. v.

Published in:
Geophysical Research Abstracts

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Khan, S. A., Kjær, K. H., Wahr, J. M., Bevis, M., Korsgaard, N., Bjørk, A. A., ... Dam, T. V. (2012). Marginal thinning in Northwest Greenland during 2002-2011. *Geophysical Research Abstracts*, 14, EGU2012-1852.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Marginal thinning in Northwest Greenland during 2002-2011

S. A. Khan (1), K. H. Kjær (2), J. M. Wahr (3), M. Bevis (4), N. Korsgaard (2), A. A. Bjørk (2), K. K. Kjeldsen (2), L. H. Timm (2), and T. v. Dam (5)

(1) DTU - Space, Geodetic, Copenhagen OE, Denmark (abbas@space.dtu.dk, +45 3536 2475), (2) Centre for GeoGenetics, Natural History Museum, University of Copenhagen, Copenhagen, Denmark., (3) Department of Physics and Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, USA., (4) Ohio State University, Columbus, Ohio, USA., (5) University of Luxembourg, Luxembourg.

Many glaciers along the southeast and northwest coast of Greenland have accelerated, increasing the Greenland ice sheet's (GrIS) contribution to global sea-level rise. Here, we map elevation changes in northwest Greenland during 2003-2009 using high-resolution Ice, Cloud and land Elevation Satellite (ICESat) laser altimeter data (Zwally, 2010) supplemented with altimeter surveys from NASA's Airborne Topographic Mapper (ATM) during 2002-2011 (Krabill, 2011). We use the measurements of elevation change to estimate catchment-wide ice volume loss (convert is to mass loss) and compare with independent measurements from GPS and the Gravity Recovery and Climate Experiment (GRACE) satellite gravity mission, launched in March, 2002. The GRACE results provide a direct measure of mass loss averaged over the entire northwest sector, while the GPS data are used to monitor crustal uplift caused by ice mass loss close to the sites.

GPS data from a long term site at Thule Airbase show accelerated uplift starting in 2005 and a minor deceleration in 2009-2010. The deceleration is more dominant at GPS stations deployed in 2007 in northwest Greenland as part of the Greenland GPS Network (GNET). Independently, all three methods suggest increased ice loss in northwest Greenland starting in 2005 and a slowdown in 2009-2010.