

Technical University of Denmark



Synthesis and Activation of Catalysts for Biofuel Synthesis in an Environmental Transmission Electron Microscope

Duchstein, Linus Daniel Leonhard; Wu, Qiong Xiao; Elkjær, Christian Fink; Sharafutdinov, Irek; Hansen, Thomas Willum; Damsgaard, Christian Danvad; Wagner, Jakob Birkedal

Publication date:
2012

[Link back to DTU Orbit](#)

Citation (APA):

Duchstein, L. D. L., Wu, Q., Elkjær, C. F., Sharafutdinov, I., Hansen, T. W., Damsgaard, C. D., & Wagner, J. B. (2012). Synthesis and Activation of Catalysts for Biofuel Synthesis in an Environmental Transmission Electron Microscope. Abstract from 2nd International Symposium on Advanced Electron Microscopy for Catalysis and Energy Storage Materials, Berlin, Germany.

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Synthesis and Activation of Catalysts for Biofuel Synthesis in an Environmental Transmission Electron Microscope

Linus D. L. Duchstein^{1, *}, Qiongxiao Wu², Christian Fink Elkjær³, Irek Sharafutdinov³, Thomas W. Hansen¹, Christian D. Damsgaard¹, Jakob B. Wagner¹

¹ Center for Electron Nanoscopy (Cen), Technical University of Denmark, Fysikvej 307, DK-2800 Kgs. Lyngby, Denmark

² Department of Chemical and Biochemical Engineering, Technical University of Denmark, Soeltofts Plads, Building 229, DK-2800 Kgs. Lyngby, Denmark

³ Department of Physics, Technical University of Denmark, Fysikvej, Building 307, DK-2800 Kgs. Lyngby, Denmark

The synthesis of transportation fuels from sustainable resources requires new and better production paths. Our approach is to use biogas to synthesize alcohols, such as methanol or higher alcohols for fuel and other chemical products. For the production of methanol a reduction of processing temperature and pressure to lower the process cost and make the product more competitive is desired. Higher alcohols are in general favorable over methanol due to their high energy density and ease of use in current internal combustion engines. However, better catalysts for this reaction are needed to increase the poor yield of higher alcohols in present production routes [1].

The Catalysis for Sustainable Energy Project (CASE) at the Technical University of Denmark aims at discovering new and improved catalysts based on density functional theory (DFT) and testing the chemical reactivity of the most promising candidates experimentally. Transmission electron microscopy (TEM) is used for microstructural characterization and provides feedback for both theory and synthesis.

We have studied the catalysts close to their working conditions in an environmental transmission electron microscope (ETEM) equipped with a differential pumping system to confine a controlled gas flow around the specimen, allowing observation in a gaseous environment. Using heating holders, dynamic information about catalysts in their working state can be gained using a variety of TEM techniques *in situ*. [2,3]. Here, we present recent ETEM studies of CuNi and NiGa catalysts for alcohol synthesis using High-Resolution TEM (HRTEM), energy electron-loss spectroscopy (EELS), Energy-Dispersive X-ray Spectroscopy (EDX). Complementary observations have been done using *in-situ* X-Ray Diffraction (XRD). We focus on structural changes during the catalysts synthesis and activation in a reducing atmosphere at elevated temperature. Changes in phase and particle size distribution with respect to the temperature can be directly observed and correlated to catalytic activity and integral phase information from the *in-situ* XRD.




Figure 1: a) EELS spectrum and b) HRTEM image with FFT of a single CuNi particle taken at 300°C in 1.3mbar H₂ atmosphere.

References

- 1) Spivey J. J., Egbibi, A. *Chem. Soc. Rev.* 2007; 36: 1514 - 1528
- 2) Datye A. K. *Journal of Catalysis* 2003; 216: 144 - 154
- 3) Hansen T. W., Wagner J. B., Hansen P. L., Dahl S., Topsøe H., Jacobsen C. J. H. *Science* 2001; 294(5546): 1508 - 1510