

Technical University of Denmark



## In Vitro Pulmonary Toxicity of Metal Oxide Nanoparticles

Cupi, Denisa; Dreher, Kevin

*Publication date:*  
2013

[Link back to DTU Orbit](#)

*Citation (APA):*

Cupi, D., & Dreher, K. (2013). In Vitro Pulmonary Toxicity of Metal Oxide Nanoparticles. Abstract from SOT Annual Meeting and ToxExpo, San Francisco, United States.

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**CONTROL ID:** 1242081

**CONTACT (NAME ONLY):** Kevin Dreher

**Abstract Details**

**PRESENTATION TYPE:** Platform or Poster

**CURRENT CATEGORY:** Nanotoxicology | Inhalants & Cardiopulmonary | Alternatives to Mammalian Models

**KEYWORDS:** nanoparticle, in vitro toxicity, pulmonary.

**DATE/TIME LAST MODIFIED:** September 29, 2011, 10:18 AM

**DATE/TIME SUBMITTED:** September 29, 2011, 10:18 AM

**Abstract**

**TITLE:** *IN VITRO* PULMONARY TOXICITY OF METAL OXIDE NANOPARTICLES

**AUTHORS (LAST NAME, FIRST NAME):** Cupi, Denisa<sup>1</sup>; Dreher, Kevin<sup>2</sup>

**SPONSOR NAME:** None

**INSTITUTIONS (ALL):** 1. Technical University of Denmark, Kongens Lyngby, Denmark.

2. US EPA, Research Triangle Park, NC, United States.

**ABSTRACT BODY:** Nanomaterials (NMs) encompass a diversity of materials with unique physicochemical characteristics which raise concerns about their potential risk to human health. Rapid predictive testing methods are needed to characterize NMs health effects as well as to screen and prioritize NMs for comprehensive toxicological assessments. BEAS2B human bronchial epithelial cells were employed to assess the *in vitro* pulmonary toxicity of 4 TiO<sub>2</sub> and 4 CeO<sub>2</sub> particles varying in size (6 - 1288nm) and crystalline structure. Exposures were conducted over several concentrations for each endpoint examined. No BEAS2B cytotoxicity was observed for any particle following a 24hr exposure to concentrations up to 100 µg/ml. The ability of TiO<sub>2</sub> and CeO<sub>2</sub> particles to induce inflammation and oxidative stress was assessed by gene induction using RT-PCR. At 50 µg/ml maximal IL-8 and IL-6 gene induction by TiO<sub>2</sub> and CeO<sub>2</sub> NMs (6 - 8nm) was observed at 6hr and 24hr post-exposure, respectively. Smaller TiO<sub>2</sub> and CeO<sub>2</sub> NMs induced greater induction of IL-8 and IL-6 mRNA levels compared to larger sized particles. CeO<sub>2</sub> 8nm NMs produced the greatest induction of IL-8 and IL-6 mRNA levels. At 50 µg/ml all TiO<sub>2</sub> and CeO<sub>2</sub> particles induced similar increases in HO-1 mRNA levels at 6hr and 24hr post-exposure, respectively. The pattern of HO-1 gene induction was inconsistent with a role of oxidative stress in metal oxide induced BEAS2B cytokine gene expression. Pretreatment of BEAS2B cells with IKK inhibitor III BMS-345541 completely inhibit 25nm TiO<sub>2</sub> and 69nm CeO<sub>2</sub> NM induction of IL-8, IL-6 and HO-1 gene expression indicating a role of NFκB in these responses. A cell-based ELISA for NFκB p65 phosphorylation revealed rapid Ser536 phosphorylation in BEAS2B cells following exposure to 50 µg/ml of TiO<sub>2</sub> and CeO<sub>2</sub> NMs with sizes ≥25nm. Results demonstrate the ability to employ *in vitro* methods to assess NM induced pulmonary toxicity. Many of the responses were found not to be totally dependent on NM size/surface area suggesting composition and surface properties play a role in mediating NM toxicity. This abstract does not reflect EPA policy.