

Technical University of Denmark



Effect of diet on the human gut microbiota

Bahl, Martin Iain

Publication date:
2013

[Link back to DTU Orbit](#)

Citation (APA):

Bahl, M. I. (2013). Effect of diet on the human gut microbiota. Abstract from 8th Danish Conference on Biotechnology and Molecular Biology, Vejle, Denmark.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract for oral presentation at 8th Danish Conference on Biotechnology and Molecular Biology, Food Biotechnology and Nutrigenomics, May 30 – 31, 2013, Hotel Munkebjerg, Vejle

Effect of diet on the human gut microbiota

Martin Iain Bahl, National Food Institute, Technical University of Denmark, Mørkhøj Bygade 19, 2860 Søborg, Denmark. E-mail: mbah@food.dtu.dk

Abstract

The gut microbiota plays an important role for humans in both health and disease. It is therefore important to understand how and to what extent choice of diet may influence the microbial community and the effects this has on the host. The variation in the normal human gut microbiota may however impede the discovery of correlations between dietary changes and compositional shifts in the microbiota by masking such effects. Although specific functional food ingredients, such as prebiotics, are known to have measurable effects on e.g. abundance of bifidobacteria, it is nevertheless clear that induced shifts in gut microbiota show large inter-individual variations. It thus seems plausible that knowing the microbiota composition could facilitate predictions as to how the community will react to dietary interventions thus moving towards some degree of personalised dietary recommendations. During a 6-month randomised, controlled dietary intervention following either the New Nordic Diet recommendations or Average Danish Diet (n=62) almost no significant differences in the gut microbiota composition caused by the different diets were observed using qPCR analysis of 33 selected bacterial groups. By stratifying subjects into two enterotypes, distinguished simply by the *Prevotella/Bacteroides* ratio (*P/B*), we were however able to detect several significant changes in the gut microbiota composition resulting from the intervention. The enterotypes were found to be stable during this 6 month study and furthermore we have observed in another study, that the fecal microbiota seems to group by *P/B* ratio during the first 3 years of life. The use of *P/B* to distinguish enterotypes appears to be a simple approach to stratify individuals, which may be useful to assess the effect of diets and other treatments on the gut microbiota.