

Technical University of Denmark



Reconstruction of the biosynthetic pathway for the core fungal polyketide scaffold rubrofusarin in *Saccharomyces cerevisiae*

Rugbjerg, Peter; Naesby, Michael; Mortensen, Uffe Hasbro; Frandsen, Rasmus John Normand

Published in:
Microbial Cell Factories

Link to article, DOI:
[10.1186/1475-2859-12-31](https://doi.org/10.1186/1475-2859-12-31)

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Rugbjerg, P., Naesby, M., Mortensen, U. H., & Frandsen, R. J. N. (2013). Reconstruction of the biosynthetic pathway for the core fungal polyketide scaffold rubrofusarin in *Saccharomyces cerevisiae*. *Microbial Cell Factories*, 12(1), [31]. DOI: 10.1186/1475-2859-12-31

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH

Open Access

Reconstruction of the biosynthetic pathway for the core fungal polyketide scaffold rubrofusarin in *Saccharomyces cerevisiae*

Peter Rugbjerg^{1,2}, Michael Naesby¹, Uffe H Mortensen² and Rasmus JN Frandsen^{2*}

Abstract

Background: Fungal polyketides include commercially important pharmaceuticals and food additives, e.g. the cholesterol-lowering statins and the red and orange monascus pigments. Presently, production relies on isolation of the compounds from the natural producers, and systems for heterologous production in easily fermentable and genetically engineerable organisms, such as *Saccharomyces cerevisiae* and *Escherichia coli* are desirable. Rubrofusarin is an orange polyketide pigment that is a common intermediate in many different fungal biosynthetic pathways.

Results: In this study, we established a biosynthetic pathway for rubrofusarin in *S. cerevisiae*. First, the *Fusarium graminearum* gene encoding polyketide synthase 12 (*PKS12*) was heterologously co-expressed with the *Aspergillus fumigatus* gene encoding phosphopantetheinyl transferase (*npgA*) resulting in production of YWA1. This aromatic heptaketide intermediate was converted into nor-rubrofusarin upon expression of the dehydratase gene *aurZ* from the aurofusarin gene cluster of *F. graminearum*. Final conversion into rubrofusarin was achieved by expression of the O-methyltransferase encoding gene *aurJ*, also obtained from the aurofusarin gene cluster, resulting in a titer of 1.1 mg/L. Reduced levels of rubrofusarin were detected when expressing *PKS12*, *npgA*, and *aurJ* alone, presumably due to spontaneous conversion of YWA1 to nor-rubrofusarin. However, the co-expression of *aurZ* resulted in an approx. six-fold increase in rubrofusarin production.

Conclusions: The reconstructed pathway for rubrofusarin in *S. cerevisiae* allows the production of a core scaffold molecule with a branch-point role in several fungal polyketide pathways, thus paving the way for production of further natural pigments and bioactive molecules. Furthermore, the reconstruction verifies the suggested pathway, and as such, it is the first example of utilizing a synthetic biological “bottom up” approach for the validation of a complex fungal polyketide pathway.

Keywords: Iterative polyketide synthase, Pigment, Heterologous production, YWA1, Nor-rubrofusarin, Aurofusarin

Background

Polyketides exhibit large structural diversity and have attracted considerable interest from the pharmaceutical industry as they form a chemical class rich in bioactive molecules. The biological activities of fungal polyketides span from potent carcinogenic toxins (aflatoxin) over immunosuppressants (dalesconols), anti-tumor compounds (aspergillide A) to block-buster cholesterol-lowering drugs (statins) [1-3]. In addition, many fungal polyketides also display vibrant colors due to their

polyaromatic structures, which have made them attractive for the textile and food industries as natural dyes [4,5].

Fungal polyketides of commercial interest are often produced in nature by species that are not easily cultured in standard liquid fermentation setups, due to their filamentous growth, and formation of dense mycelium clumps resulting in increased viscosity and reduced oxygen solubility. The transfer of such biosynthetic pathways to standard industrial cell factories, such as *Saccharomyces cerevisiae*, could offer an attractive alternative as it would allow easier fermentation and process optimization via the existing metabolic engineering strategies. However, only a few examples of successful heterologous expression

* Correspondence: rasf@bio.dtu.dk

²Søltøfts Plads Building 223, Kgs Lyngby DK-2800, Denmark

Full list of author information is available at the end of the article

of genes encoding fungal polyketide synthases in yeast have been reported in literature. These include the 6-MSA synthase gene from *Penicillium patulum* resulting in 6-methylsalicylic acid (6-MSA) production [6-8], *rdc1/rdc5* from *Pochonia chlamydosporia* [9] resulting in (*R*)-monocillin II production, and *hpm3/hpm8* resulting in hypothemycin production [10]. Most recently the individual expression of three PKSs from *Chaetomium globosum* and one from *Coprinopsis cinerea* was reported in an effort to characterize orphan PKSs [11]. These results show that yeast heterologous expression can provide an avenue for production of industrially relevant polyketides and for characterizing orphan polyketide synthase genes from non-model organisms. However, common for these examples is that the studies only aimed to produce the first stable intermediate in the respective biosynthetic pathway and have not included the heterologous expression of any tailoring enzymes responsible for modification of the synthase product. To test the feasibility of using yeast for production of complex polyketides we aimed to reconstruct a fungal polyketide pigment biosynthetic pathway, which leads to the formation of rubrofusarin (Figure 1). Rubrofusarin is an orange-brown pigment produced by a wide range of different fungal species, such as *Fusarium graminearum*, *Aspergillus niger*, *Aspergillus parasiticus* and *Ustilaginoidea virens* [12-14]. The biological activity of rubrofusarin is reported to include inhibition of human DNA topoisomerase II- α [15] and general antibiotic effects on *Mycobacterium tuberculosis* and various filamentous fungi [16]. Interestingly, the fate of this polyketide core scaffold differs in the particular fungal species due to their unique repertoires of tailoring enzymes, resulting in production of such different end products as rubrofusarin B, aurofusarin, nigerone, nigerasperone A, chaetochromin, ustilaginoidin and parasperone A [12,17].

Currently, the best described biosynthetic pathway, which includes rubrofusarin as an intermediate, is the aurofusarin pathway in *F. graminearum*. In this species, the first stable pathway intermediate on the path to rubrofusarin is the yellow naphthopyrone YWA1, which is synthesized by the condensation of one acetyl-CoA and six malonyl-CoA molecules, catalyzed by the type I iterative non-reducing PKS12 [18]. Similar reactions have been reported for *Aspergillus nidulans* [19]. The formed YWA1 is dehydrated by AurZ to form the orange pigment nor-

rubrofusarin, which is then O-methylated by AurJ to yield rubrofusarin (Figure 1) [12,18]. Polyketide biosynthetic pathways are typically encoded by gene clusters in fungi and mapping of the aurofusarin pathway in *F. graminearum* has therefore primarily been based on targeted deletion of the genes surrounding *PKS12* [12,18,20]. Thus, the possibility exists that genes located elsewhere in the genome might be directly involved in the formation of rubrofusarin. Hence, an appropriate bottom-up proof of the sole involvement of *PKS12*, *aurZ* and *aurJ* in rubrofusarin formation would be their heterologous expression in a phylogenetically distant organism.

Heterologous production in *S. cerevisiae* of other important compound classes with a complex biosynthetic origin, e.g. terpenoids [21] and plant type III polyketides [22], has already been well established. To provide a proof-of-concept that this approach is also a viable option for polyketide pathways with a fungal origin, and to prove the proposed biosynthetic pathway, we here present the stepwise construction of a rubrofusarin pathway in the heterologous host *S. cerevisiae*.

Results and discussion

Rubrofusarin has previously been described to have anti-fungal properties, and to avoid problems with toxic effects of intermediates or the end-product during strain construction, the open reading frame (ORF) encoding the wild type PKS12 was placed under control of the inducible *CUP1* promoter to allow control of expression onset and strength. Codon optimized ORFs for *aurZ*, *aurJ* and *npgA* were expressed using unique constitutive promoters and terminators (Table 1). The genes were inserted individually into different plasmids, derived from pRS413-6 [23], each carrying unique auxotrophic selection marker genes along with the unique promoter and terminator elements. This plasmid-based approach was chosen to allow for easy construction of any desired combination of genes in *S. cerevisiae* to study the sequential order of individual catalytic steps in the biosynthetic pathway (Table 1).

Heterologous production of YWA1

Expression of *PKS12* and *npgA* in *S. cerevisiae* resulted in the production of a yellow pigment which was not seen in the control strain carrying empty plasmids. In contrast to plates without added copper, yellow-colored




Figure 1 The rubrofusarin biosynthetic pathway. Biosynthetic pathway of rubrofusarin in *F. graminearum* established by targeted gene deletion experiments [12,18].

Table 1 *S. cerevisiae* strains used in the study

Strain	Plasmids carried	Relevant genotype
<i>S. cerevisiae</i> control	pRS416 (<i>URA3</i>)	-
	pRS413 (<i>HIS3</i>)	
	pRS414 (<i>TRP1</i>)	
	pRS415 (<i>LEU2</i>)	
<i>S. cerevisiae</i> <i>PKS12/npaA</i>	pRS416::P _{CUP1} - <i>PKS12</i> -t _{ADH1} , <i>PKS12</i> , <i>npaA</i>	
	pRS413::P _{PYK1} - <i>npaA</i> -t _{TEF1}	
	pRS414	
	pRS415	
<i>S. cerevisiae</i> <i>PKS12/npaA/aurZ</i>	pRS416::P _{CUP1} - <i>PKS12</i> -t _{ADH1} , <i>PKS12</i> , <i>npaA</i> , <i>aurZ</i>	
	pRS413::P _{PYK1} - <i>npaA</i> -t _{TEF1}	
	pRS414::P _{TEF1} - <i>aurZ</i> -t _{ENO2}	
	pRS415	
<i>S. cerevisiae</i> <i>PKS12/npaA/aurJ</i>	pRS416::P _{CUP1} - <i>PKS12</i> -t _{ADH1} , <i>PKS12</i> , <i>npaA</i> , <i>aurJ</i>	
	pRS413::P _{PYK1} - <i>npaA</i> -t _{TEF1}	
	pRS415::P _{GPD1} - <i>aurJ</i> -t _{CYC1}	
	pRS414	
<i>S. cerevisiae</i> <i>PKS12/npaA/aurZ/aurJ</i>	pRS416::P _{CUP1} - <i>PKS12</i> -t _{ADH1} , <i>PKS12</i> , <i>npaA</i> , <i>aurZ</i> , <i>aurJ</i>	
	pRS413::P _{PYK1} - <i>npaA</i> -t _{TEF1}	
	pRS414::P _{TEF1} - <i>aurZ</i> -t _{ENO2}	
	pRS415::P _{GPD1} - <i>aurJ</i> -t _{CYC1}	

S. cerevisiae strains that were analyzed for production of rubrofusarin and precursors. Final plasmids transformed into *S. cerevisiae* strains showing the vector backbone in which the constructed promoter (p) – ORF – terminator (t) gene cassettes were inserted. The auxotrophic marker gene is also shown. All plasmids were maintained by the low copy number sequences ARS/CEN.

pigment was visually observed after approx. 24 hours of cultivation on synthetic complete (SC) agar plates supplemented with copper. Following incubation for an additional 2-3 days, a yellow pigment became visible in the medium surrounding the *S. cerevisiae* colonies, indicating that the produced pigment was escaping from the cells. In SC liquid cultures a yellow color became visually observable after 2-3 days of cultivation (Figure 2). Chemical analysis, using LC-MS, of the growth medium from the induced *PKS12/npaA* strain revealed a new peak with a retention time (RT) of 5.3 min and a [M + H]⁺ of 277.07 m/z. Based on the exact mass, the compound was identified as the pigment YWA1 (Figure 3). The intensity of the peak was, however, relatively weak suggesting either a low production level, poor extraction or ionization of YWA1, or rapid degradation as was previously reported by Fujii and colleagues working with YWA1 in *A. oryzae* [24]. This compound is also known from *Aspergillus fumigatus* in which it is converted to 1,8-dihydroxynaphthalene (DHN) in multiple steps, including a chain-shortening reaction catalyzed by Ayl1 [25], and in *A. nidulans* where it is the precursor for the green conidial pigment [19].

Heterologous production of nor-rubrofusarin


Introduction of the dehydratase-encoding *aurZ* gene in the YWA1-producing strain resulted in colonies that visually appeared more intensely orange after 3 days of cultivation as compared to the parent YWA1-producing *PKS12/npaA* strain (Figure 2). The yellow pigment observed in the plate medium surrounding *S. cerevisiae* colonies of the *PKS12/npaA* strain was no longer visible, suggesting conversion of YWA1 into another orange pigment, which did not readily escape the cells (Figure 2).

Chemical analysis revealed a single new peak with a RT of 10.0 min and a [M + H]⁺ = 259.06 m/z which displayed a distinct UV absorption spectrum (see UV spectrum in Additional file 1). Based on a comparison to nor-rubrofusarin extracted from a nor-rubrofusarin-accumulating *F. graminearum* Δ *aurJ* reference strain, and by comparison to a previously published UV spectrum, the compound was identified as nor-rubrofusarin (Figure 3) [18]. Nor-rubrofusarin could be detected in both the cell pellet and growth medium, while YWA1 was still only detectable in the growth medium, further indicating that the observed change in colour is linked to the compounds of the identified peaks (Figure 4). A low level of nor-rubrofusarin was also detected in the cell pellet of the *PKS12/npaA* parent strain likely resulting from spontaneous conversion of YWA1. Spontaneous dehydration of YWA1 and YWA1-like compounds has previously been observed with purified compounds [18,26].

Heterologous production of rubrofusarin


Introduction of *aurJ*, encoding an O-methyltransferase, into the *S. cerevisiae* *PKS12/npaA/aurZ* strain changed the visual phenotype to appear less orange during growth on plates (strain *PKS12/npaA/aurZ/aurJ*, Figure 2). Chemical analysis showed that nor-rubrofusarin was no longer accumulating. Instead, a new compound accumulated which had a RT of 13.3 min, [M + H]⁺ of 273.07 m/z and a unique UV absorption spectrum (see UV spectrum in Additional file 1). The compound was identified as rubrofusarin by comparison to rubrofusarin extracted from a rubrofusarin-accumulating *F. graminearum* Δ *aurF* reference strain (Figure 3) and by comparison to the previously published UV spectrum for rubrofusarin [18]. The rubrofusarin could be detected both in the extracted cell pellet and growth medium. YWA1 was still measurable in the growth medium, however at a reduced level (Figure 4A).

Surprisingly, introduction of *aurJ* in the *PKS12/npaA* strain resulted in colonies that were more orange than the *PKS12/npaA* strain which solely accumulates YWA1 (Figure 2). Chemical analysis showed the production of rubrofusarin, evidenced by the appearance of a peak at the same RT and [M + H]⁺ = 273.07 m/z as in the *PKS12/*



npgA/aurZ/aurJ strain, but at an approx. six-fold lower level. These results are contrary to what has previously been observed in *F. graminearum* ΔaurZ mutants which only accumulate YWA1 even though they still express *aurJ* [18]. However, as we observed in the *PKS12/npgA* strain, low levels of YWA1 may undergo spontaneous conversion to nor-rubrofusarin, and subsequent methylation of this by AurJ may then explain

the formation of rubrofusarin without the presence of AurZ. The difference in rubrofusarin concentration observed between the *PKS12/npgA/aurZ/aurJ* and *PKS12/npgA/aurJ* strains shows that AurZ is responsible for catalyzing the specific dehydration of YWA1 to yield nor-rubrofusarin, as previously suggested [18]. It is noteworthy that no methylated YWA1 was observed in the *PKS12/npgA/aurJ* strain at $[\text{M} + \text{H}]^+ = 291.08 \text{ m/z}$,



suggesting that formation of rubrofusarin only proceeds through the nor-rubrofusarin intermediate, as previously proposed by Frandsen *et al* [12].

The presented heterologous production of rubrofusarin supports the biosynthetic pathway suggested by Frandsen *et al.* 2011 i.e. that *PKS12*, *AurZ*, and *AurJ* are stepwise responsible for formation of *YWA1*, nor-rubrofusarin, and rubrofusarin, respectively. In addition, it rules out the possibility that further, as yet unidentified enzymes are required for completion of the pathway.

The accumulated rubrofusarin from the strains *PKS12/npG/A/aurZ/aurJ* and *PKS12/npG/A/aurJ* was quantified by HPLC. Integration of rubrofusarin peaks from strain culture extracts showed a titer of 1.1 mg/L rubrofusarin produced by the *PKS12/npG/A/aurZ/aurJ* strain, while 0.19 mg/L rubrofusarin was measured in the strain extract, when *AurZ* was excluded from the pathway.

In the present study no optimization strategies for production were employed. However, several simple optimization strategies would be applicable, such as increasing the gene copy number, tuning the individual expression level of pathway enzymes, and increasing the acetyl-CoA and malonyl-CoA precursor pools in the cell. In addition, optimizing the fermentation conditions is likely to substantially increase the titer. Production of cell pigmentation would provide a simple and convenient read-out for successful optimization.

The structurally simpler polyketide 6-MSA has previously been heterologously produced in *S. cerevisiae* reaching a titer of 1.7 g/L [6]. In another study, the 6-MSA titer was increased 60% by increasing the malonyl-CoA supply, via overexpression of the acetyl-CoA carboxylase *ACC1*, reaching a final 554 mg/L with their particular strain background [8]. These results suggest that the potential for rubrofusarin production in yeast may be significantly higher than what we detected in this pathway reconstruction study.

***S. cerevisiae* as a heterologous host for fungal natural or synthetic polyketide pathways**

Simple eukaryotic hosts, such as the unicellular yeast *S. cerevisiae*, hold great potential for expression and study of heterologous biosynthetic pathways, including those of entire gene clusters typical of secondary metabolism of filamentous fungi. In synthetic biology terms, *S. cerevisiae* represents a more “minimal host” than filamentous fungi, due to the absence of any significant endogenous secondary metabolites [27], which could otherwise result in pleiotropic effects, complicating product formation and the ensuing characterization of the heterologous pathways. Hence, a yeast-based system for heterologous expression of fungal polyketide biosynthetic pathways is very well-suited for efficient characterization of the continuously increasing pool of orphan fungal polyketide




Figure 4 Detection of intermediates inside and outside of the cells. Detection of the rubrofusarin pathway intermediates in the constructed *S. cerevisiae* strains, shown by mean areas of the integrated peaks from extracted ion chromatograms for **A**) YWA1, **B**) Nor-rubrofusarin, and shown by mean titer (mg/L) calculated for **C**) Rubrofusarin. Measurements performed for the analyzed cell pellet (C) and growth medium (GM) of the constructed *S. cerevisiae* strains (see Table 1). Error bars indicate the std. deviation of each integrated peak (n = 3). Note that the mean areas are not comparable between different compounds.

gene clusters being revealed by genome sequencing projects [28,29].

Reconstruction of the rubrofusarin pathway further opens the door for heterologous production of a diversity of related, but novel polyketides in *S. cerevisiae*, for example by co-expressing combinations of tailoring enzymes such as p450s, glycosyl-, acyl- and prenyl-transferases etc. selected from other secondary metabolite pathways and gene clusters. The genes of such enzymes could efficiently be laid “on top” of the rubrofusarin pathway for example by random assembly and expression on yeast artificial chromosomes [22]. The readily observable phenotype of the developed rubrofusarin-producing strain may also serve as a tool for optimizing yeast for general polyketide production, e.g. via enhanced precursor supply or screening for optimized enzymes such as PPTases.

Conclusions

S. cerevisiae was engineered to produce the orange-brown polyketide pigment rubrofusarin, a common intermediate in a range of biosynthetic pathways found in different filamentous fungal species, by introduction of the three *F. graminearum* genes *PKS12*, *aurZ* and *aurJ* together with the PKS-activating PPTase gene *npgA*. A titer of 1.1 mg/L rubrofusarin was detected. The biosynthetic reconstruction further provides the evidence that *PKS12* is responsible for formation of the pigment YWA1, whereas *AurZ* catalyzes the formation of the pigment nor-rubrofusarin from YWA1. The results confirm the previously proposed biosynthetic pathway for the formation of rubrofusarin using a bottom-up approach and show that no other enzymes are necessary. The study also demonstrates that it is possible to produce complex fungal polyketide products in *S. cerevisiae*, which opens up for the industrial production of a diversity of active and commercially interesting polyketides in the future.

Materials and methods

Strains

Molecular cloning was performed using *Escherichia coli* XL1 Blue, *E. coli* XL10 Gold (Stratagene) and *E. coli* DH5 α (Invitrogen) competent cells. Final plasmids carried by strains in the chemical analysis are listed in Table 1. All *S. cerevisiae* strains constructed had the original genotype *mat α* , *trp1 Δ* , *lys2 Δ 0*, *leu2 Δ* , *his3 Δ 1*, *ura3 Δ 0*, *arg4::loxP* and were derived from *S. cerevisiae* BY4742 (Euroscarf). The constructed strains are listed in Table 1.

Enzymes and oligonucleotides

All restriction enzymes were purchased from New England Biolabs. Oligonucleotides for subcloning and mutagenesis of *PKS12* were synthesized by Microsynth AG, while

oligonucleotides for the USER friendly cloning were synthesized by Integrated DNA Technologies.

Design and synthesis of genes

A wild type *F. graminearum* PH1 *PKS12* ORF was constructed using USER friendly cloning [30] by the assembly of the six exons predicted by MIPS [31] [FGDB: FGSG_02324] into the recipient vector BGHA-P1 [32] with *in vivo* ligation in *E. coli*. All exon fragments were PCR amplified from *F. graminearum* PH1 genomic DNA using primers designed with the PHUSER web tool (see oligonucleotides in Additional file 1) [33]. An internal HindIII restriction site in the ORF was silently mutated by mixing approx. 200 ng plasmid harboring the gene with 10 nmol of the primer Mut1Fwd and 10 nmol Mut1Rev in an otherwise standard 50 μ L PCR mixture using iProof DNA polymerase and HF Buffer (BioRad) as the manufacturer specifies. The reaction was PCR thermocycled for 18 cycles followed by addition of 1 μ L DpnI and 3 hours of incubation at 37°C to degrade the dam methylated original (non-mutated) plasmid strands. 2 μ L reaction was transformed into 25 μ L *E. coli* XL10 Gold competent cells and plasmids isolated with QIAprep Miniprep (Qiagen). A correctly mutated plasmid was subsequently validated by the absence of HindIII digestion.

The *AurZ* [FGDB: FGSG_02326] and *AurJ* [FGDB: FGSG_02325] protein sequences from *F. graminearum* PH1 were used for synthesizing the two gene ORFs with verification by sequencing and proprietary “GeneGPS” codon optimization for *S. cerevisiae* by DNA2.0. The *A. fumigatus* *npgA* gene [GenBank: XM_750100.1] was synthesized by DNA2.0 with codon-optimization for *S. cerevisiae* and verification by sequencing.

Cloning of ORFs in pRS vectors

A set of pRS vectors [31] were used in which a unique *AscI* site had been inserted between the two *PvuII* sites, thus replacing the original multi-cloning site. All ORFs were subcloned with HindIII and SacII into vectors containing the respective, constitutive promoters and terminators (specified in Table 1) to form vectors with gene cassettes. The *PKS12* was provided with flanking HindIII and SacII restriction sites by PCR with iProof DNA polymerase using primers *PKS12Fwd* and *PKS12Rev* (see Additional file 1). All cassettes were subsequently cloned into the specified pRS plasmids (Table 1) with *AscI*. All ligations were performed using T4 DNA Ligase (New England Biolabs) and confirmed by sequencing (Microsynth). The pRS plasmids each harbour a unique auxotrophic marker gene (Table 1) and the ARS/CEN sequences for stable maintenance in *S. cerevisiae* at approx. one copy per cell, thus allowing strain-to-strain comparability. The *PKS12* ORF was controlled by the inducible *CUPI* promoter,

which is induced from its low basal expression level by addition of Cu^{2+} to the medium [34].

Transformation of *S. cerevisiae* strains

The *S. cerevisiae* strains (Table 1) were constructed by transformation with the specified plasmids (Table 1) according to the protocol described by Gietz & Schiestl, 2007 [35]. Transformants were plated on selective synthetic complete (SC) agar plates lacking the appropriate supplements uracil, histidine, tryptophan and leucine.

Cultivation of the *S. cerevisiae* strains

The *S. cerevisiae* strains were grown for characterization on 2% agar plates with selective SC medium plus 500 μM CuSO_4 for induction at 30°C. For chemical analysis, the strains were cultured by inoculating 5 mL selective SC medium from single colonies. These pre-cultures were incubated overnight at 30°C, with 160 rpm horizontal shaking. Main cultures of 10 mL selective SC medium, plus 300 μM CuSO_4 for induction, were inoculated from the pre-cultures to an $\text{OD}_{600} = 0.10$ in three replicates of each pre-cultured strain. The main cultures were incubated for 96 hours at 30°C with 160 rpm horizontal shaking.

Cultivation of reference *F. graminearum* strains

The reference *F. graminearum* strains ΔaurZ , ΔaurJ and ΔaurF were grown on Potato Dextrose Agar (Difco) plates at 20°C for 7 days.

Chemical extraction and LC-MS analysis

Cells and growth medium of the main cultures were separated by centrifugation. The cells were resuspended in 1 mL dH_2O and extracted by vigorous shaking with 5 mL extraction solvent (methanol:dichloromethane:ethyl acetate (1:2:3) + 1% formic acid). The mixtures were sonicated for 1 hour in an ultrasonic bath, centrifuged, filtered through filter paper and dried in a SpeedVac. Eight mL of the growth medium were extracted by vigorous shaking with 10 mL extraction solvent after which the organic phases were dried in a SpeedVac. Three 10 mm plugs were cut out of the mycelium of each reference *F. graminearum* strain. The plugs of each replicate were mashed in 10 mL ethyl acetate and shaken vigorously to facilitate extraction. The organic phase was recovered following centrifugation and filtering, at which point it was dried in a SpeedVac. The extractions from the *F. graminearum* reference strains were performed in triplicates.

All samples were reconstituted in 150 μL dimethyl sulfoxide. Five μL was injected and analyzed by LC-UV/ELSD on a Waters Acquity UPLC (Waters Corporation, Milford, MA, USA), running a gradient of mobile phases A (water + 0.1% formic acid) and B (acetonitrile + 0.1% formic acid), going from 1 to 100% B in 15 min, flow rate of 0.4 mL/min. The column was a Waters

Acquity BEH C18 (2.1 mm \times 100 mm \times 1.7 μm (Waters Corporation, Milford, MA, USA) temperature-controlled at 35°C. Liquid chromatography was coupled to a Bruker Daltonics microQTOF II (Bruker Daltonics, Bremen, Germany) using electrospray ionization source in positive mode: End plate Offset: -500 V, Capillary voltage: 4500 V, Nebulizer: 1.6 Bar, Dry Gas: 8.0 L/min, Dry temperature: 180°C. Extracted ion chromatograms for YWA1 (m/z 277.0707 \pm 0.01), nor-rubrofusarin (m/z 259.0601 \pm 0.01) and rubrofusarin (m/z 273.0753 \pm 0.01) were created using the Bruker DataAnalysis 4.0 software. The integration of the peaks was performed with the DataAnalysis Integrate functions (with default settings).

Quantification of rubrofusarin production

Cells and growth medium from 8 mL main culture were separated by centrifugation. The cell pellet was resuspended in 1 mL dH_2O and 5 mL ethyl acetate. The cell pellet solution was sonicated for 30 min. in an ultrasonic bath followed by vigorous shaking. The growth medium supernatant was shaken vigorously with 8 mL ethyl acetate. The ethyl acetate phases of growth medium and cell pellet were collected individually and allowed to evaporate in a fume hood. The extracted compounds of growth medium and cell pellet were reconstituted in 1.5 mL and 1.0 mL methanol, respectively. A rubrofusarin standard, purified from *F. graminearum*, was purchased from BioViotica Naturstoffe GmbH (Liestal, Switzerland) and reconstituted to 1 mg/mL in dimethyl sulfoxide. A standard curve was prepared by dissolving the rubrofusarin standard in methanol in a range from 4 $\mu\text{g/mL}$ to 10 $\mu\text{g/mL}$. Twenty μL of each standard concentration and the extracted samples were analyzed by HPLC using an Ascentis C18 column (particle size: 5 μm , length \times I.D.: 15 cm \times 4.6 mm) (Sigma-Aldrich) running a gradient of mobile phases C (10 mM ammonium formate pH 3.0) and D (acetonitrile) at a flow rate of 1 mL/min, starting at 25% D (0-3 min), going to 80% D (3-15 min), remaining at 80% D (15-16 min), going to 25% D (16-18 min) and remaining at 25% D (18-20 min).

Additional file

Additional file 1: Oligonucleotides, UV spectra and HPLC chromatograms.

Competing interests

PR, UM and RF declare no competing interests. MN is employed by Evolva SA.

Authors' contributions

PR and MN planned the experimental procedures of the study. PR carried out the experiments, except for cloning the exons of *PKS12*, which RF did. PR analyzed and interpreted the data and wrote the manuscript together with RF, MN and UM. All authors read and approved the final manuscript.

Acknowledgements

The authors wish to thank Evolva SA for supplying the laboratory facilities and materials including chemical gene synthesis. For helpful technical assistance and guidance we especially wish to thank Anders Hansson, Christophe Folly and Thiamo Boussemgoune, all employed at Evolva. Further, we wish to thank Sailesh Malla for helpful advice regarding compound quantification. RF's involvement was funded by the Danish Research Council for Independent Research — Technology and Production grant no. 09-069707 and Young Elite Researcher grant no. 09-076147.

Author details

¹Evolva SA, Duggingerstrasse 23, Reinach CH-4153, Switzerland. ²Søltofts Plads Building 223, Kgs Lyngby DK-2800, Denmark.

Received: 18 December 2012 Accepted: 1 April 2013

Published: 4 April 2013

References

- Weissman KJ, Leadlay PF: **Combinatorial biosynthesis of reduced polyketides.** *Nat Rev Microbiol* 2005, **3**:925–936.
- Tao K, Du L, Sun X, Cai M, Zhu T, Zhou X, Gu Q, Zhang Y: **Biosynthesis of aspergiolide A, a novel antitumor compound by a marine-derived fungus *Aspergillus glaucus* via the polyketide pathway.** *Tetrahedron Lett* 2009, **50**:1082–1085.
- Zhang YL, Ge HM, Zhao W, Dong H, Xu Q, Li SH, Li J, Zhang J, Song YC, Tan RX: **Unprecedented immunosuppressive polyketides from *Daldinia eschscholzii*, a Mantis-Associated Fungus.** *Angew Chem* 2008, **120**:5907–5910.
- Mapari SA, Meyer AS, Thrane U, Frisvad JC: **Identification of potentially safe promising fungal cell factories for the production of polyketide natural food colorants using chemotaxonomic rationale.** *Microb Cell Fact* 2009, **8**:24.
- Wang T, Lin T: **Monascus Rice Products.** *Adv Food Nutr Res* 2007, **53**:123–159.
- Kealey J, Liu L, Santi D, Betlach M, Barr P: **Production of a polyketide natural product in nonpolyketide-producing prokaryotic and eukaryotic hosts.** *Proc Natl Acad Sci USA* 1998, **95**:505–509.
- Wattanachaisaereekul S, Lantz AE, Nielsen ML, Andre S: **Optimization of Heterologous Production of the Polyketide 6-MSA in *Saccharomyces cerevisiae*.** *Biotechnology* 2007, **97**:893–900.
- Wattanachaisaereekul S, Lantz AE, Nielsen ML, Nielsen J: **Production of the polyketide 6-MSA in yeast engineered for increased malonyl-CoA supply.** *Metab Eng* 2008, **10**:246–254.
- Zhou H, Qiao K, Gao Z, Vederas JC, Tang Y: **Insights into radical biosynthesis via heterologous synthesis of intermediates and analogs.** *J Biol Chem* 2010, **285**:41412–41421.
- Zhou H, Gao Z, Qiao K, Wang J, Vederas JC, Tang Y: **A fungal ketoreductase domain that displays substrate-dependent stereospecificity.** *Nat Chem Biol* 2012, **8**:331–333.
- Ishiyuchi K, Nakazawa T, Ookuma T, Sugimoto S, Sato M, Tsunematsu Y, Ishikawa N, Noguchi H, Hotta K, Moriya H, Watanabe K: **Establishing a New Methodology for Genome Mining and Biosynthesis of Polyketides and Peptides through Yeast Molecular Genetics.** *Chem Bio Chem* 2012, **13**:846–854.
- Frandsen RJN, Nielsen NJ, Maolanon N, Sørensen JC, Olsson S, Nielsen J, Giese H: **The biosynthetic pathway for aurofusarin in *Fusarium graminearum* reveals a close link between the naphthoquinones and naphthopyrones.** *Mol Microbiol* 2006, **61**:1069–1080.
- Ghosal S, Biswas K, Chakrabati D: **Toxic naphtho-gamma-pyrones from *Aspergillus niger*.** *J Agric Food Chem* 1979, **27**:1347–1351.
- Shibata S, Ohta A, Ogihara Y: **The reactions of ustilaginoidin A.** *Bull. Chem. Pharm.* 1963.
- Branco A, Pinto AC, Braz-Filho R, Silva EF, Grynberg NF, Echevarria A: **Rubrofusarin, a natural polyketide as new human topoisomerase II- α ; inhibitor.** *Revista Brasileira de Farmacognosia* 2008, **18**:703–708.
- Graham JG, Zhang H, Pendland SL, Santarsiero BD, Mesecar AD, Cabieses F, Farnsworth NR: **Antimycobacterial Naphthopyrones from *Senna obliqua*.** *J Nat Prod* 2004, **67**:225–227.
- Brown D, Hauser F, Tommasi R: **Structural Elucidation of a Putative Conidial Pigment Intermediate in *Aspergillus parasiticus*.** *Tetrahedron Lett* 1993, **34**:5–8.
- Frandsen RJN, Schütt C, Lund BW, Staerk D, Nielsen J, Olsson S, Giese H: **Two novel classes of enzymes are required for the biosynthesis of aurofusarin in *Fusarium graminearum*.** *J Biol Chem* 2011, **286**:10419–10428.
- Watanabe A, Sankawa U, Mayorga E, Timberlake WE, Ebizuka Y: **Re-identification of *Aspergillus nidulans* wA Gene to Code for a Polyketide Synthase of Naphthopyrone.** *Tetrahedron Lett* 1999, **40**:91–94.
- Kim J, Han K, Jin J, Kim H, Kim J, Yun S, Lee Y: **Putative Polyketide Synthase and Laccase Genes for Biosynthesis of Aurofusarin in *Gibberella zeae*.** *Appl Environ Microbiol* 2005, **71**:1701–1708.
- Ukibe K, Hashida K, Yoshida N, Takagi H: **Metabolic engineering of *Saccharomyces cerevisiae* for astaxanthin production and oxidative stress tolerance.** *Appl Environ Microbiol* 2009, **75**:7205–7211.
- Naesby M, Nielsen SV, Nielsen CA, Green T, Tange TO, Simón E, Knechtle P, Hansson A, Schwab MS, Titz O, Folly C, Archila RE, Maver M, Van Sint Fiet S, Boussemgoune T, Janes M, SS Ka, Sonkar SP, Mitra PP, Benjamin VAK, Korrapati N, Suman I, Hansen EH, Thybo T, Goldsmith N, Sorensen AS: **Yeast artificial chromosomes employed for random assembly of biosynthetic pathways and production of diverse compounds in *Saccharomyces cerevisiae*.** *Microb Cell Fact* 2009, **8**:45.
- Sikorski R, Hieter P: **A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*.** *Genetics* 1989, **122**:19–27.
- Fujii I, Watanabe A, Sankawa U, Ebizuka Y: **Identification of Claisen cyclase domain in fungal polyketide synthase WA, a naphthopyrone synthase of *Aspergillus nidulans*.** *Chem Biol* 2001, **8**:189–197.
- Fujii I, Yasuoka Y, Tsai H-F, Chang YC, Kwon-Chung KJ, Ebizuka Y: **Hydrolytic polyketide shortening by *ayg1p*, a novel enzyme involved in fungal melanin biosynthesis.** *J Biol Chem* 2004, **279**:44613–44620.
- Chiang Y-M, Meyer KM, Praseuth M, Baker SE, Bruno KS, Wang CCC: **Characterization of a polyketide synthase in *Aspergillus niger* whose product is a precursor for both dihydroxynaphthalene (DHN) melanin and naphtho- γ -pyrone.** *Fungal genetics and biology: FG & B* 2011, **48**:430–437.
- Siddiqui MS, Thodey K, Trenchard I, Smolke CD: **Advancing Secondary Metabolite Biosynthesis in Yeast with Synthetic Biology Tools.** *FEMS Yeast Res* 2011, **12**:144–170.
- Chiang Y-M, Chang S-L, Oakley BR, Wang CCC: **Recent advances in awakening silent biosynthetic gene clusters and linking orphan clusters to natural products in microorganisms.** *Curr Opin Chem Biol* 2011, **15**:137–143.
- Bok JW, Hoffmeister D, Maggio-Hall La, Murillo R, Glasner JD, Keller NP: **Genomic mining for *Aspergillus* natural products.** *Chem Biol* 2006, **13**:31–37.
- Geu-Flores F, Nour-Eldin HH, Nielsen MT, Halkier BA: **USER fusion: a rapid and efficient method for simultaneous fusion and cloning of multiple PCR products.** *Nucleic Acids Res* 2007, **35**:e55.
- Wong P, Walter M, Lee W, Mannhaupt G, Münsterkötter M, Mewes H-W, Adam G, Güldener U: **FGDB: revisiting the genome annotation of the plant pathogen *Fusarium graminearum*.** *Nucleic Acids Res* 2011, **39**:D637–D639.
- Hansen BG, Salomonsen B, Nielsen MT, Nielsen JB, Hansen NB, Nielsen K, Regueira TB, Nielsen J, Patil KR, Mortensen UH: **Versatile enzyme expression and characterization system for *Aspergillus nidulans*, with the *Penicillium brevicompactum* polyketide synthase gene from the mycophenolic acid gene cluster as a test case.** *Appl Environ Microbiol* 2011, **77**:3044–3051.
- Olsen LR, Hansen NB, Bonde MT, Genee HJ, Holm DK, Carlsen S, Hansen BG, Patil KR, Mortensen UH, Wernersson R: **PHUSER (Primer Help for USER): a novel tool for USER fusion primer design.** *Nucleic Acids Res* 2011, **39**(Suppl 2):W61–W67.
- Labbé S, Thiele D: **Copper ion inducible and repressible promoter systems in yeast.** *Methods Enzymol* 1999, **306**:145–153.
- Gietz RD, Schiestl RH: **High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method.** *Nat Protoc* 2007, **2**:31–34.

doi:10.1186/1475-2859-12-31

Cite this article as: Rugbjerg et al.: Reconstruction of the biosynthetic pathway for the core fungal polyketide scaffold rubrofusarin in *Saccharomyces cerevisiae*. *Microbial Cell Factories* 2013 **12**:31.