

Technical University of Denmark



## Modelling the fate of ciprofloxacin in activated sludge systems - The relevance of the sorption process

Polesel, Fabio; Lehnberg, Kai; Dott, Wolfgang; Trapp, Stefan; Thomas, Kevin H.; Plósz, Benedek G.

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Polesel, F., Lehnberg, K., Dott, W., Trapp, S., Thomas, K. H., & Plósz, B. (2013). Modelling the fate of ciprofloxacin in activated sludge systems - The relevance of the sorption process. Abstract from 14th EuCheMS International Conference on Chemistry and the Environment (ICCE 2013), Barcelona, Spain.

**DTU Library**  
Technical Information Center of Denmark

---

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Modelling the fate of ciprofloxacin in activated sludge systems – The relevance of the sorption process

Fabio Polese<sup>1</sup>, Kai Lehnberg<sup>2</sup>, Wolfgang Dott<sup>2</sup>, Stefan Trapp<sup>1</sup>, Kevin H. Thomas<sup>3</sup>, Benedek Gy. Plósz<sup>1</sup>

(1) Department of Environmental Engineering, Technical University of Denmark, Kgs. Lyngby, Denmark

(2) Institute of Hygiene and Environmental Medicine, RWTH Aachen University, Aachen, Germany.

(3) Department of Ecotoxicology and Risk Assessment, Norwegian Institute for Water Research (NIVA), Oslo, Norway.

[fabp@env.dtu.com](mailto:fabp@env.dtu.com)

The sorption process can impact the removal of specific pharmaceuticals in municipal wastewater treatment plants (WWTPs). Ionic interactions (e.g., pH-driven equilibria and complexation), rather than hydrophobic interactions, are known to affect the sorption of zwitterionic pharmaceuticals. In a previous study [1], a daily systematic reduction of ciprofloxacin removal in a full-scale WWTP (Bekkelaget, Norway) was associated to deteriorated sorption. Therefore, in this study we further investigated the sorption of ciprofloxacin onto activated sludge at laboratory- and full-scale. Targeted batch experiments were performed to estimate sorption model parameters using Freundlich isotherms under specific pH and iron salt dosing (used for chemical phosphorus removal) conditions. We used the previously tested activated sludge framework model for xenobiotic trace chemicals (ASM-X) to assess the fate of ciprofloxacin in a full-scale activated sludge system. Sorption was described by linear kinetics and, in an extended version of ASM-X, using a Freundlich-based submodel. In the latter case, Freundlich parameter values estimated from the batch experiments were used for model calibration. The prediction accuracy was statistically evaluated in the two cases by comparing the model output with measured data.

Batch experiments showed that maximum sorption capacity occurred at pH=7.4, corresponding to the isoelectric point of ciprofloxacin. A pH increase resulted in a significant reduction of sorption capacity as compared to the effect of the pH decrease applied in the experiment. Additionally, iron salt dosing was found to enhance sorption under both aerobic and anoxic conditions. Using the extended ASM-X model, results obtained in scenario simulations – based on the batch experimental Freundlich parameters – suggest that pH conditions, rather than reduced salt dosing, can be responsible for the decrease of ciprofloxacin sorption in the full-scale WWTP. The most accurate predictions were obtained for Freundlich parameter values of  $K=0.01$  ( $\mu\text{g}^{(1-1/n)} \text{L}^{1/n} \text{mg}^{-1}$ ) and  $1/n=1.33$ . A pH increase was therefore estimated to cause reduced sorption in the anoxic and the aerobic reactors, possibly being a consequence of the lower sorption extent exhibited by the anionic ciprofloxacin species. Comparable prediction accuracy was obtained using linear sorption. A 20-fold decrease of the anoxic and aerobic  $K_D$  values (1.1 and  $0.42 \text{ L gX}_{SS}^{-1}$  under normal conditions, respectively) was estimated in the time interval when deteriorated sorption was hypothesized.

[1] Plósz, B.G.; Leknes, H.; Thomas, K.V. (2010a). *Impacts of competitive inhibition, parent compound formation and partitioning behaviour on antibiotic micro-pollutants removal in activated sludge*. Environmental Science & Technology, 44 (2), 734–742.