



Open Research Online

The Open University's repository of research publications and other research outputs

Spore chemistry as a proxy for UV-B flux

Conference or Workshop Item

How to cite:

Watson, Jonathan; Sephton, Mark; Sephton, Sarah; Beerling, David; Self, Stephen; Gilmour, Iain and Wellman, Charles (2005). Spore chemistry as a proxy for UV-B flux. In: 22nd International Meeting on Organic Geochemistry, 12-16 Sep 2005, Seville, Spain.

For guidance on citations see [FAQs](#).

© [not recorded]

Version: [not recorded]

Link(s) to article on publisher's website:

<http://linkinghub.elsevier.com/retrieve/pii/S0146638006002397>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Spore chemistry as a proxy for UV-B flux

Jonathan S. Watson¹, Mark A. Sephton^{1,a}, Sarah V. Sephton², David J. Beerling³, Stephen Self², Iain Gilmour¹, Charles H. Wellman³

¹Planetary and Space Sciences Research Institute, Open University, Milton Keynes, MK7 6AA, UK

²Department of Earth Sciences, Open University, Milton Keynes, Buckinghamshire, MK7 6AA, UK

³Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

^aPresent address: Department of Earth Science and Engineering, Royal School of Mines, Imperial College, London, SW7 2BP, UK

Placing current increased terrestrial UV-B fluxes, due to the seasonal depletion of the stratospheric ozone layer, in a historical context is difficult due to a lack of long-term (century or more) instrumental records and necessitates developing proxy indicators. One promising line of enquiry derives from the response of plants to increased near-surface solar fluxes of harmful ultraviolet radiation in the 280 – 315 nm wavelength (UV-B) (Rozema et al. 2001, 2002). Plants exposed to increased UV-B radiation typically experience a number of detrimental effects, including damage to proteins, membrane lipids and DNA. To reduce this damage, many plants, animals and microbes accumulate UV-B protecting pigments (Cockell & Knowland 1999; Rozema et al. 2001, 2002).

Here, we evaluate the potential of a promising candidate for such a proxy, which is based on changes in the chemical composition of spores in response to variations in near-surface UV-B fluxes, in a field setting. We obtained spores from five populations of the tropical lycopsid *Lycopodium cernuum* growing across an altitudinal gradient (650-1981 m a.s.l.) in S.E. Asia with the assumption that they experienced a range of UV-B radiation doses. Spores from each population were analyzed for UV-B protecting compounds using micro-Fourier transform infrared spectroscopy (micro-FTIR) and thermochemolysis-GC-MS.

The data reveal the presence of various functional groups associated with UV-B protecting pigments including OH, C=O and C=C. Thermochemolysis and subsequent pyrolysis liberated UV-B pigments (ferulic and *para*-coumaric acid) from the spores. All of the aromatic compounds liberated from spores by thermochemolysis and pyrolysis were active in UV-B protection. We show systematic increases in micro-FTIR aromatic absorption

(1520 cm^{-1}) and olefinic or aromatic absorption (829 cm^{-1}) with altitude that reflect a chemical response to higher UV-B flux. Our results indicate that detailed chemical analyses of historical spore samples could provide a proxy for stratospheric O_3 layer variability and UV-B flux over historical (century to millennia) timescales.




Figure 1: Total ion chromatogram of products from a typical *Lycopodium cernuum* sample liberated at 300 °C in the presence of TMAH

REFERENCES

Cockell, C. S., Knowland, J., 1999. Ultraviolet radiation screening compounds. *Biological Reviews*. 74, 311-345.

Rozema, J., Broekman, R.A., Blokker, P., Meijkamp, B.B., de Bakker, N., van de Staaij, J., van Beem, A., Ariese, F., Kars, S.M., 2001. UV-B absorbance and UV-B absorbing compounds (para-coumaric acid) in pollen and sporopollenin: the perspective to track historic UV-B levels. *Journal of Photochemistry and Photobiology B* 62, 108-117.

Rozema, J., van Geel, B., Björn L.O., Lean, J., Madronich, S., 2002. Towards solving the UV puzzle. *Science* 296, 1621-1622.