



University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): Shengxing Zheng, Qian Wang, Qian He, Xiaorong Song, Duyun Ye, Fang Gao, Shengwei Jin, and QingQuan Lian
Article Title: Novel Biphasic Role of LipoxinA4 on Expression of Cyclooxygenase-2 in Lipopolysaccharide-Stimulated Lung Fibroblasts
Year of publication: 2011

Link to published article: <http://dx.doi.org/10.1155/2011/745340>
Publisher statement: Open Access authors retain the copyrights of their papers, and all open access articles are distributed under the terms of the Creative Commons Attribution license, which permits unrestricted use, distribution and reproduction in any medium, provided that the original work is properly cited.

Zheng, S. et al. (2011). Novel Biphasic Role of LipoxinA4 on Expression of Cyclooxygenase-2 in Lipopolysaccharide-Stimulated Lung Fibroblasts. *Mediators of Inflammation*, Article ID 745340.

Research Article

Novel Biphasic Role of LipoxinA₄ on Expression of Cyclooxygenase-2 in Lipopolysaccharide-Stimulated Lung Fibroblasts

Shengxing Zheng,¹ Qian Wang,¹ Qian He,¹ Xiaorong Song,¹ Duyun Ye,² Fang Gao,^{3,4} Shengwei Jin,¹ and QingQuan Lian¹

¹ Department of Anaesthesia, Second Affiliated Hospital of WenZhou Medical College, Zhejiang 325003, China

² Department of Pathophysiology, Tongji Medical College, Huazhong University of Science and Technology, 13 Hangkong Road, Wuhan, Hubei Province 430030, China

³ Warwick Medical School, University of Warwick, Coventry CV4 7AL, UK

⁴ Academic Department of Critical Care, Resuscitation, Anaesthesia, and Pain, Heart of England NHS Foundation Trust, Birmingham B9 5SS, UK

Correspondence should be addressed to

Shengwei Jin, jinshengwei69@yahoo.com.cn and QingQuan Lian, lianqingquan@yahoo.com.cn

Received 15 February 2011; Accepted 3 May 2011

Academic Editor: Donna-Marie McCafferty

Copyright © 2011 Shengxing Zheng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fibroblasts are important to host defence and immunity, can also as initiators of inflammation as well. As the endogenous “braking signal”, Lipoxins can regulate anti-inflammation and the resolution of inflammation. We investigated the effect of lipoxinA₄ on the expression of cyclooxygenase-2 in lipopolysaccharide-stimulated lung fibroblasts. We demonstrated that the expression of cyclooxygenase-2 protein was significantly increased and peaked initially at 6 hours, with a second increase, with maximal levels occurring 24 hours after lipopolysaccharide challenge. ProstaglandinE₂ levels also peaked at 6 hours, and prostaglandinD₂ levels were increased at both 6 and 24 hours. Exogenous lipoxinA₄ inhibited the first peak of cyclooxygenase-2 expression as well as the production of prostaglandinE₂ induced by lipopolysaccharide in a dose-dependent manner. In contrast, exogenous lipoxinA₄ increased the second peak of cyclooxygenase-2 expression as well as the production of prostaglandinD₂ induced by lipopolysaccharide in a dose-dependent manner. LipoxinA₄ receptor mRNA expression was markedly stimulated by lipopolysaccharide but inhibited by lipoxinA₄. We present evidence for a novel biphasic role of lipoxinA₄ on the expression of cyclooxygenase-2 in lipopolysaccharide-stimulated lung fibroblasts, whereby LXA₄ has an anti-inflammatory and proresolving activity in lung fibroblasts following LPS stimulation.

1. Introduction

The acute inflammatory reaction in the lung is a complex response but is usually self-limiting and resolves. Traditionally, immune effector cells such as lymphocytes and macrophages have been considered to have a fundamental role in the development of inflammation. Traditionally, fibroblasts have only been considered as a structural element. Recent studies, however, demonstrate that pulmonary fibroblasts, far from being merely bystander cells, are important to host defence but may also promote lung injury.

Recent evidence has shown that fibroblasts can produce proinflammatory cytokines and prostaglandins (PGs) and can act as initiators of inflammation as well as regulators of immunity [1–3]. When activated, fibroblasts are capable of producing inflammatory mediators, including interleukin-8 (IL-8), monocyte chemoattractant protein-1, express cyclooxygenase-2 (COX-2), with the resultant release of proinflammatory PGs such as prostaglandinE₂ (PGE₂) [3]. Moreover, fibroblasts can be directly activated by exposure to lipopolysaccharide (LPS) [4]. Gram-negative bacteria can be responsible for the failure of early treatment and

significantly increased morbidity and mortality in patients with pneumonia [5, 6]. LPS, as a major constituent of Gram-negative bacterial cell walls, is recognized by the innate immune system of cells, such as resident tissue fibroblasts [7]. Specifically, Toll receptor expression on these fibroblasts may be able to direct the tissues' response to injury to promote inflammation resolution.

Cyclooxygenase is a key enzyme that catalyzes the conversion of arachidonic acid to prostaglandin [8]. There are two cyclooxygenase isoforms, COX-1 and COX-2. COX-1 is produced constitutively in most cell types, whereas COX-2 is inducible [8]. Prostaglandins are lipid mediators synthesized from arachidonic acid by the actions of COX enzymes [9]. They can be secreted by resident fibroblasts, as well as inflammatory cells, in response to TNF- α , IL-1 β , or LPS [10, 11]. Prostaglandins also contribute to the signs and symptoms of inflammation [12]. ProstaglandinE₂, the main PG produced during inflammatory response, is a proinflammatory lipid mediator of inflammation and participates in initiation of inflammation [13]. Previous studies suggest PGD₂, as a proresolution mediator, also actively contributes to the resolution of tissue injury and inflammation [14].

Lipoxins (LXs) are trihydroxytetraene-containing eicosanoids mainly formed through transcellular biosynthesis involving either 5- and 15-lipoxygenases (LOXs) or 5- and 12-lipoxygenases (LOXs) as well as COX-2 in respiratory tissues [15, 16]. Lipoxins were the first proresolving mediators to be recognized; they not only have anti-inflammatory properties, but also promote the resolution of inflammation [17]. Lipoxins have been described as the endogenous "braking signal" for inflammation [18–20]. Lipoxins have been extensively studied in asthma [21, 22], cystic fibrosis [23–25], and in various infections. These studies have highlighted LXs as potential novel therapeutic agent for the treatment of inflammatory disease.

A recent study reported that injured bronchial epithelial cells upregulated lipoxinA₄ (LXA₄) receptor (LXA₄R) in a COX-2-dependent manner to promote LXA₄-mediated resolution of airway inflammation [26]. In response to acid injury, epithelial cells rapidly increased COX-2 and PGE₂ expression [26]. The COX-2 enzyme has also been implicated as an important mediator of pulmonary fibrosis, with COX-2^{-/-} mice having increased fibrotic lung responses [27]. More recently, Medeiros et al. reported that LXA₄ also repressed the expression and activity of COX-2 on endotoxin-induced uveitis (EIU) in rats [28]. Moreover, LXA₄ has also been shown to inhibit connective tissue growth factor- (CTGF) stimulated proliferation of human lung fibroblasts [25], and fibroblasts directly simulated by LPS are capable of producing COX-2 and PGE₂ [3, 4]. However, the time course of COX-2 expression in lung fibroblasts stimulated by LPS and the effect of LXA₄ on expression of COX-2 PGE₂ and PGD₂ remain unclear.

In this study, we examined the expression of COX-2 and the production of PGE₂ and PGD₂ in lung fibroblasts after LPS challenge. Additionally, we also investigated the effect of LXA₄ on the expression of COX-2 and the production of PGE₂ and PGD₂. Finally, we investigated the effect of LXA₄

or LPS on LXA₄R mRNA expression in lung fibroblasts; we present evidence for a novel biphasic role of LXA₄ on expression of COX-2 in LPS-stimulated lung fibroblasts.

2. Materials and Methods

2.1. Materials. LipoxinA₄, from Cayman Chemical Company, was stored at -80°C until being diluted in serum-free culture medium immediately before use. Lipopolysaccharide (LPS; E. coli serotype 055 : B5) was purchased from Sigma. DMEM, FCS, Trypsin EDTA, and enzyme-free cell dissociation buffer were purchased from Gibco. Penicillin and streptomycin in saline citrate buffer were from Invitrogen. Hoechst 33258 was obtained from Novus. Anti-CD31, anti-Vimentin and anti-COX-2 were purchased from Abcam. Anti-Cytokeratin-8 and anti-F4/80 were purchased from Santa Cruz.

2.2. Cell Culture. Rat pulmonary fibroblasts were isolated from Sprague-Dawley rats as previously described [29]. Lung tissue was cut into <1-mm³ pieces and dissociated in Hanks buffered saline solution (HBSS) containing 0.25% trypsin at 37°C for 1.5 min. Trypsin was inhibited by DMEM with 15% fetal calf serum (FCS) and dissociated tissue centrifuged at 1000 g for 5 minutes at 4°C. The dissociated tissue pieces were placed into a culture plate with DMEM containing 15% FCS and left to allow fibroblast outgrowth. After fibroblasts had grown out from the tissues, usually 2–3 days, the remaining tissue was removed by aspiration, and the cells were allowed to reach confluence. Confluent fibroblasts were then passaged with a split ratio of 1 : 2 by trypsin treatment and used for the experiments at passages 3–5. The purity of fibroblast cultures were consistently over 99% as established morphologically by their typical spindle shape and characteristics and by expression of the fibroblast marker vimentin and negative expression of endothelial (CD31), macrophage (F4/80), and epithelial (cytokeratin-8) cell markers.

For all experiments, cells were subcultured into six-well plates and maintained until subconfluence (80%), confluent cells (100%) were serum deprived for 24 hours with low-serum medium (DMEM supplemented with 0.1% FCS) prior to the addition of LPS and/or LXA₄. The cells were then incubated with LPS (1 μ g/mL) for 6, 12, 24, 48, and 72 hours. For LXA₄ experiments, the cells were incubated in the low-serum medium containing 1 μ g/ml LPS in the presence or absence of 100, 200, or 400 nmol/mL of LXA₄ for 6 or 24 hours.

2.3. Haematoxylin and Eosin (H&E) Staining. Fibroblasts grow in culture on coverslips were fixed with 4% paraformaldehyde in PBS for 10 min, stained with Haematoxylin (BDH, Lutterworth, UK) for 10 minutes, incubated in Scott's tap water (tap water with a few drops of 1 M sodium hydroxide) for 5 min and stained with alcoholic Eosin solution for 5 min. They were washed by immersion in tap water for 2 minutes after every step. Cells were then differentiated by immersion in 0.1% hydrochloric acid-ethanol each for 30 s and mounted by inversion onto glass slides dotted with Gel/Mount. Images were taken by an

inverted microscope (IX70, Olympus America, Inc., Melville, NY, USA) with a 1.40NA 60 × objective sets. Image size was 2560 × 1920.

2.4. Indirect-Immunofluorescence. Fibroblasts were grown to approximately 70% confluence on poly-D-lysine-coated glass coverslips in 24-well plates, fixed with 4% paraformaldehyde in PBS for 10 minutes, rinsed three times with PBS, and permeabilized by 0.2% Triton X-100/PBS for 2 minutes followed by 0.5% Triton X-100 (Pierce) in PBS for 10 minutes. Nonspecific binding of antibodies was prevented by the addition of 5% bovine serum albumin in PBS for 30 minutes at 37°C. The samples were then incubated overnight at 4°C with antivimentin (1:100), anti-CD31 (1:100), anti-F4/80 (1:100) or anti-Cytokeratin-8 (1:200) in 2% BSA/PBS. Following three PBS washes, cells were incubated for 2 hours with fluorescein-conjugated immunopure goat anti-mouse IgG (H+L) or goat antirabbit IgG (H+L) (1:200) respectively, in 5% BSA/PBS at room temperature. After washing three times with PBS, cell nuclei were counter stained with Hoechst (1:1000) for 15 minutes, followed by three PBS washes. Cells were then mounted by inversion onto glass slides dotted with Gel/Mount. Images were taken by an inverted microscope (IX70, Olympus America, Inc., Melville, NY) with a 1.40NA 60 × objective and FITC, rhodamine and Cy5 filter sets. Image size was 2560 × 1920.

2.5. PGE₂ and PGD₂ Protein Expression. Fibroblast supernatants were collected following treatments, centrifuged (1500 g, 5 minutes), aliquoted, and stored at -80°C. PGE₂ and PGD₂ protein expression was measured by ELISA according to the manufacturer's instructions (R&D systems). Assays were run in triplicate and repeated twice.

2.6. COX-2 Protein Expression. Fibroblasts were lysed and homogenized in 200 μL of cold lysis buffer (150 mM NaCl, 1% Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 50 mmol/L Tris-HCl (pH 7.2), 0.2 mM sodium vanadate, 1% phenylmethylsulfonyl fluoride, and 0.2% aprotinin). Samples were incubated on ice for 20 minutes and then centrifuged at 12,000 rpm for 10 minutes. Protein concentrations of the supernatants were determined by using a BCA protein assay (Pierce). Proteins were separated in 10% SDS polyacrylamide gels and transferred onto nitrocellulose membranes. Expression of COX-2 was determined using primary rabbit anti-COX-2 antibody (1:750) and secondary horseradish peroxidase-conjugated goat antirabbit IgG. Protein expression of β-actin served as a loading control. The bound antibody was detected by enhanced chemiluminescence on an X-ray film.

2.7. RNA Isolation, Reverse Transcription and PCR. Total RNA was extracted using TriZol reagent (Life Technologies) followed by phenol-chloroform extraction and ethanol precipitation (Fisher Scientific). RNA purity was checked by spectrophotometry, and RNA integrity was confirmed by visualization of 28 S and 18 S bands on an agarose gel. 1 μg of RNA was reverse transcribed using avian myeloblastosis

virus reverse transcriptase (Promega). PCR analysis was performed with the following sets of primers: for rat LXA₄ receptor 5'-TGTTGGGCCCTGGATTTTAGC-3' (sense) and 5'-TGTTACCCCAGGATGCGAAGTT-3' (antisense), amplifying a 116-bp fragment [30] and for β-actin, used as an internal control 5'-AACAGTCCGCCTAGAAGCAC-3' (sense) and 5'-CGTTGACATCCGTAAGACC-3' (antisense), generating a 281-bp fragment. PCR for the rat LXA₄ receptor consisted of 35 repetitive cycles of predenaturing at 95°C for 4 minutes, denaturing at 94°C for 30 seconds, annealing at 59°C for 40 seconds, extension at 72°C for 40 seconds, and a final extension at 72°C for 5 minutes. For r-actin an annealing temperature of 56°C was used. Amplified cDNA was separated on a 1.6% agarose gel and visualized using ethidium bromide. Semiquantitative analysis was performed using UVP-gel densitometry (SanGabriel, Calif. USA).

3. Results

3.1. Purification and Identification of Primary Lung Fibroblasts. Untreated fibroblast were stained with hematoxylin and eosin for conventional morphological evaluation under light microscope (Nikon eclipse 90i, Tokyo, Japan) (Figure 1(a)) or stained by indirect immunofluorescence for Vimentin, CD31, Cytokeratin-8 and F4/80 expressions (Figure 1(b)). Vimentin was used as marker of fibroblast cells, CD31 as marker of endothelial cells [31], F4/80 as a surface marker of macrophages [32, 33], and Cytokeratin-8, as a marker of epithelial cells [34]. We observed only cells with fibroblast morphology which stained only for Vimentin, therefore, only purified fibroblasts were cultured.

3.2. The Effect of LPS on COX-2, PGE₂ and PGD₂ Expression in Lung Fibroblasts. To determine the dynamic expression of COX-2 in rat lung fibroblasts, our isolated fibroblasts were incubated with LPS (1 μg/mL) for 6, 12, 24, 48, and 72 hours. The expression of COX-2 protein was significantly increased and peaked initially 6 hours after LPS stimulation, with maximal levels occurring at 24 hours (Figure 2(a)). In contrast, PGE₂ levels were increased only at 6 hours (Figure 2(b)), with the precursor of prostaglandin J series, PGD₂ [35], levels increased at both 6 and 24 hours, (Figure 2(c)).

3.3. The Effect of LXA₄ on LPS-Induced Expression of COX-2 Protein Expression and PGE₂, and PGD₂ Production at 6 Hours in Primary Lung Fibroblasts. To determine whether exogenous LXA₄ modulates COX-2 expression after LPS stimulation, we reassessed COX-2 protein at 6 hours with various concentrations of LXA₄ treatment in our isolated lung fibroblasts. Using LXA₄ at 100, 200, or 400 nmol/ml we observed inhibition of COX-2 protein expression in a dose-dependent manner (Figure 3(a)). Moreover, after cells were incubated with LXA₄ for 6 hours, PGE₂ and PGD₂ protein levels in the supernatant were measured by ELISA (Figure 3(b) and 3(c), resp.). PGE₂ secretion was inhibited by LXA₄ in a dose-dependent manner, decreasing




FIGURE 1: Purification and identification of primary lung fibroblasts. (a) Untreated fibroblasts isolated from rat lungs were stained with hematoxylin and eosin for conventional morphological evaluation under light microscope (Nikon eclipse 90i, Tokyo, Japan). Primary cultures (A1), still contained some non-fibroblasts cells. At passage 3 (A2), fibroblast purity was consistently over 99% as established morphologically by their typical spindle shape and characteristics. (b) Cultures were stained by indirect-immunofluorescence assay for Vimentin, CD31, cytokeratin-8, or F40/80. Cultures contained only Vimentin positive cells indicative of fibroblasts.

from 411.734 ± 1.364 pg/mL in 0 nM LXA₄ treated cells to 307.075 ± 2.151 pg/mL in 100 nM LXA₄-treated fibroblasts and then further still to 108.089 ± 4.851 pg/mL in 400 nM LXA₄-treated fibroblasts ($P < 0.05$). In contrast although PGD₂ levels, increased with LXA₄ treatment, this was not observed to be dose dependent.

3.4. The Effect of LXA₄ on LPS-Induced Expression of COX-2 Protein Expression and PGE₂ and PGD₂ Production at 24 Hours in Primary Lung Fibroblasts. To determine whether treatment with exogenous LXA₄ affected the secondary increase of COX-2 expression after LPS stimulation, we also reassessed COX-2 protein at 24 hours after various concentrations of lipoxinA₄ treatment. Using LXA₄ at 100, 200, or 400 nmol/mL, we observed an increase in COX-2 protein expression in a dose-dependent manner (Figure 4(a)). We also measure secretion of PGE₂ and PGD₂ following LPS and LXA₄ treatments (Figures 4(b) and 4(c), resp.).

Interestingly, in contrast to our result at 6 hr LPS treatment although levels of PGE₂ increased with LXA₄, it was not dose-dependent. Furthermore, PGD₂ secretion following this treatment regime was enhanced by LXA₄ in a dose-dependent manner, increasing from 367.170 ± 4.773 pg/mL in 0 nM LXA₄-treated cells to 417.916 ± 3.251 pg/mL following 100 nM LXA₄-treated fibroblasts and 584.307 ± 15.478 pg/mL in 400 nM LXA₄-treated fibroblasts ($P < 0.05$).

3.5. LipoxinA₄ Receptor is Expressed in Rat Lung Fibroblasts and Upregulated by LPS. LipoxinA₄ interactions with its receptor, LXA₄R, play a significant role in regulating leukocyte functions [36]. Therefore, we tested whether LXA₄R mRNA expression altered following LPS and LXA₄ treatment in rat lung fibroblasts by using semiquantitative RT-PCR. A single band corresponding to LXA₄R mRNA expression was amplified (Figure 5, lane 1), which when analysed by densitometry was markedly stimulated by LPS treatment (Figure 5, lane 2). Interestingly, we observed that cotreatment with LPS and LXA₄ reduced expression of LXA₄R back to that seen in untreated controls (Figure 5, lane 3).

4. Discussion

Acute lung injury (ALI)/ARDS is an inflammatory lung disease with high mortality [36–39]. Treatment of inflammatory diseases today is largely based on interrupting the synthesis or action of mediators that also decrease the host's ability to successfully deal with infection, given that the innate inflammatory response is a beneficial defensive event [40]. Recently, resolution of acute inflammation was shown to be an active, rather than a passive process, and endogenous chemical mediators play key roles in its programmed resolution and returning to homeostasis [41]. Among them, lipoxins and aspirin-triggered lipoxins evoke bioactions in a range of physiologic and pathophysiologic processes and serve as endogenous lipid/chemical mediators that stop neutrophilic infiltration and initiate resolution [17]. Development of strategies that promote the resolution of inflammation is a novel therapeutic measure to attenuate inflammatory lung injury.

In a previous study, we clearly demonstrated that post-treatment with lipoxinA₄ (LXA₄) significantly reduces LPS-induced ALI in mice [42]. Lipoxin also promotes gradual resolution of fibrosis in lung [43]. In addition, LXA₄ repressed the expression and the activity of COX-2 on endotoxin-induced uveitis in rats [28]. In carrageenin-induced pleurisy in rats, COX-2 protein expression peaked initially at 2 hours, and at 48 hours, there was a second increase in COX-2 expression in inflammatory cells separated from the inflammatory exudates [44]. Taking these data together, our purpose was to find out whether COX-2 expression in lung fibroblasts stimulated by LPS also has two peaks and, if so, how LXA₄ affect the expression of COX-2 and the production of prostaglandins, specifically PGE₂ and PGD₂.

Our data clearly demonstrated that the expression of COX-2 protein was significantly increased and peaked initially 6 hours after LPS stimulation in lung fibroblasts.




FIGURE 2: The effect of LPS on COX-2, PGE₂, and PGD₂ expression in lung fibroblasts. (a) Rat lung fibroblasts were incubated with LPS (1 μg/mL) for 6, 12, 24, 48, and 72 hours. The expression of COX-2 protein was assessed by western blot and analysed by densitometry compared to β-actin expression. COX-2 expression peaked initially at 6 hours, and then with maximal levels at 24 hours after-LPS treatment (**P* < 0.05 versus non-LPS group #*P* < 0.05 versus 12, 48, 72 hours groups). (b) Supernatants were collected after LPS (1 μg/mL) treatment for 6, 12, 24, 48 and 72 hours. PGE₂ protein was measured by ELISA. Data are expressed as mean ± SE for each group. (**P* < 0.05 versus non-LPS group; &*P* < 0.01 versus 12, 24, 48, and 72 hours groups; #*P* < 0.05 versus 6, 24, 48, and 72 hours groups). (c): Supernatants were collected after LPS (1 μg/mL) treatment for 6, 12, 24, 48 and 72 hours. PGD₂ protein was measured by ELISA. Data are expressed as mean ± SE for each group. (**P* < 0.05 versus non-LPS group; #*P* < 0.05 versus 12, 48, 72 hours groups).

This was also associated with maximal PGE₂ synthesis. However, following 24 hours of LPS stimulation, there was a second increase in COX-2 expression, this time associated with maximal PGD₂ synthesis. Thus, as inflammation progresses into resolution, PGE₂ synthesis declines, giving way to a prominence of COX-2-derived PGD₂, both of which play important roles in mediating resolution. This data indicates that COX-2 may be proinflammatory (via PGE₂ expression) during the development of inflammation, but anti-inflammatory (via PGD₂ expression) during resolution in lung fibroblasts. Recent studies have also highlighted a role for COX-2-derived PGs serving anti-inflammatory and anti-fibrotic roles in the resolution of inflammation [13, 45, 46]. In a model of spontaneously

resolving ALI, selective COX-2 inhibition results in prolonged inflammation, in part, by decreasing production of pro-resolving mediators, including LXA₄ and 15-epimer-LXA₄ [44, 45]. So, a late, anti-inflammatory effect of COX-2, instead of the more widely appreciated early, proinflammatory action, was crucial to the timely recovery from ALI [45].

Our results also demonstrated that the expression of COX-2 as well as PGE₂ production by fibroblast cells was significantly inhibited by LXA₄ in a dose-dependent manner 6-hour LPS treatment, suggesting that LXA₄ has a potential anti-inflammatory role in lung fibroblasts during the onset of inflammation. Consistent with our findings, similar results have shown that LXA₄ also repressed the expression and




FIGURE 3: The effect of LXA₄ on LPS-induced expression of COX-2 protein expression and PGE₂ and PGD₂ production at 6 hours in primary lung fibroblasts. (a) Rat lung fibroblast cells were treated with LXA₄ at 0, 100, 200 or 400 nM in the presence of LPS (1 μ g/mL) for 6 hours. Cells were then harvested, sonicated and COX-2 protein detected by western blot. * $P < 0.05$ versus control group, # $P < 0.05$ versus (100, 200 or 400 nM) LXA₄ groups; & $P < 0.05$ versus (200 or 400 nM) LXA₄ groups. (b) Supernatants from rat lung fibroblast cells treated with LXA₄ at 0, 100, 200, or 400 nM in the presence of LPS (1 μ g/mL) for 6 hours were collected and PGE₂ protein measured by ELISA. Data are expressed as mean \pm SE for each group. * $P < 0.05$ versus control group, # $P < 0.05$ versus (100, 200, or 400 nM) LXA₄ groups; & $P < 0.05$ versus (200 or 400 nM) LXA₄ groups. (c) Supernatants from rat lung fibroblast cells treated with LXA₄ at 0, 100, 200 or 400 nM in the presence of LPS (1 μ g/mL) for 6 hours were collected and PGD₂ protein measured by ELISA. Data are expressed as mean \pm SE for each group. * $P < 0.05$ versus control group.

the activity of COX-2 on endotoxin-induced uveitis in rats [28]. Interestingly, the expression of COX-2 by fibroblast cells of its second increase (24 hours) was significantly promoted by LXA₄ in a dose-dependent manner. In addition, and consistent with the results above, LXA₄ inhibited the production of PGE₂ while promoted the production of PGD₂ in the supernatants. Therefore, our study demonstrates a novel biphasic role of LXA₄ on the expression of COX-2 and the production of PGE₂ and PGD₂, suggesting that LXA₄ has a potential anti-inflammatory and proresolving roles in LPS-stimulated lung fibroblasts.

As the endogenous “braking signals” in inflammation [18–20], lipoxins are produced locally in the lung to regulate inflammatory cells. Furthermore, the specific receptor with high affinity for LXA₄ (LXA₄R) has been cloned from myeloid lineages [15, 16]. Expression of LXA₄R is required to evoke actions of lipoxins in each tissue therefore, the receptor expression can also control biological function of lipoxins *in vivo*. LXA₄R belongs to the G-protein coupled receptor superfamily of proteins and is widely distributed in cells and tissues [15]. Our results indicate clearly for the first time that LXA₄R mRNA was expressed in rat




FIGURE 4: The effect of LXA₄ on LPS-induced expression of COX-2 protein expression and PGE₂ and PGD₂ production at 24 hours in primary lung fibroblasts. (a) Rat lung fibroblast cells were treated with LXA₄ at 0, 100, 200, or 400 nM in the presence of LPS (1 μg/mL) for 24 hours. Cells were then harvested, sonicated, and COX-2 protein detected by Western blot. **P* < 0.05 versus control group, #*P* < 0.05 versus non-LXA₄ groups; & *P* < 0.05 versus (100 or 200 nM) LXA₄ groups. (b) Supernatants from rat lung fibroblast cells treated with LXA₄ at 0, 100, 200, or 400 nM in the presence of LPS (1 μg/mL) for 24 hours were collected and PGE₂ protein measured by ELISA. Data are expressed as mean ± SE for each group. **P* < 0.05 versus control group. (c) Supernatants from rat lung fibroblast cells treated with LXA₄ at 0, 100, 200 or 400 nM in the presence of LPS (1 μg/mL) for 24 hours were collected and PGD₂ protein measured by ELISA. Data are expressed as mean ± SE for each group. **P* < 0.05 versus control group, #*P* < 0.05 versus (0, 100 or 200 nM) LXA₄ groups.

lung fibroblasts and is upregulated by LPS stimulation. Moreover, possibly due to a negative feedback mechanism, LXA₄R mRNA was inhibited by cotreatment of LPS and LXA₄.

In summary, this study demonstrated that COX-2 protein expression peaks initially at 6 hours but then also at 24 hours after LPS stimulation in isolated lung fibroblasts. Moreover, LXA₄ has a novel biphasic role on expression of COX-2 and production of PGE₂ and PGD₂, whereby LXA₄ has an anti-inflammatory and proresolving activity in lung fibroblasts following LPS stimulation. Therefore, our study

may provide a novel target for future therapies for controlling LPS-induced ALI.

Abbreviations

LPS: Lipopolysaccharide
 COX-2: Cyclooxygenase-2
 PGE₂: ProstaglandinE₂
 PGD₂: ProstaglandinD₂
 PGs: Prostaglandins
 LXA₄: LipoxinA₄
 LXA₄R: LipoxinA₄ receptor.




FIGURE 5: LipoxinA₄ receptor is expressed in rat lung fibroblasts and upregulated by LPS. Rat lung fibroblast cells were treated with LPS (1 μg/mL) in the presence or absence of LX A₄ (100 nM) for 24 hours. mRNA was isolated and semiquantitative RT-PCR for LX A₄R and β-actin were performed. Amplified cDNA was separated by 1.6% agarose gel electrophoresis, visualized with ethidium bromide and analyzed by gel densitometry relative to β-actin expression. Lane 1: untreated control. Lane 2: LPS only (1 μg/mL). Lane 3: LPS (1 μg/mL) and LX A₄ (100 nM). **P* < 0.05 versus control group, #*P* < 0.05 versus (100 nM) LX A₄ group. These data represent 6 individual experimental repeats.

Acknowledgment

This work was sponsored by Zhejiang Provincial Program for the Cultivation of High-level Innovative Health talents and supported by the grants from the National Natural Science Foundation of China (no. 30772088, no. 81070061).

References

- [1] H. Kirpalani and J. Gauldie, "Differentiation and effector function of pulmonary fibroblasts," in *Asthma and Rhinitis*, W. W. Busse and S. T. Holgate, Eds., pp. 539–553, Blackwell Scientific, Boston, Mass, USA, 1995.
- [2] C. D. Buckley, D. Pilling, J. M. Lord, A. N. Akbar, D. Scheel-Toellner, and M. Salmon, "Fibroblasts regulate the switch from acute resolving to chronic persistent inflammation," *Trends in Immunology*, vol. 22, no. 4, pp. 199–204, 2001.
- [3] R. S. Smith, T. J. Smith, T. M. Blieden, and R. P. Phipps, "Fibroblasts as sentinel cells. Synthesis of chemokines and regulation of inflammation," *The American Journal of Pathology*, vol. 151, no. 2, pp. 317–322, 1997.
- [4] A. Zdařilová, A. R. Svobodová, K. Chytilová, V. Šimánek, and J. Ulrichová, "Polyphenolic fraction of *Lonicera caerulea* L. fruits reduces oxidative stress and inflammatory markers induced by lipopolysaccharide in gingival fibroblasts," *Food and Chemical Toxicology*, vol. 48, no. 6, pp. 1555–1561, 2010.
- [5] B. Rosón, J. Carratalà, N. Fernández-Sabé, F. Tubau, F. Manresa, and F. Gudiol, "Causes and factors associated with early failure in hospitalized patients with community-acquired pneumonia," *Archives of Internal Medicine*, vol. 164, no. 5, pp. 502–508, 2004.
- [6] C. W. Chow, M. T. H. Abreu, T. Suzuki, and G. P. Downey, "Oxidative stress and acute lung injury," *American Journal of Respiratory Cell and Molecular Biology*, vol. 29, no. 4, pp. 427–431, 2003.
- [7] J. Beck, R. Garcia, G. Heiss, P. S. Vokonas, and S. Offenbacher, "Periodontal disease and cardiovascular disease," *Journal of Periodontology*, vol. 67, no. 10, pp. 1123–1137, 1996.
- [8] W. L. Smith, R. Michael Garavito, and D. L. DeWitt, "Prostaglandin endoperoxide H synthases (cyclooxygenases)-1 and -2," *The Journal of Biological Chemistry*, vol. 271, no. 52, pp. 33157–33160, 1996.
- [9] C. D. Funk, "Prostaglandins and leukotrienes: advances in eicosanoid biology," *Science*, vol. 294, no. 5548, pp. 1871–1875, 2001.
- [10] C. Y. Wu, P. L. Chi, H. L. Hsieh, S. F. Luo, and C. M. Yang, "TLR4-dependent induction of vascular adhesion molecule-1 in rheumatoid arthritis synovial fibroblasts: roles of cytosolic phospholipase a 2α/ cyclooxygenase-2," *Journal of Cellular Physiology*, vol. 223, no. 2, pp. 480–491, 2010.
- [11] I. Morita, "Distinct functions of COX-1 and COX-2," *Prostaglandins and Other Lipid Mediators*, vol. 68–69, pp. 165–175, 2002.
- [12] H. Mizuno, C. Sakamoto, K. Matsuda et al., "Induction of cyclooxygenase 2 in gastric mucosal lesions and its inhibition by the specific antagonist delays healing in mice," *Gastroenterology*, vol. 112, no. 2, pp. 387–397, 1997.
- [13] D. W. Gilroy, P. R. Colville-Nash, D. Willis, J. Chivers, M. J. Paul-Clark, and D. A. Willoughby, "Inducible cyclooxygenase may have anti-inflammatory properties," *Nature Medicine*, vol. 5, no. 6, pp. 698–701, 1999.
- [14] L. Vong, J. G. P. Ferraz, R. Panaccione, P. L. Beck, and J. L. Wallace, "A pro-resolution mediator, prostaglandin D₂, is specifically up-regulated in individuals in long-term remission from ulcerative colitis," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 26, pp. 12023–12027, 2010.
- [15] H. Kühn and V. B. O'Donnell, "Inflammation and immune regulation by 12/15-lipoxygenases," *Progress in Lipid Research*, vol. 45, no. 4, pp. 334–356, 2006.

- [16] B. D. Levy, "Lipoxins and lipoxin analogs in asthma," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 73, no. 3-4, pp. 231-237, 2005.
- [17] C. N. Serhan, "Controlling the resolution of acute inflammation: a new genus of dual anti-inflammatory and proresolving mediators," *Journal of Periodontology*, vol. 79, no. 8, pp. 1520-1526, 2008.
- [18] S. J. O'Meara, K. Rodgers, and C. Godson, "Lipoxins: update and impact of endogenous pro-resolution lipid mediators," *Reviews of Physiology, Biochemistry and Pharmacology*, vol. 160, pp. 47-70, 2008.
- [19] C. N. Serhan, "Resolution phase of inflammation: novel endogenous anti-inflammatory and proresolving lipid mediators and pathways," *Annual Review of Immunology*, vol. 25, pp. 101-137, 2007.
- [20] C. N. Serhan, S. Yacoubian, and R. Yang, "Anti-inflammatory and proresolving lipid mediators," *Annual Review of Pathology*, vol. 3, pp. 279-312, 2008.
- [21] B. D. Levy, "Lipoxins and lipoxin analogs in asthma," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 73, no. 3-4, pp. 231-237, 2005.
- [22] C. Bandeira-Melo, M. F. Serra, B. L. Diaz et al., "Cyclooxygenase-2-derived prostaglandin E₂ and lipoxin A₄ accelerate resolution of allergic edema in *Angiostrongylus costaricensis*-infected rats: relationship with concurrent eosinophilia," *Journal of Immunology*, vol. 164, no. 2, pp. 1029-1036, 2000.
- [23] C. L. Karp, L. M. Flick, K. W. Park et al., "Defective lipoxin-mediated anti-inflammatory activity in the cystic fibrosis airway," *Nature Immunology*, vol. 5, no. 4, pp. 388-392, 2004.
- [24] C. L. Karp, L. M. Flick, R. Yang, J. Uddin, and N. A. Petasis, "Cystic fibrosis and lipoxins," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 73, no. 3-4, pp. 263-270, 2005.
- [25] S. H. Wu, X. H. Wu, C. Lu, L. Dong, and Z. Q. Chen, "Lipoxin A₄ inhibits proliferation of human lung fibroblasts induced by connective tissue growth factor," *American Journal of Respiratory Cell and Molecular Biology*, vol. 34, no. 1, pp. 65-72, 2006.
- [26] C. Bonnans, K. Fukunaga, M. A. Levy, and B. D. Levy, "Lipoxin A₄ regulates bronchial epithelial cell responses to acid injury," *American Journal of Pathology*, vol. 168, no. 4, pp. 1064-1072, 2006.
- [27] J. C. Bonner, A. B. Rice, J. L. Ingram et al., "Susceptibility of cyclooxygenase-2-deficient mice to pulmonary fibrogenesis," *American Journal of Pathology*, vol. 161, no. 2, pp. 459-470, 2002.
- [28] R. Medeiros, G. B. Rodrigues, C. P. Figueiredo et al., "Molecular mechanisms of topical anti-inflammatory effects of lipoxin A₄ in endotoxin-induced uveitis," *Molecular Pharmacology*, vol. 74, no. 1, pp. 154-161, 2008.
- [29] M. Tamm, M. Roth, M. Malouf et al., "Primary fibroblast cell cultures from transbronchial biopsies of lung transplant recipients," *Transplantation*, vol. 71, no. 2, pp. 337-339, 2001.
- [30] E. Motohashi, H. Kawachi, H. Endo et al., "Regulatory expression of lipoxin A₄ receptor in physiologically estrus cycle and pathologically endometriosis," *Biomedicine and Pharmacotherapy*, vol. 59, no. 6, pp. 330-338, 2005.
- [31] J. P. Louboutin, B. A. S. Reyes, L. Agrawal, C. R. Maxwell, E. J. Van Bockstaele, and D. S. Strayer, "Blood-brain barrier abnormalities caused by exposure to HIV-1 gp120 - Protection by gene delivery of antioxidant enzymes," *Neurobiology of Disease*, vol. 38, no. 2, pp. 313-325, 2010.
- [32] M. Raval, P. G. Frank, L. Laury-Kleintop, G. Yan, and S. Lanza-Jacoby, "Celecoxib combined with atorvastatin prevents progression of atherosclerosis," *Journal of Surgical Research*, vol. 163, no. 2, pp. e113-e122, 2010.
- [33] J. O'Brien, T. Lyons, J. Monks et al., "Alternatively activated macrophages and collagen remodeling characterize the postpartum involuting mammary gland across species," *American Journal of Pathology*, vol. 176, no. 3, pp. 1241-1255, 2010.
- [34] A. Lührmann, K. Grote, M. Stephan, T. Tschernig, and R. Pabst, "Local pulmonary immune stimulation by the Toll-like receptor 2 and 6 ligand MALP-2 in rats is age dependent," *Immunology Letters*, vol. 108, no. 2, pp. 167-173, 2007.
- [35] J. U. Scher and M. H. Pillinger, "15d-PGJ₂: the anti-inflammatory prostaglandin?" *Clinical Immunology*, vol. 114, no. 2, pp. 100-109, 2005.
- [36] E. Estenssoro, A. Dubin, E. Laffaire et al., "Incidence, clinical course, and outcome in 217 patients with acute respiratory distress syndrome," *Critical Care Medicine*, vol. 30, no. 11, pp. 2450-2456, 2002.
- [37] F. Frutos-Vivar, N. Nin, and A. Esteban, "Epidemiology of acute lung injury and acute respiratory distress syndrome," *Current Opinion in Critical Care*, vol. 10, no. 1, pp. 1-6, 2004.
- [38] G. D. Rubenfeld, E. Caldwell, E. Peabody et al., "Incidence and outcomes of acute lung injury," *The New England Journal of Medicine*, vol. 353, no. 16, pp. 1685-1693, 2005.
- [39] L. B. Ware and M. A. Matthay, "The acute respiratory distress syndrome," *The New England Journal of Medicine*, vol. 342, no. 18, pp. 1334-1349, 2000.
- [40] D. W. Gilroy, T. Lawrence, M. Perretti, and A. G. Rossi, "Inflammatory resolution: new opportunities for drug discovery," *Nature Reviews Drug Discovery*, vol. 3, no. 5, pp. 401-416, 2004.
- [41] C. N. Serhan, "Resolution phase of inflammation: novel endogenous anti-inflammatory and proresolving lipid mediators and pathways," *Annual Review of Immunology*, vol. 25, pp. 101-137, 2007.
- [42] S. W. Jin, L. Zhang, Q. Q. Lian et al., "Posttreatment with aspirin-triggered lipoxin A₄ analog attenuates lipopolysaccharide-induced acute lung injury in mice: the role of heme oxygenase-1," *Anesthesia and Analgesia*, vol. 104, no. 2, pp. 369-377, 2007.
- [43] D. Takai, T. Nagase, and T. Shimizu, "New therapeutic key for cystic fibrosis: a role for lipoxins," *Nature Immunology*, vol. 5, no. 4, pp. 357-358, 2004.
- [44] A. Tomlinson, I. Appleton, A. R. Moore et al., "Cyclooxygenase and nitric oxide synthase isoforms in rat carrageenin-induced pleurisy," *British Journal of Pharmacology*, vol. 113, no. 3, pp. 693-698, 1994.
- [45] K. Fukunaga, P. Kohli, C. Bonnans, L. E. Fredenburgh, and B. D. Levy, "Cyclooxygenase 2 plays a pivotal role in the resolution of acute lung injury," *Journal of Immunology*, vol. 174, no. 8, pp. 5033-5039, 2005.
- [46] C. Bandeira-Melo, M. F. Serra, B. L. Diaz et al., "Cyclooxygenase-2-derived prostaglandin E₂ and lipoxin A₄ accelerate resolution of allergic edema in *Angiostrongylus costaricensis*-infected rats: relationship with concurrent eosinophilia," *Journal of Immunology*, vol. 164, no. 2, pp. 1029-1036, 2000.