



University
of Glasgow

McMillan, D.C. and Hole, D.J. and McArdle, C.S. (2008) *The impact of old age on cancer-specific and non-cancer-related survival following elective potentially curative surgery for Dukes A/B colorectal cancer.* British Journal of Cancer, 99 . pp. 1046-1049.

<http://eprints.gla.ac.uk/4822/>

17th December 2008

The impact of old age on cancer specific and non-cancer related survival following elective potentially curative surgery for Dukes A/B colorectal cancer.

Donald C McMillan¹, David J Hole², Colin S McArdle¹.

University Departments of Surgery¹ and Public Health², Faculty of Medicine- University of Glasgow, Glasgow G12 8QQ

David J Hole died March 26th 2007

Running title: Old age and outcome following elective curative colorectal cancer surgery

Key words: colorectal cancer, age, elective, curative surgery, survival

Support: Clinical Resource and Audit Group (CRAG), Chief Scientist Office, Scotland.

Correspondence to:

Dr Donald C McMillan,

University Department of Surgery,

Royal Infirmary,

Glasgow, G31 2ER,

United Kingdom.

Tel No. 0141 211 5435

Fax. No. 0141 552 3229

E-mail d.c.mcmillan@clinmed.gla.ac.uk

Abstract

Previous studies have suggested that survival following surgery for colorectal cancer is poorer in the elderly. However, the findings were inconsistent and none of the studies adjusted for case mix. The aim of the present study was to establish whether there were age-related differences in cancer (colorectal) specific and non-cancer (colorectal) related survival in patients undergoing elective potentially curative resection for Dukes stage A/ B colorectal cancer. One thousand and forty three patients who underwent elective potentially curative resection for Dukes' A/B colorectal cancer between 1991-1994 in 11 hospitals in Scotland were included in the study. Ten year cancer specific and non-cancer related survival and the hazard ratios were calculated according to age groups (<64, 65-74/ >74 years). On follow-up 273 patients died of their cancer and 328 died of non-cancer related causes. At 10 years overall survival was 45%, cancer specific was 70% and non-cancer related survival was 64%. On multivariate analysis of all factors, age (HR 1.38, 95%CI 1.18-1.62, $p<0.001$), sex (HR 1.74, 95%CI 1.36-2.23, $p<0.001$), site (HR 1.42, 95%CI 1.11-1.81, $p<0.01$) and Dukes' stage (HR 1.71, 1.19-2.47, $p<0.01$) were independently associated with cancer specific survival. On multivariate analysis of all factors, age (HR 2.14, 1.84-2.49, $p<0.001$), sex (HR 1.43, 1.15-1.79, $p<0.01$) and deprivation (HR 1.30, 1.09-1.55, $p<0.01$) were independently associated with non-cancer related survival. The results of the present study show that increasing age impacts negatively both on cancer specific and non-cancer related survival following elective potentially curative resection for node negative colorectal cancer. However, the effect of increasing age is greater on the non-cancer related survival. These results suggest that cancer specific and non-cancer related mortality should be considered separately in survival analysis of these cancer patients.

Introduction

Colorectal cancer is the second commonest cause of cancer death in Western Europe and North America. Many patients have evidence of locally advanced or metastatic disease at the time of initial presentation. Even in those undergoing apparently curative resection, only half survive 5 years (McArdle and Hole, 2002a).

It has long been recognised that there are a number of factors, in addition to pathological stage, which contribute to poor outcome following potentially curative surgery for colorectal cancer. Age (Mulcahy et al., 1994; Shankar and Taylor, 1998; Colorectal Cancer Collaborative Group, 2000), gender (McArdle et al., 2003), deprivation (Hole and McArdle, 2002), tumour site (McArdle and Hole, 2002b), emergency presentation (McArdle and Hole, 2004a) and specialisation (McArdle and Hole, 2004b) have been shown to impact on long term survival in these patients.

However, whether old age is associated with poorer survival, independent of these factors, remains unclear. The aim of the present study was to establish whether there were age-related differences in cancer (colorectal) specific and non-cancer (colorectal) related survival in patients undergoing elective potentially curative resection for Dukes stage A/ B colorectal cancer.

Patients and methods

One thousand and forty three patients who underwent an elective potentially curative resection for Dukes A/ B colorectal cancer between 1st January 1991 and 31st December 1994 in eleven hospitals in the central belt of Scotland were included in the study. Information was abstracted from casenotes by two specially trained data managers. Details included age, sex, deprivation category, site of tumour, extent of tumour spread, the nature of surgery, post-operative mortality, Dukes' stage and adjuvant therapy. Data for 1991 and 1992 were collected retrospectively, and those for 1993 and 1994 were collected prospectively. There was no difference in baseline characteristics of the patients between the two periods.

The extent of deprivation was defined using the Carstairs Index (Carstairs and Morris, 1991), an area-based measure derived from the 1991 census data based on the postcode of residents at diagnosis. Carstairs divides the scores into a seven-point scale ranging from most affluent (category 1) to most deprived (category 7).

Tumours were classified according to site colon or rectum. The extent of tumour spread was assessed by conventional Dukes' classification based on histological examination of the resected specimen.

Patients were deemed to have had a curative resection if the surgeon considered that there was no macroscopic residual tumour once resection had been completed.

Individual surgeons were defined as specialists or non-specialists by a panel of six senior consultants and one of the authors (C.S.McA.). Two of the six consultants were specialist colorectal surgeons from teaching hospitals and four were district general hospital consultants. These assessments were made without knowledge of the outcome and before any analysis was performed.

Approval was obtained for information on date and cause of death to be checked with that received by the cancer registration system through linkage with the Registrar General (Scotland). Deaths up to the end of 2003 have been included in the analysis, providing an average length of follow-up of 11 years (minimum 9 years, maximum 13 years).

Statistical analysis

The percentages of patients surviving 10 years were calculated using the Kaplan-Meier technique. Comparison of the association between age and other variables was made using the X^2 test or a X^2 for trend where appropriate. The effect of age on cancer and non-cancer related survival was examined using Cox's proportional hazards model. Analysis was performed using the SPSS software package (SPSS Inc., Chicago, IL, USA).

Results

Of the 1043 patients included in the analysis, 33% were aged 75 or over, 21% were socio-economically deprived, 59% had colonic tumours and 82% had Dukes' B disease at the time of surgery. 275 (26%) patients were treated by a specialist surgeon. Three percent of patients received adjuvant therapy.

The baseline characteristics of the patients included in the study are shown in Table 1. With increasing age there was an increase in the proportion of patients who had colonic tumours ($p<0.001$). Few patients over the age of 74 years received adjuvant therapy ($p<0.001$).

273 patients died of their cancer and 328 died of non-cancer related causes (Table 2). Of the non-cancer related deaths 212 (65%) patients died of cardiovascular or respiratory disease. At 10 years overall survival was 45%, cancer specific was 70% and non-cancer related survival was 64%. An increased proportion of cancer deaths at 10 years was associated with increased Dukes' stage ($p<0.05$) and non-specialist surgeons ($p<0.10$). An increased proportion of non-cancer related deaths at 10 years was associated with older age ($p<0.001$), female gender ($p<0.05$) and deprivation ($p<0.10$).

On univariate analysis, age ($p<0.01$), sex ($p<0.001$), site ($p<0.05$) and Dukes' stage ($p<0.01$) were significantly associated with cancer specific survival (Table 3). On multivariate analysis of all factors, age (HR 1.38, 95%CI 1.18-1.62, $p<0.001$), sex (HR 1.74, 95%CI 1.36-2.23, $p<0.001$), site (HR 1.42, 95%CI 1.11-1.81, $p<0.01$) and Dukes' stage (HR 1.71, 1.19-2.47, $p<0.01$) were independently associated with cancer specific survival.

On univariate analysis, age ($p<0.001$), sex ($p<0.05$), deprivation ($p<0.01$) and adjuvant therapy ($p<0.10$) were significantly associated with non-cancer related survival (Table 4). On multivariate analysis of all factors, age (HR 2.14, 1.84-2.49, $p<0.001$), sex

(HR 1.43, 1.15-1.79, $p < 0.01$) and deprivation (HR 1.30, 1.09-1.55, $p < 0.01$) were independently associated with non-cancer related survival.

The relationship between age and the hazard ratios for cancer and non-cancer related survival in patients with Dukes' A/ B disease colon and rectum cancer is shown in Table 5. Compared with patients under the age 64years who had colonic tumours, those patients over the age of 74years had a hazard ratio of 2.2 ($P < 0.01$) and 5.4 ($p < 0.001$) for cancer and non cancer related survival respectively. Compared with patients under the age 64years who had rectal tumours, those patients over the age of 74years had a hazard ratio of 5.1 ($p < 0.001$) for non cancer related survival.

Discussion

The results of the present study show that, in patients undergoing elective potentially curative surgery for Dukes' A/ B colorectal cancer, with 10-year follow-up, more patients die of intercurrent disease, in particular cardiovascular or respiratory disease, than die of cancer. Increasing age, 75years and over, impacted on cancer related and non-cancer related survival. Although, it was notable that the effect on non-cancer related survival was approximately twice that of cancer related survival.

Given that colorectal cancer is a common cancer and that large numbers of patients undergoing elective potentially curative resection will die of intercurrent comorbid disease, the use of overall survival as an end-point may give misleading results. Therefore, future studies of these patients should specify the associations between factors and cancer specific survival and non-cancer related survival separately in order to establish whether the factor impacts equally on cancer specific and non-cancer related survival.

The results of the present study have a number of important implications. Firstly, with the introduction of screening programmes for colorectal cancer and the consequent increase in early stage disease it is likely that the issue of age-related comorbid death will become a major issue in these patients. If this were proven to be the case it would be important that within screening programmes for colorectal cancer, patients should be also screened for the presence of comorbid disease. For example, in patients aged of 75 years and over undergoing potentially curative resection for node negative colorectal cancer, it would be important not only to treat the cancer, but also the comorbid disease, in particular it would appear from the present study that this is cardiovascular or respiratory disease.

In the context of the present study it is of interest that a strong association between colorectal cancer and vascular disease has been reported (Chan et al., 2006; Chan et al., 2007). This may be due to shared environmental risk factors such as age, smoking, diabetes

mellitus and obesity (Giovannucci and Martínez, 1996; Larsson et al., 2005; McMillan et al., 2006). It has also been hypothesised that these diseases develop through a common pathway of chronic inflammatory disease (Chan et al., 2006; Chan et al., 2007). Indeed, there is evidence that such environmental factors are directly associated with C-reactive protein concentrations, the prototypical marker of the systemic inflammatory response, in apparently healthy subjects (Koenig et al., 1999; Hutchinson et al., 2000; Chenillot et al., 2000; Wong et al., 2001). However, it remains to be determined whether the systemic inflammatory response impacts equally on cancer specific and non-cancer related survival in patients undergoing potentially curative surgery for colorectal cancer.

In summary, the results of the present show that although age impacts both on cancer specific and non-cancer related survival following elective potentially curative resection for node negative colorectal cancer, the effect is greater on non-cancer related survival. These results suggest that cancer specific and non-cancer related deaths should be considered separately in survival analysis of these patients.

Acknowledgements

The authors thank the surgeons who participated in the study and Heather Wotherspoon and Janette Stevenson, who collected the baseline data. They also thank the Clinical Resource and Audit Group (CRAG), Scottish Home and Health Department, Scotland who funded the collection of the baseline data.

References

- Chan AO, Lam KF, Tong T, Siu DC, Jim MH, Hui WM, Lai KC, Yuen MF, Lam SK, Wong BC (2006). Coexistence between colorectal cancer/adenoma and coronary artery disease: results from 1382 patients. *Aliment Pharmacol Ther.* 24:535-9.
- Chan AO, Jim MH, Lam KF, Morris JS, Siu DC, Tong T, Ng FH, Wong SY, Hui WM, Chan CK, Lai KC, Cheung TK, Chan P, Wong G, Yuen MF, Lau YK, Lee S, Szeto ML, Wong BC, Lam SK (2007). Prevalence of colorectal neoplasm among patients with newly diagnosed coronary artery disease. *JAMA.* 298:1412-9.
- Carstairs V, Morris R (1991). *Deprivation and Health in Scotland.* Aberdeen University Press: Aberdeen.
- Chenillot O, Henny J, Steinmetz J, Herbeth B, Wagner C, Siest G (2000). High sensitivity C-reactive protein: biological variations and reference limits. *Clin Chem Lab Med* 10:1003–11.
- Colorectal Cancer Collaborative Group (2000) Surgery for colorectal cancer in elderly patients: a systematic review. *Lancet.* 356(9234):968-74.
- Giovannucci E, Martínez ME (1996). Tobacco, colorectal cancer, and adenomas: a review of the evidence. *J Natl Cancer Inst.* 88:1717-30.
- Hole DJ, McArdle CS (2002). Impact of socioeconomic deprivation on outcome after surgery for colorectal cancer. *Br J Surg.* 89:586-90.
- Hutchinson WL, Koenig W, Frohlich M, Sund M, Lowe GD, Pepys MB. (2000) Immunoradiometric assay of circulating C-reactive protein: age-related values in the adult general population. *Clin Chem* 7: 934–8.
- Koenig W, Sund M, Frohlich M, Fischer HG, Lowel H, Doring A, Hutchinson WL, Pepys MB. (1999) C-reactive protein, a sensitive marker of inflammation, predicts future risk of coronary heart disease in initially healthy middle-aged men: results from the MONICA

(Monitoring Trends and Determinants in Cardiovascular Disease) Augsburg Cohort Study, 1984–1992. *Circulation* 99: 237–242.

Larsson SC, Orsini N, Wolk A (2005). Diabetes mellitus and risk of colorectal cancer: a meta-analysis. *J Natl Cancer Inst.* 97:1679-87.

McArdle CS, Hole DJ (2002a) Outcome following surgery for colorectal cancer: analysis by hospital after adjustment for case-mix and deprivation. *Br J Cancer* 86: 331–335.

McArdle CS, Hole DJ (2002b). Outcome following surgery for colorectal cancer. *Br Med Bull.* 64:119-25.

McArdle CS, McMillan DC, Hole DJ (2003). Male gender adversely affects survival following surgery for colorectal cancer. *Br J Surg.* 90:711-5.

McArdle CS, Hole DJ (2004a). Emergency presentation of colorectal cancer is associated with poor 5-year survival. *Br J Surg.* 91:605-9.

McArdle CS, Hole DJ (2004b). Influence of volume and specialization on survival following surgery for colorectal cancer. *Br J Surg.*2004; 91:610-7.

McMillan DC, Sattar N, McArdle CS (2006). ABC of obesity. Obesity and cancer. *BMJ.* 333(7578):1109-11.

Mulcahy HE, Patchett SE, Daly L, O'Donoghue DP (1994). Prognosis of elderly patients with large bowel cancer. *Br J Surg.* 81:736-8.

Shankar A, Taylor I (1998). Treatment of colorectal cancer in patients aged over 75. *Eur J Surg Oncol.* 24:391-5.

Wong ND, Pio J, Valencia R, Thakal G (2001). Distribution of C-reactive protein and its relation to risk factors and coronary heart disease risk estimation in the National Health and Nutrition Examination Survey (NHANES) III. *Prev Cardiol* 4: 109–114.

Table 1: Baseline characteristics of patients undergoing elective curative resection for Dukes A/B colorectal cancer by age (n= 1043)

	Age (years)			p value
	<64yrs n= 305 (29%)	65-74yrs n= 398 (38%)	>74 n= 340 (33%)	
Female	143 (47%)	186 (47%)	184 (56%)	0.061
Male	162 (53%)	212 (53%)	156 (46%)	
Depcat Score				0.625
1, 2	53 (17%)	70 (18%)	69 (20%)	
3, 4, 5	191 (63%)	245 (62%)	201 (59%)	
6, 7	61 (20%)	83 (21%)	70 (21%)	
Site				<0.001
Colon	160 (52%)	228 (57%)	223 (66%)	
Rectum	145 (48%)	170 (43%)	117 (37%)	
Dukes' Stage				0.822
A	52 (17%)	76 (19%)	56 (17%)	
B	253 (83%)	322 (81%)	284 (83%)	
Specialisation				0.402
Specialist	76 (25%)	104 (27%)	95 (28%)	
Non-specialist	223 (75%)	288 (73%)	240 (72%)	
Adjuvant therapy				<0.001
Yes	15 (5%)	16 (4%)	3 (1%)	
No	290 (95%)	382 (96%)	337 (99%)	

Table 2: Baseline characteristics of patients undergoing elective curative resection for Dukes A/B colorectal cancer by cancer and non-cancer death after 10 years follow-up

	Cancer deaths (n= 273)	Non-cancer deaths (n= 328)	p value
Age			
<64yrs	68 (25%)	42 (13%)	
65-74yrs	109 (40%)	129 (39%)	
>74yrs	96 (35%)	157 (48%)	<0.001
Female	104 (38%)	153 (47%)	
Male	169 (62%)	175 (53%)	0.035
Depcat Score			
1, 2	53 (19%)	45 (14%)	
3, 4, 5	167 (62%)	206 (63%)	
6, 7	53 (19%)	77 (24%)	0.052
Site			
Colon	148 (54%)	192 (59%)	
Rectum	125 (46%)	136 (41%)	0.287
Dukes' Stage			
A	34 (13%)	61 (19%)	
B	239 (87%)	267 (81%)	0.040
Specialisation			
Specialist	61 (23%)	94 (27%)	
Non-specialist	209 (77%)	226 (71%)	0.062
Adjuvant therapy			
Yes	10 (4%)	6 (2%)	
No	263 (96%)	322 (98%)	0.165

Table 3. The relationship between clinicopathological characteristics and cancer specific survival in patients undergoing elective curative resection for Dukes A/B colorectal cancer: Univariate and multivariate analysis.

	Patients (n= 1043)	Univariate analysis		Multivariate analysis	
		Hazard ratio (95% CI)	<i>P</i> -value	Hazard ratio (95% CI)	<i>P</i> -value
Age (<64/ 65-74/ >74years)	305/ 398/ 340	1.29 (1.11-1.50)	0.001	1.38 (1.18-1.62)	<0.001
Sex (female/ male)	513/ 530	1.75 (1.37-2.23)	<0.001	1.74 (1.36-2.23)	<0.001
Deprivation (1-2/ 3-5/ 6-7)*	192/ 637/ 214	0.98 (0.81-1.19)	0.840		0.322
Site (colon/ rectum)	611/ 432	1.28 (1.01-1.62)	0.044	1.42 (1.11-1.81)	0.006
Dukes' stage (A/ B)	184/ 859	1.70 (1.19-2.44)	0.004	1.71 (1.19-2.47)	0.004
Specialisation (yes/ no)	275/ 751	1.27 (0.95-1.69)	0.101		0.104
Adjuvant therapy (yes/ no)	34/ 1009	0.94 (0.50-1.77)	0.845		0.750

* Individual deprivation categories were used in the statistical analysis

Table 4. The relationship between clinicopathological characteristics and non-cancer survival in patients undergoing elective curative resection for Dukes A/B colorectal cancer: Univariate and multivariate analysis.

	Patients (n= 1043)	Univariate analysis		Multivariate analysis	
		Hazard ratio (95% CI)	<i>P</i> -value	Hazard ratio (95% CI)	<i>P</i> -value
Age (<64/ 65-74/ >74years)	305/ 398/ 340	2.11 (1.82-2.45)	<0.001	2.14 (1.84-2.49)	<0.001
Sex (female/ male)	513/ 530	1.32 (1.06-1.64)	0.012	1.43 (1.15-1.79)	0.002
Deprivation (1-2/ 3-5/ 6-7)*	192/ 637/ 214	1.30 (1.09-1.53)	0.003	1.30 (1.09-1.55)	0.003
Site (colon/ rectum)	611/ 432	1.11 (0.88-1.37)	0.407		0.094
Dukes' stage (A/ B)	184/ 859	1.07 (0.81-1.42)	0.676		0.346
Specialisation (yes/ no)	275/ 751	0.89 (0.70-1.13)	0.332		0.548
Adjuvant therapy (yes/ no)	34/ 1009	2.11 (0.96-4.87)	0.061		0.302

* Individual deprivation categories were used in the statistical analysis

Table 5: Hazard ratios for cancer specific and non-cancer survival at 10 years in patients undergoing elective curative resection for Dukes A/B colorectal cancer by age.

Colon (n= 611)	Cancer specific (deaths= 148)		Non-cancer (deaths= 192)	
	Hazard ratio (95%CI)	p-value	Hazard ratio (95%CI)	p-value
Age <64yrs	1	0.004	1	<0.001
65-74yrs	1.71 (1.09-2.69)	0.019	3.01 (1.85-4.91)	<0.001
>74yrs	2.16 (1.38-3.37)	0.001	5.39 (3.36-8.64)	<0.001
Rectum (n= 432)	Cancer specific (deaths= 125)		Non-cancer (deaths= 136)	
	Hazard ratio (95%CI)	p-value	Hazard ratio (95%CI)	p-value
Age <64yrs	1	0.286	1	<0.001
65-74yrs	1.28 (0.85-1.94)	0.237	3.02 (1.84-4.50)	<0.001
>74yrs	1.42 (0.90-2.25)	0.132	5.09 (3.08-8.41)	<0.001