

Morach 2017

1 **Population structure and virulence gene profiles of *Streptococcus***
2 ***agalactiae* collected worldwide from different hosts**

3
4 Running Title: DNA microarray typing of GBS

5

6

7 Marina Morach^a, Roger Stephan^a, Sarah Schmitt^b, Christa Ewers^c, Michael Zschöck^d, Julian
8 Reyes-Velez^{e,f}, Urs Gilli^g, María del Pilar Crespo-Ortiz^h, Margaret Crumlishⁱ, Revathi
9 Gunturu^j, Claudia A. Daubenberger^{k,l}, Margaret Ip^m, Walter Regliⁿ, Sophia Johler^{a*}

10

11 Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Zurich,
12 Switzerland^a

13 Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Zurich, Zurich,
14 Switzerland^b

15 Institute of Hygiene and Infectious Diseases in Animals, Justus-Liebig University, Giessen,
16 Germany^c

17 Hessian State Laboratory, Department of Veterinary Medicine, Giessen, Germany^d

18 Department of Health Management, Atlantic Veterinary College, University of Prince Edward
19 Island, Charlottetown, PE, Canada^e

20 Tropical Medicine Institute of Colombia, Antonio Roldan Betancur - CES University,

21 Tropical Medicine Research Group, Medellin, Colombia^f

22 IDEXX Diavet Laboratories, Bäch, Switzerland^g

23 Department of Microbiology, University of Valle, Department of Biomedical Sciences,

24 Santiago de Cali University, Cali, Colombia^h

- 25 Institute of Aquaculture, University of Stirling, Stirling, United Kingdomⁱ
26 Aga Khan University Hospital, Nairobi, Kenya^j
27 Swiss Tropical and Public Health Institute, Basel, Switzerland^k
28 University of Basel, Basel, Switzerland^l
29 Department of Microbiology, Chinese University of Hong Kong, Prince of Wales Hospital,
30 Shatin, Hong Kong, China^m
31 labor-zentral.ch, Geuensee, Switzerlandⁿ

32

33

34

35 Accepted for publication in *European Journal of Clinical Microbiology and Infectious*

36 Diseases published by Springer. The final publication is available at Springer via

37 <https://doi.org/10.1007/s10096-017-3146-x>

38

39

40

41

42

43

44

45

46

47

48 *Corresponding author. Mailing address: Winterthurerstrasse 272, CH-8057 Zurich,
49 Switzerland. Phone: +41-44-635-8655. Fax: +41-44-635-8908. E-mail: sophia.johler@uzh.ch,
50 ORCID: 0000-0003-4299-5651

51 **Abstract**

52 *Streptococcus (S.) agalactiae* is a leading cause of morbidity and mortality among neonates
53 and causes severe infections in pregnant women and nonpregnant predisposed adults, as well
54 as various animal species worldwide. Still, information on the population structure of *S.*
55 *agalactiae* and the geographical distribution of different clones is limited. Further data is
56 urgently needed to identify particularly successful clones and obtain insights into possible
57 routes of transmission within one host species and across species borders. We aimed to
58 determine the population structure and virulence gene profiles of *S. agalactiae* strains from a
59 diverse set of sources and geographical origins. To this end, 373 *S. agalactiae* isolates
60 obtained from humans and animals from five different continents were typed by DNA
61 microarray profiling. A total of 242 different *S. agalactiae* strains were identified and further
62 analyzed. Particularly successful clonal lineages, hybridization patterns, and strains were
63 identified that were spread across different continents and/or were present in more than one
64 host species. In particular, several strains were detected both in humans and cattle, and several
65 canine strains were also detected in samples from human, bovine, and porcine hosts. The
66 findings of our study suggest that while *S. agalactiae* is well adapted to various hosts
67 including humans, cattle, dogs, rodents, and fish, interspecies transmission is possible and
68 occurs between humans and cows, dogs, and rabbits. The presented virulence and resistance
69 gene profiles enable new insights into interspecies transmission and make a crucial
70 contribution in the identification of suitable targets for therapeutic agents and vaccines.

71

72

73 Keywords: genotype B *Streptococci*, GBS, transmission, capsular serotype, resistance,
74 clonality

75 **Introduction**

76 *Streptococcus (S.) agalactiae*, also known as group B *Streptococcus* (GBS), emerged
77 in the 1970s as a major cause of morbidity and mortality in neonates and pregnant women.
78 The organism leads to meningitis and septicemia in newborns and severe peripartum
79 complications in pregnant women [1]. *S. agalactiae* has been linked to disease in the elderly
80 and in nonpregnant adults suffering from chronic diseases [2,3]. The organism is also
81 commonly found in food [4] and there are some indications for foodborne/ feedborne
82 transmission [5–7]. In spite of numerous eradication programs, *S. agalactiae* is still a common
83 cause of bovine intramammary infections in many countries [8], with particularly high herd
84 prevalence levels in countries with emerging dairy industries [9].

85 Capsular polysaccharide (CPS) was recognized as a major virulence factor of *S.*
86 *agalactiae* and plays an important role in the evasion of host defence mechanisms. CPS has
87 also been used to type GBS and assign isolates to distinct CPS serotypes (Ia, Ib, and II to IX),
88 with serotypes Ia, Ib, II, III and V being highly prevalent in human invasive GBS isolates in
89 many regions of the world [10–12]. Vaccines combining these serotypes can be highly
90 effective, they fail however to offer protection against other GBS serotypes, which cause the
91 majority of GBS infections in some regions of the world such as Japan [11,12].

92 GBS strains can harbour a wide range of genes encoding virulence factors such as Bac
93 involved in immune evasion, the alpha-like proteins involved in invasion, or the pilus islands,
94 which play a role in host adaptation and specificity. GBS also frequently exhibit resistance
95 genes, including genes conferring resistance to macrolide, lincosamide, and tetracycline.
96 Recently, several studies typing and characterizing *S. agalactiae* isolates have been published
97 [13–19] and a tool for rapid GBS typing based on DNA microarray hybridization patterns
98 (HPs) has been introduced [13]. However, comprehensive information on the population
99 structure and virulence gene profiles of *S. agalactiae* and the geographical distribution of

100 different clonal lineages is extremely scarce. In particular, comprehensive data on the
101 population structure and virulence gene profile of isolates from a broad range of host species
102 is missing. This data would be crucial to obtain further insights into host adaptation, to
103 identify particularly successful clones, and to determine the geographical distribution of
104 different clonal lineages. It could also be used to identify suitable targets for vaccines and
105 antimicrobial agents, and to further elucidate possible routes of transmission.

106 A prospective cross-sectional cohort study found that exposure to cattle is a predictor
107 of human colonization with *S. agalactiae* [20]. Case reports and some GBS typing data
108 indicate possible transmission not only between human hosts and cows, but also human hosts
109 and dogs, cats, and crocodiles [21–25]. In addition, experimental studies have evidenced
110 transmission of bovine and human *S. agalactiae* strains to fish [26–28]. Still, data on
111 interspecies transmission is scarce and strain typing studies involving a diverse set of hosts
112 and geographical areas are missing.

113 Therefore, here we provide data on the population structure and virulence gene
114 profiles of *S. agalactiae* strains isolated from a diverse set of hosts and a wide variety of
115 geographical areas.

116

117

118 **Material and methods**119 **Bacterial isolates**

120 In this study, a total of 373 *S. agalactiae* isolates from 5 different continents were
121 analyzed. Countries of origin represented in this study were: Belgium (n = 1), Colombia (n =
122 86), Costa Rica (n = 1), Germany (n = 109), Honduras (n = 3), Hong Kong SAR, China (n =
123 30), Kenya (n = 33), Switzerland (n = 103), Thailand (n = 6), Vietnam (n = 1). Isolates
124 included in this study originated from human hosts (n = 225), cattle (n = 84), dogs (n = 16),
125 fish (n = 15), mice (n = 11), elephants (n = 7), guinea pigs (n = 3), emerald monitors (n = 3),
126 rats (n = 2), snakes (n = 2) and one isolate each was collected from a rabbit, a goat, a pig, a
127 turtle, and a frog. A full summary stating the host species, geographical source, and sample
128 type is provided as Online Resource 1.

129

130 **DNA extraction and DNA microarray**

131 All isolates other than fish isolates were cultivated on 5% sheep blood agar (Oxoid
132 Limited, Hampshire, UK) and incubated for 48 to 72 hours at 37°C. *S. agalactiae* isolates
133 obtained from fish were streaked on both sheep blood agar and Tryptic Soy Agar (Becton
134 Dickinson), and incubated for 72 hours at 30°C. Subsequent DNA extraction was performed
135 using a Qiagen DNeasy kit and following the recommendations of the DNA microarray
136 *S.agaType* AS-1 kit provider (Alere Technologies, Jena, Germany). As this protocol proved
137 unsuccessful in fish isolates, these isolates were cultivated in 10 mL Tryptic Soy Broth and/or
138 10 mL Brain Heart Infusion and incubated at 28°C and at 200 rpm/min for 48h or until
139 clouding of the broth culture was visible. The following day, cells were harvested by
140 centrifugation and dissolved in A1 lysis buffer, before transfer to the A2 lysis enhancer
141 Eppendorf tube, to which 400 U achromopeptidase was added. Subsequent steps were
142 performed according to the manufacturer`s protocol (Alere Technologies). A ND-100 UV-Vis

143 spectrophotometer (NanoDrop Technologies, Wilmington, Germany) was used to measure
144 DNA concentrations in all samples.

145 The DNA microarray used in this study provides data on the presence/absence of typing
146 markers (capsule/pilus-associated genes and *alp* genes), as well as genes conferring resistance
147 (resistance to macrolide/ lincosamide antibiotics, tetracycline, heavy metals) or encoding
148 virulence factors, enzymes and other metabolic functions [13]. Linear PCR amplification and
149 DNA microarray hybridization, washing steps, and staining were performed as suggested by
150 the DNA microarray manufacturer. Hybridization patterns and signal intensities were
151 measured applying an ArrayMate reader (Alere Technologies) and were used for *S. agalactiae*
152 species confirmation, assignment to a clonal complex and capsule type, hybridization pattern,
153 and strain, where possible [13].

154

155 **SplitsTree analysis**

156 Similar to Coombs et al., DNA microarray hybridization profiles were used to
157 calculate unrooted phylogenetic networks from molecular sequence data [29,30]. Stringent
158 inclusion criteria were applied to avoid bias. Multiple isolates were considered to represent
159 the same strain (*e.g.* S1) if DNA microarray hybridization results were identical for all
160 positive/negative signals. In these cases, only one *S. agalactiae* DNA microarray profile was
161 considered for construction of the SplitsTree and was included in the statistical analysis. This
162 resulted in a total number of 161 strains from humans, 52 strains from ruminants, 15 strains
163 from dogs, 8 strains from rodents, 8 strains from fish, and 12 strains from other hosts being
164 included in the statistical analysis. SplitsTree4 (www.splitstree.org) was used to depict the
165 degree of similarity of the different *S. agalactiae* hybridization patterns [31].

166

167 **Statistical analysis**

168 Statistically significant differences ($p \leq 0.050$) in the distribution of virulence and
169 resistance genes between isolates from different sources (hosts or host groups) were
170 determined either by Chi squared test or Fisher's exact test (in case $n < 5$) using SPSS 24.0
171 (IBM Corp., Armonk, NY, USA).

172

173

Results

174 The 373 GBS isolates included in this study could be assigned to 242 different strains.
175 Multiple isolates representing the same strain were detected in many host species and across
176 different countries or continents (see Table 1). We observed particularly high rates of
177 duplicates assigned to the same strain among murine (64%), piscine (47%), and bovine
178 isolates (39%). In addition, isolates representing the same *S. agalactiae* strains were not only
179 detected multiple times within one host species, but in some cases also across different host
180 species (see Fig. 1).

181 We determined pronounced host-specific differences in the frequency of different
182 clonal complexes (Table 2). In GBS from human hosts, CC19-19 was most prevalent (35%),
183 followed by CC23 (20%). In contrast, GBS strains isolated from ruminants were most
184 commonly assigned to CC23 (21%), strains from dogs to CC19-10 (40%), strains from
185 rodents to CC19-01 (75%), and strains from fish to CC260/261 (75%). Some host-specific
186 differences were also visible in the prevalence of capsular serotypes (Table 3). While serotype
187 IB was highly prevalent in GBS strains from fish (63%), it was only rarely detected in isolates
188 from other hosts. In contrast, serotypes IA, II, III, and V were common in GBS from different
189 host species. As illustrated in the SplitsTree (see Fig. 2), the *S. agalactiae* strains investigated
190 in this study also exhibited highly heterogeneous DNA microarray hybridization profiles.
191 With the exception of *S. agalactiae* isolated from fish, no distinct clustering of strains based
192 on host species, geographical origin, or clonal complex assignment could be observed.

193 The prevalence of selected virulence and resistance genes among different host groups
194 is presented in Table 4. Depending on the host, different combinations and variants of the
195 pilus island gene clusters were observed. The *speM* gene encoding exotoxin M was detected
196 in only one isolate (S209, CC19-19), originating from a recto-vaginal swab from a patient in
197 China. With regard to the allelic variants of the alpha-like GBS surface proteins, the allele
198 *alp_rib (R4)* was significantly more prevalent in strains of human origin than in strains from
199 all other sources. The *bac* gene encoding a GBS surface protein was frequently present in
200 isolates from dogs. In addition, the genes of the first pilin gene cluster (*pilA/B/C-I*) were more
201 common in canine GBS isolates, whereas prevalence was low in fish isolates. In contrast, the
202 *pilA/B/C-2b* genes of the second pilin gene cluster were significantly more prevalent in GBS
203 from fish compared with GBS isolated from humans, dogs, and rodents. The vast majority of
204 human isolates (94%) harbored *scpB*, which encodes for C5a peptidase and is used as a
205 diagnostic marker.

206 As for genes conferring resistance to antimicrobial agents, the *emrB/qacA* multidrug
207 resistance transporter gene was present in all tested strains. The majority of strains also
208 exhibited *tetM*, a gene associated with tetracycline resistance, and *cadD*, involved in cadmium
209 resistance. Among human and canine strains, we frequently detected *merA/R*, genes involved
210 in mercuric resistance. Online Resource 2 provides a comprehensive overview of the
211 frequency of all virulence and resistance genes detected among the different host groups, as
212 well as *p*-values for statistically significant differences. Full DNA microarray hybridization
213 patterns of all strains are included in Online Resource 3.

214

215

Discussion

216 To date, data on GBS interspecies transmission is limited. In particular, the zoonotic
217 potential and the directionality of transmission of GBS infections are poorly understood.

218 Experimental studies showed the transmissibility of various bovine and human GBS strains to
219 fish [26–28] and characterization and genotyping studies suggested occasional transmission
220 between humans and cattle [23,24]. Very recently, transmission of *S. agalactiae* through
221 ingestion of raw fish sushi was reported to have led to severe infections in humans
222 (Kalimuddin et al., 2017). In addition, cases of GBS infections acquired through contact with
223 GBS from other host species have been reported: necrotizing fasciitis and endocarditis cases
224 in humans occurred after a dog [25] and a cat bite [21], respectively, and necrotizing fasciitis
225 cases in a group of crocodiles were likely of human origin [22].

226 In our study, isolates from various hosts were assigned to the same strain, suggesting
227 interspecies transmission. Five GBS strains were detected in at least one bovine and one
228 human host, and another strain was detected in a human, a bovine, and two canine hosts. In
229 addition, a canine and a porcine isolate were assigned to the same strain. The relatively high
230 number of *S. agalactiae* strains identified in both a sample from a dog and at least one other
231 host species is particularly striking, considering that only 15 canine strains were included in
232 this study. However, the data provided in this study does not allow for conclusions regarding
233 the directionality of transmission. In addition, it needs to be taken into consideration that the
234 strain collection tested predominantly comprises human and bovine GBS strains originating
235 from Europe, which may bias results.

236 Nitschke et al. [13] introduced GBS typing based on DNA microarray hybridization
237 patterns and provided data on human GBS from Germany and the Caribbean, as well as
238 bovine GBS from Germany: The most prevalent hybridization patterns detected were HP-01
239 (CC19-01), HP-30 (CC19-17), HP-35 (CC19-19), and HP-48 (CC23), corresponding to the
240 whole-genome sequenced reference strains CJB111, COH1, Gottschalk 1003A, and Strain
241 515, respectively. All four hybridization patterns were also frequently detected in our study,
242 with HP-01 being linked to the most diverse set of hosts. GBS of HP-01 originated from

243 humans (n = 5), cows (n = 3), dogs (n = 2), mice (n = 3), emerald monitors (n = 2), a rat (n =
244 1), and a snake (n = 1). GBS of HP-30 originated from human hosts (n = 10), a rabbit (n = 1),
245 a cow (n = 1), and a goat (n = 1). GBS of HP-35 originated from humans (n = 8), a dog (n =
246 1), and a cow (n = 1), and GBS of HP-48 were detected in human (n = 15), bovine (n = 3),
247 and canine (n = 2) hosts.

248 The versatility and wide spread of these strains becomes evident, when considering the hosts
249 and geographical locations, in which some of the strains investigated in this study were
250 isolated: S60/S250/S256 (HP-01) was detected in a sample from the skin of a dog in Germany,
251 as well as in a human vaginal swab from China, and bovine mastitis milk in Germany.
252 S117/S254 (HP-30) was identified in a sample from a rabbit in Germany, as well as in human
253 samples in Germany and Colombia. S185/S255 (HP-35) was detected in a sample from the
254 paw of a dog in Germany, and vaginal swabs from women in Colombia and Switzerland.

255 This study provides comprehensive data on the occurrence of capsular serotypes
256 among human and animal GBS isolates. CPS typing data is not only essential for
257 epidemiological purposes, but is also needed in the development of effective CPS-based
258 vaccines [11,12,32].

259 Among the GBS strains investigated in this study, we frequently detected genes
260 conferring resistance to antimicrobial agents and heavy metal resistance markers. Genes
261 associated with macrolide/ clindamycin resistance were exclusively found among GBS from
262 humans, ruminants, dogs, and a pig. Various recent studies report that 15-21% of GBS strains
263 isolated from pregnant women or cases of neonatal GBS infections are resistant to macrolide
264 and/or lincosamide [33–35]. The high prevalence of *tetM* detected in our study in human
265 (76%) and ruminant (48%) strains is consistent with findings of Nitschke and colleagues,
266 which reported prevalence rates of 78% and 71% in human GBS from Germany and the
267 Caribbean, as well as 48% in bovine GBS from Germany [13].

268 In our study, 40% of the canine strains and 25% of fish strains exhibited *bac*, while the
269 gene was only detected in 13% of GBS strains from human origin. The *bac* gene encodes the
270 C protein beta antigen (Bac), which is able to simultaneously bind to the Fc fragment of IgA
271 and the complement regulator factor H, thus likely contributing to immune evasion [32,36]. In
272 addition, increased Bac expression was reported in invasive strains compared to strains
273 collected from vaginal carriers [37]. Previous studies have associated *bac* sequence types with
274 capsular serotype assignment [37,38]. In contrast to our findings, a study investigating human
275 GBS from Asia, Australia, Europe, New Zealand, and North America found that *bac* was
276 present in 97% of serotype Ib isolates and 37% of serotype II isolates, while being largely
277 absent in GBS assigned to other serotypes [38].

278 Low prevalence of the *speM* gene encoding exotoxin M has been reported among GBS
279 from human and bovine sources [13]. This is consistent with our findings. In this study, we
280 detected *speM* in only one isolate (S209, CC19-19) originating from a recto-vaginal swab
281 from a patient in China.

282 In our study, the alpha-like GBS surface protein allele *alp_rib* (*R4*) (= R4, rib) was
283 significantly more prevalent in strains of human origin than in strains from all other sources.
284 The alpha-like proteins are chimeras forming mosaic structures on the surface of the organism
285 [39]. While the function of many alpha-like proteins is still poorly understood, they may act
286 as invasins mediating adherence to cervical epithelial cells, as well as transmembrane passage
287 and translocation of the organism [39].

288 In our study, different hosts were associated with different combinations and allelic
289 variants of genes of the pilus islands. Each of the three pilus islands (PI-1, PI-2a, PI2b)
290 encodes one backbone and two ancillary proteins that mediate interactions with host cells. The
291 pilus islands and their combinations were shown to play an important role in host adaptation
292 and specificity, as well as disease presentation [40].

293 The findings of our study suggest that while *S. agalactiae* is well adapted to various hosts
294 including humans, cattle, dogs, rodents, and fish, interspecies transmission is possible and
295 occurs amongst others between humans and cows, dogs, and rabbits. Involvement of a canine
296 host in interspecies transmission events may be particularly frequent, with the directionality of
297 transmission still being unclear. The virulence and resistance gene patterns determined in our
298 study significantly extend the limited current knowledge on interspecies transmission. They
299 could also be utilized in the identification of suitable targets for therapeutic agents, as well as
300 vaccines.

301

302

Acknowledgements

303 We thank the Institute of Medical Microbiology, University of Zurich, for providing strains.

304

305

Funding

306 All materials used for this study were funded by the Institute for Food Safety, University of
307 Zurich. No third party funding or external grants were involved.

308

309

Conflict of Interest

310 The authors declare that the research was conducted in the absence of any commercial or
311 financial relationships that could be construed as a potential conflict of interest.

312

313

Ethical Approval

314 This study was carried out in accordance with ethical clearance and informed consent
315 regulations of the locally cognizant ethics commission. All isolates were part of existing strain
316 collections with anonymized sample information. No animal or human hosts were subjected
317 to sampling for the purpose of the present study.

318

319

Informed consent

320 This was a retrospective study. For this type of study formal consent is not required.

321 **References**

- 322 1. Krohn MA, Hillier SL, Baker CJ. Maternal peripartum complications associated with
323 vaginal Group B *Streptococci* colonization. J Infect Dis 1999;179:1410–1415
- 324 2. Farley M, Harvey C, Stull T et al. A population-based assessment of invasive disease
325 due to group B *Streptococcus* in nonpregnant adults. N Engl J Med 1993;328:1807–
326 1811
- 327 3. Jackson LA, Hilsdon R, Farley MM et al. Risk factors for group B streptococcal
328 disease in adults. Ann Intern Med 1995;123:415–420
- 329 4. van der Mee-Marquet N, Domelier A-S, Salloum M et al. Molecular characterization of
330 temporally and geographically matched *Streptococcus agalactiae* strains. Foodborne
331 Pathog Dis 2009;6:1177–1183
- 332 5. Foxman B, Gillespie BW, Manning SD, Marrs CF. Risk factors for group B
333 streptococcal colonization: potential for different transmission systems by capsular type.
334 Ann Epidemiol 2007;17:854–862
- 335 6. Hetzel U, König A, Yildirim AÖ, Lämmle C, Kipar A. Septicaemia in emerald
336 monitors (*Varanus prasinus* Schlegel 1839) caused by *Streptococcus agalactiae*
337 acquired from mice. Vet Microbiol 2003;95:283–293
- 338 7. Duremdez R, Al-Marzouk A, Qasem JA, Al-Harbi A, Gharabally H. Isolation of
339 *Streptococcus agalactiae* from cultured silver pomfret, *Pampus argenteus* (Euphrasen),
340 in Kuwait. J Fish Dis 2004;27:307–10
- 341 8. Barkema HW, Green MJ, Bradley AJ, Zadoks RN. Invited review: The role of
342 contagious disease in udder health. J Dairy Sci 2009;92:4717–4729
- 343 9. Keefe G. Update on control of *Staphylococcus aureus* and *Streptococcus agalactiae* for
344 management of mastitis. Vet Clin North Am - Food Anim Pract 2012;28:203–216
- 345 10. Le Doare K, Heath PT. An overview of global GBS epidemiology. Vaccine

- 346 2013;31:D7–12
- 347 11. Johri AK, Paoletti LC, Glaser P, Dua M, Sharma PK, Grandi G, et al. Group B
348 *Streptococcus*: global incidence and vaccine development. Nat Rev Microbiol
349 2006;4:932–942
- 350 12. Nuccitelli A, Rinaudo CD, Maione D. Group B *Streptococcus* vaccine: state of the art.
351 Ther Adv Vaccines 2015;3:76–90
- 352 13. Nitschke H, Slickers P, Müller E, Ehricht R, Monecke S. DNA microarray-based
353 typing of *Streptococcus agalactiae*. J Clin Microbiol 2014;52:3933–43
- 354 14. Yao K, Poulsen K, Maione D et al. Capsular gene typing of *Streptococcus agalactiae*
355 compared to serotyping by latex agglutination. J Clin Microbiol 2013;51:503–507.
- 356 15. Yildirim AÖ, Lämmler C, Weiß R. Identification and characterization of *Streptococcus*
357 *agalactiae* isolated from horses. Vet Microbiol 2002;85:31–35
- 358 16. Skov Sørensen UB, Poulsen K, Ghezzi C, Margarit I, Kilian M. Emergence and global
359 dissemination of host-specific *Streptococcus agalactiae* clones. MBio 2010;1:1–9.
- 360 17. Delannoy CM, Crumlish M, Fontaine MC et al. Human *Streptococcus agalactiae*
361 strains in aquatic mammals and fish. BMC Microbiol 2013;13:41
- 362 18. Ip M, Cheuk ESC, Tsui MHY, Kong F, Leung TN, Gilbert GL. Identification of a
363 *Streptococcus agalactiae* serotype III subtype 4 clone in association with adult invasive
364 disease in Hong Kong. J Clin Microbiol 2006;44:4252–4254
- 365 19. Belard S, Toepfner N, Capan-Melser M et al. *Streptococcus agalactiae* serotype
366 distribution and antimicrobial susceptibility in pregnant women in Gabon, Central
367 Africa. Sci Rep 2015;5:17281
- 368 20. Manning SD, Springman AC, Million AD et al. Association of group B *Streptococcus*
369 colonization and bovine exposure: a prospective cross-sectional cohort study. PLoS
370 One 2010;5

- 371 21. Ashrafian H, Griselli M, Rubens MB, Mullen MJ, Sethia B. Pulmonary Homograft
 372 Endocarditis 19 Years after a Ross Procedure. *Thorac Cardiovasc Surg* 2007;55:55–56
- 373 22. Bishop EJ, Shilton C, Benedict S et al. Necrotizing fasciitis in captive juvenile
 374 *Crocodylus porosus* caused by *Streptococcus agalactiae*: an outbreak and review of the
 375 animal and human literature. *Epidemiol. Infect.* 2007;135:1248–1255
- 376 23. Dogan B, Schukken YH, Santisteban C, Boor J, Boor KJ. Distribution of serotypes and
 377 antimicrobial resistance genes among *Streptococcus agalactiae* isolates from bovine
 378 and human hosts. *Society* 2005;43:5899–5906
- 379 24. Oliveira ICM, Mattos MC De, Pinto TA, Benchetrit LC. Genetic relatedness between
 380 group B *Streptococci* originating from bovine mastitis and a human group B
 381 *Streptococcus* type V cluster displaying an identical pulsed-field gel electrophoresis
 382 pattern. *Clin Microbiol Infect* 2006;12:887–893
- 383 25. Lee S, Roh KH, Kim CK et al. A case of necrotizing fasciitis due to *Streptococcus*
 384 *agalactiae*, *Arcanobacterium haemolyticum*, and *Fingoldia magna* in a dog-bitten
 385 patient with diabetes. *Korean J Lab Med* 2008;28:191–195
- 386 26. Chen M, Wang R, Luo FG et al. *Streptococcus agalactiae* isolates of serotypes Ia, III
 387 and V from human and cow are able to infect tilapia. *Vet Microbiol* 2015;180:129–135
- 388 27. Evans JJ, Klesius PH, Pasnik DJ, Bohnsack JF. Human *Streptococcus agalactiae*
 389 isolate in Nile tilapia (*Oreochromis niloticus*). *Emerg Infect Dis* 2009;15:774–776
- 390 28. Pereira UP, Mian GF, Oliveira ICM, Benchetrit LC, Costa GM, Figueiredo HCP.
 391 Genotyping of *Streptococcus agalactiae* strains isolated from fish, human and cattle
 392 and their virulence potential in Nile tilapia. *Vet. Microbiol.* 2010;140:186–192
- 393 29. Coombs GW, Monecke S, Ehrlich R et al. Differentiation of clonal complex 59
 394 community-associated methicillin-resistant *Staphylococcus aureus* in Western
 395 Australia. *Antimicrob Agents Chemother* 2010;54:1914–1921.

- 396 30. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies.
397 Mol Biol Evol 2006;23:254–267
- 398 31. Wattering L, Stephan R, Layer F, Johler S. Comparison of *Staphylococcus aureus*
399 isolates associated with food intoxication with isolates from human nasal carriers and
400 human infections. Eur J Clin Microbiol Infect Dis 2012;31:455–464
- 401 32. Lindahl G, Stålhammar-Carlemalm M, Areschoug T. Surface proteins of *Streptococcus*
402 *agalactiae* and related proteins in other bacterial pathogens. Clin Microbiol Rev
403 2005;18:102–127
- 404 33. Bolukaoto JY, Monyama CM, Chukwu MO et al. Antibiotic resistance of
405 *Streptococcus agalactiae* isolated from pregnant women in Garankuwa, South Africa.
406 BMC Res Notes 2015;8:364
- 407 34. Creti R, Imperi M, Berardi A et al. Neonatal group B *Streptococcus* infections. Pediatr
408 Infect Dis J 2017;36:256–262
- 409 35. Rojo-Bezares B, Azcona-Gutiérrez JM, Martín C, Jareño MS, Torres C, Sáenz Y.
410 *Streptococcus agalactiae* from pregnant women: antibiotic and heavy-metal resistance
411 mechanisms and molecular typing. Epidemiol Infect 2016;144:3205–3214
- 412 36. Areschoug T, Linse S, Stålhammar-Carlemalm M, Hedén LO, Lindahl G. A proline-
413 rich region with a highly periodic sequence in streptococcal beta protein adopts the
414 polyproline II structure and is exposed on the bacterial surface. J Bacteriol
415 2002;184:6376–6383
- 416 37. Nagano N, Nagano Y, Taguchi F. High expression of a C protein beta antigen gene
417 among invasive strains from certain clonally related groups of type Ia and Ib group B
418 *Streptococci*. Infect Immun 2002;70:4643–4649
- 419 38. Kong F, Gidding HF, Berner R, Gilbert GL. *Streptococcus agalactiae* C β protein gene
420 (*bac*) sequence types, based on the repeated region of the cell-wall-spanning domain:

- 421 Relationship to virulence and a proposed standardized nomenclature. *J Med Microbiol*
422 2006;55:829–837
- 423 39. Maeland JA, Afset JE, Lyng R V., Radtke A. Survey of immunological features of the
424 alpha-like proteins of *Streptococcus agalactiae*. *Clin Vaccine Immunol* 2015;22:153–
425 159
- 426 40. Springman A, Lacher DW, Waymire EA et al. Pilus distribution among lineages of
427 group b *Streptococcus*: an evolutionary and clinical perspective. *BMC Microbiol*
428 2014;14:159
429
430

431 **Figure legends**

432

433 **Fig 1 Interspecies transmission.** Several *S. agalactiae* strains were detected in samples from
434 more than one host species, indicating interspecies transmission. This figure provides an
435 overview of the links detected and their frequency.

436

437 **Fig 2 SplitsTree.** SplitsTree illustrating the degree of similarity of virulence and resistance
438 gene profiles of *S. agalactiae* strains from different sources: Human host (pink), ruminant
439 (green), dog (orange), elephant (grey), fish (blue), rodent/rabbit (yellow), other (purple).
440 Strains detected in two or more host species are marked by red circles.

441

442

Tables

443

444 **Table 1: Clonal lineages and strains identified in more than one continent and across**
 445 **multiple host species.** In some clonal complexes, strains were isolated more than once, some
 446 of them beyond country borders and from different host species.

Clonal complex	Strain	Source	Sample	Country^a
CC19-01	S48/S244/S245	Rat (n = 1)	Abscess	CH
		Monitor (n = 2)	Lung/ kidney/ liver/ intestine	DE
		Mouse (n = 3)	Intestine	DE
	S53	Mouse (n = 5)	Intestine	DE
	S57/ S249	Snake (n = 2)	Liver, skin	DE
		Monitor (n = 1)	Liver	DE
	S60/S250/S256	Dog (n = 2)	Skin	DE
		Human (n = 4)	Vaginal swab	HK
		Bovine (n = 2)	Milk	DE
	S61/S251	Rat (n = 1)	Trachea	DE
		Mouse (n = 2)	Prepuce	DE
	S63	Bovine (n = 4)	Milk	DE
	S64	Bovine (n = 3)	Milk	DE
	S65	Bovine (n = 2)	Milk	DE
	S69	Bovine (n = 5)	Milk	DE
	S81	Human (n = 2)	Vaginal swab	HK
	S84	Human (n = 2)	Vaginal swab	HK

Morach 2017

	S92	Human (n = 2)	Vaginal swab	HK
	S102	Human (n = 2)	Vaginal swab, abdominal tissue	HK, KY
CC19-02	S3	Guinea pig (n = 2)	Nose, liver	DE
	S7	Bovine (n = 2)	Milk	DE
CC19-10	S58/S252	Bovine (n = 1)	Organs	DE
		Human (n = 3)	Urine, vaginal swab, wound	CO, CH, KY
	S66	Bovine (n = 2)	Uterus, milk	DE
	S68	Bovine (n = 3)	Milk	DE
	S73	Human (n = 2)	Pus, urine	CO
	S85	Human (n = 2)	Vaginal swab	HK
	S90	Tilapia (n = 4)	Kidney	TH
	S91	Tilapia (n = 2)	Kidney	TH, VN
	S112	Human (n = 2)	Urine, blood	KY
CC19-17	S116	Human (n = 4)	Mastitis, blood, vaginal swab	DE, CO, CH
	S117/S254	Rabbit (n = 1)	Unknown	DE
		Human (n = 4)	Vaginal swab, urine	CO, CH
	S120	Elephant (n = 3)	Abscess/ foot	DE
	S126	Bovine (n = 2)	Milk	DE
	S152	Human (n = 3)	Vaginal swab	HK
	S153	Human (n = 2)	Vaginal swab	CH
	S157	Human (n = 2)	Vaginal swab	CH
	S169	Human (n = 4)	Vaginal swab	CH
	S175	Human (n = 4)	Blood, urine, vaginal swab	KY

Morach 2017

CC19-19	S186/S255	Dog (n = 1)	Paw	DE	
		Human (n = 4)	Vaginal swab, urine	CO, CH	
	S190	Human (n = 2)	Urine, vaginal swab	CO	
	S193	Human (n = 2)	Urine	CO	
	S195	Human (n = 3)	Urine, vaginal swab	CO	
	S197	Human (n = 2)	Vaginal swab	CO	
	S198	Human (n = 2)	Urine, blood	CO	
	S218	Human (n = 2)	Vaginal swab	CH	
	S222	Human (n = 2)	Vaginal swab	CH, KY	
	S227	Human (n = 3)	Vaginal swab	CH	
	S235	Human (n = 2)	Vaginal swab	CH	
	S237	Human (n = 2)	Blood, vaginal swab	KY, CO	
	CC19-67	S5/S243	Dog (n = 1)	Skin	DE
			Bovine (n = 1)	Milk	CH
S17		Bovine (n = 4)	Milk	CO	
S23		Bovine (n = 2)	Milk	CH	
CC23	S124/248	Dog (n = 1)	Skin	DE	
		Pig (n = 1)	Milk	DE	
	S128	Bovine (n = 2)	Milk	DE	
	S130	Bovine (n = 2)	Milk	DE	
	S133	Bovine (n = 3)	Milk	DE	
	S134/S253	Bovine (n = 1)	Milk	DE	
		Human (n = 3)	Urine, vaginal swab	CO, HK	
S135	Bovine (n = 2)	Milk	DE		

Morach 2017

	S137	Human (n = 12)	Vaginal swab, biopsy, urine, blood	CO, CH, KY
	S139	Human (n = 3)	Urine, blood, secretion	CO
	S141	Human (n = 2)	Vaginal swab, urine	CO
	S142	Human (n = 2)	Vaginal swab	CO
	S145	Human (n = 3)	Vaginal swab	CH, CO
	S162	Human (n = 3)	Vaginal swab	CH
CC103	S11/S247	Bovine (n = 1)	Milk	DE
		Human (n = 1)	Pus	CO
	S14	Bovine (n = 5)	Milk	DE
	S16/S246	Bovine (n = 1)	Milk	DE
		Human (n = 1)	Urine	CO
CC260/261	S31	Tilapia (n = 2)	Spleen, kidney	HN, CO
	S32	Tilapia (n = 3)	Spleen, kidney	HN, CO
CC298	S19	Bovine (n = 3)	Milk	CO
not assigned	S10	Bovine (n = 2)	Milk	DE
	S18	Bovine (n = 2)	Milk	CO

447 ^a Country abbreviations: CH = Switzerland, CO = Colombia, DE = Germany, HK = Hong

448 Kong SAR (China), HN = Honduras, KY = Kenya, TH = Thailand, VN = Vietnam

449

450

451

452

453

454 **Table 2. Clonal complex distribution.** This table provides an overview of the prevalence of
 455 different clonal complexes among *S. agalactiae* strains from various hosts (in percent).

Clonal complex	Hosts (% of strains)					
	Human (n = 161)	Ruminant (n = 52)	Dog (n = 15)	Rodent (n = 8)	Fish (n = 8)	Other (n = 12)
CC19-01	12	19	13	75	0	25
CC19-02	4	2	7	25	0	0
CC19-04	1	0	0	0	0	0
CC19-10	12	12	40	0	25	17
CC19-17	10	6	0	0	0	17
CC19-19	35	4	13	0	0	0
CC19-22	2	0	0	0	0	0
CC19-67	1	13	7	0	0	8
CC23	20	21	20	0	0	33
CC26	1	0	0	0	0	0
CC103	2	10	0	0	0	0
CC130	1	0	0	0	0	0
CC260/261	0	0	0	0	75	0
CC298	0	2	0	0	0	0
not assigned	0	12	0	0	0	0

456

457 **Table 3: Prevalence of capsular serotypes.**

Capsular serotype	Hosts (% of strains)					
	Human (n = 161)	Ruminant (n = 52)	Dog (n = 15)	Rodent (n = 8)	Fish (n = 8)	Other (n = 12)
IA	16	35	20	0	13	25
IB	9	8	7	0	63	0
II	17	19	20	50	0	0
III	22	21	13	0	13	33
IV	4	10	13	0	0	0
V	21	8	20	50	0	25
VI	1	0	0	0	0	0
VII	2	0	0	0	0	0
IX	1	0	0	0	0	0
negative	2	0	0	0	13	17
not assignable	5	0	7	0	0	0

458

Table 4. Virulence and resistance genes. Prevalence of selected virulence and resistance genes among GBS strains isolated from different hosts: humans, ruminants, dogs, rodents, fish, and other (snake, turtle, frog, elephant, pig, rabbit). A comprehensive list of DNA microarray results including *p*-values is provided as Supplementary Table 2.

Gene	Function	Host (% of strains)					
		Human (n = 161)	Ruminant (n = 52)	Dog (n = 15)	Rodent (n = 8)	Fish (n = 8)	Other (n = 12)
Virulence genes							
<i>speM</i>	Exotoxin M	1	0	0	0	0	0
<i>cylD</i>	Beta hemolysin locus	96 ^{*F}	100 ^{*F}	100 ^{*F}	100 ^{*F}	25 ^{*HRDXY}	100 ^{*F}
<i>cylE</i>	Beta hemolysin locus	87 ^{*F}	94 ^{*F}	100 ^{*F}	100 ^{*F}	25 ^{*HRDXY}	100 ^{*F}
<i>alp_3</i>	Allele of the α -like protein/ α -antigenic cell wall protein	7 ^{*X}	10 ^{*X}	13 ^{*X}	75 ^{*HRDF}	0 ^{*X}	25
<i>alp_rib (R4)</i>	Allele of the α -like protein/ α -antigenic cell wall protein	52 ^{*RXFY}	24 ^{*H}	20	0 ^{*H}	0 ^{*H}	9 ^{*H}
<i>bac</i>	β -antigenic cell wall protein	13 ^{*D}	15	40 ^{*H}	0	25	8
<i>pilA1</i>	Pilin gene cluster 1	51	50	80 ^{*F}	71	25 ^{*D}	50

Morach 2017

<i>pilB1</i>	Pilin gene cluster 1	63	48 ^{*D}	80 ^{*RF}	75	25 ^{*D}	50
<i>pilC1</i>	Pilin gene cluster 1	66	56	80	75	38	50
<i>pilA2a</i>	Pilin gene cluster 2a	81 ^{*RFY}	52 ^{*HDX}	87 ^{*RF}	100 ^{*RFY}	13 ^{*HDX}	50 ^{*HX}
<i>pilC2a</i>	Pilin gene cluster 2a	82 ^{*RF}	52 ^{*HDX}	93 ^{*RF}	100 ^{*RF}	13 ^{*HDX}	58
<i>pilA2b</i>	pilin gene cluster 2b	15 ^{*RFY}	48 ^{*HDX}	7 ^{*RF}	0 ^{*RF}	63 ^{*HDX}	42 ^{*H}
<i>pilB2b</i>	Pilin gene cluster 2b	15 ^{*RFY}	48 ^{*HDX}	7 ^{*RF}	0 ^{*RF}	67 ^{*HDX}	42 ^{*H}
<i>pilC2b</i>	Pilin gene cluster 2b	14 ^{*RFY}	48 ^{*HDX}	7 ^{*RF}	0 ^{*RF}	75 ^{*HDX}	42 ^{*H}
<i>scpB-var1</i>	Complement-inactivating C5a peptidase	94 ^{*RDXFY}	50 ^{*H}	67 ^{*HF}	25 ^{*H}	13 ^{*HD}	27 ^{*H}
<i>scpB-var2</i>	Complement-inactivating C5a peptidase	94 ^{*RDXFY}	48 ^{*H}	67 ^{*HF}	25 ^{*H}	13 ^{*HD}	25 ^{*H}
<i>fsb-var3</i>	Allele of a fibrinogen binding protein	61 ^{*X}	46 ^{*X}	73 ^{*F}	100 ^{*HRFY}	25 ^{*DX}	33 ^{*X}

Resistance genes

<i>cadC</i>	Cadmium efflux system accessory protein	21 ^{*R}	2 ^{*H}	13	0	0	0
<i>cadD</i>	Cadmium resistance protein	75 ^{*F}	77 ^{*F}	93 ^{*FY}	100 ^{*FY}	14 ^{*HRDX}	50 ^{*DX}
<i>emrB/qacA</i>	Multidrug resistance transporter	100	100	100	100	100	100
<i>ermA</i>	Macrolide/clindamycin resistance	9	2	13	0	0	8

Morach 2017

<i>ermB</i>	Macrolide/clindamycin resistance	19	10	7	0	0	0
<i>ermC</i>	Macrolide/clindamycin resistance	0	0	0	0	0	0
<i>merA</i>	Mercuric reductase	58 ^{*RXFY}	11 ^{*HD}	45 ^{*R}	0 ^{*H}	13 ^{*H}	10 ^{*H}
<i>merR</i>	Mercuric resistance operon regulatory protein	57 ^{*RXFY}	12 ^{*H}	33	0 ^{*H}	13 ^{*H}	17 ^{*H}
<i>tetM</i>	Tetracycline resistance	76 ^{*RF}	48 ^{*HD}	93 ^{*RF}	75	25 ^{*HD}	58

*The distribution of the respective gene differed significantly between strains from the stated hosts (with $p \leq 0.050$). Host groups are indicated as follows:

humans (H), ruminants (R), dogs (D), rodents (X), fish (F), and other (Y).

459

460

461