

CROP FERTILIZATION ON COAST PRAIRIE AND COASTAL BEND SOILS

Charles D. Welch, Carl Gray, Dwayne Westfall,
and Marvin E. Riewe*

Land resource regions, Figure 1, represent about 11 million acres of forest, crop and rangeland. Annual rainfall ranges from 26 inches in the west to 55 inches in the east. A portion of this region is coastal marsh, poorly drained, with limited productive use.

Soil Characteristics

Soils range from light brown to black and from sands to clay in the surface. Subsoils, generally, are higher in clay but some coastal marsh soils show only slight changes from the surface downward. A few areas have salt accumulations that limit use for crop production. Some soils in the Coastal Bend have caliche outcrops or exposed subsurface layers, creating nutritional as well as management problems often difficult to correct.

Base status of soils varies from acid to alkaline influenced by parent materials and rainfall. Clay

*Respectively, Extension soil chemists, Texas Agricultural Extension Service; associate professor, Texas A&M University Research and Extension Center, Beaumont; and associate professor in charge of Texas A&M University Research Center, Angleton.



Figure 1. Location of Coast Prairie, Coastal Marsh and Coastal Bend Regions.

type varies but montmorillonitic types generally predominate. Such clays are difficult to maintain in a desirable physical condition, often creating management problems.

Soil Fertility Levels

Soil characteristics, past fertilization and cropping practices have resulted in a wide range of soil fertility levels. Soil test summary data, Table 1, show 80 percent of soils in the Coast Prairie and 33 percent in the Coastal Bend low in phosphorus. More Coast Prairie soils are low in potassium than Coastal Bend.

N, P₂O₅ and K₂O for Major Crops

The wide range in soil fertility levels and management practices in this region call for fertilization adapted to specific production requirements. Therefore, soil tests properly calibrated to express available nutrients and correlated with crop response are the best guide to profitable fertilization and liming. Two important criteria needed for selecting the profitable rate of nutrient are: (1) the level of available nutrient in the soil and (2) the expected yield or production goal.

Many soil properties, as well as extractable nutrients, must be evaluated in grouping soils, as a means of expressing the level of available nutrients. Depth of sampling is important especially for perennial sod crops. For established pastures, collect soil to a depth of 3 to 4 inches. See D-494, *Soil Sample Information Sheet for Field Crops*. The expected yield, which expresses potential productivity, includes anticipated moisture and management conditions.

Rates of N, P₂O₅ and K₂O at varying soil test levels and expected yields for major crops are shown in Table 2 through 12. The soil test levels are used by Texas A&M University. To use these tables, determine the soil test level in the left column and read across to the expected yield column for nutrient rate. For example in Table 2 a soil low (L) in nitrogen, low (L) in phosphorus and medium (M) in potassium would show a 60-40-30 for 4,500 pounds of grain sorghum.

Table 1. Percentage distribution of Coast Prairie and Coastal Bend soils in five levels for pH, organic matter, phosphorus and potassium¹

Soil test level	Organic matter		Phosphorus		Potassium		Soil pH range	CP ²	CB ²
	CP ²	CB ²	CP ²	CB ²	CB ²	CB ²			
VL	4	0	61	18	19	1	Below 5.0	1	0
L	16	12	17	15	26	1	5.1 - 5.5	5	0
M	25	45	13	35	22	2	5.6 - 6.0	18	0
H	23	28	5	20	15	31	6.1 - 6.5	28	3
VH	32	15	4	12	18	65	Above 6.5	48	97

¹From soil test summaries, Soil Testing Laboratory, Agricultural Extension Service, Texas A&M University.

²CP—Coast Prairie, CB—Coastal Bend.

Table 2. Application rates of nutrients for grain sorghum—three production levels

Soil test level	Expected yield 3,000 lb/A			Expected yield 4,500 lb/A			Expected yield 6,000 lb/A		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
VL	40	30	30-50	70	50	50-60	100	70	80-100
L	30	20	20-30	60	40	30-50	90	60	60-80
M	20	0	0-20	50	30	20-30	80	50	40-60
H	—	0	0	—	0	0	—	30	30-40
VH	—	0	0	—	0	0	—	0	0

Table 3. Application rates of nutrients for cotton—three production levels

Soil test level	Expected yield 1 bale/A			Expected yield 1 1/2 bale/A			Expected yield 2 bale/A		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
VL	60	60	60-80	80	80	80-100	100	100	100-120
L	40	50	40-60	60	70	60-80	80	90	80-100
M	20	40	30-50	40	50	40-60	60	70	60-80
H	—	0	0	—	0	0	—	0	0
VH	—	0	0	—	30	20-40	—	60	30-60

Table 4. Application rates of nutrients for rice—two levels of management

Soil test level	Regular varieties			High N varieties			For second crop		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
VL	100	40	40-60	120	40	40-60	60	0	0
L	80	40	30-50	100	40	30-50	50	0	0
M	60	20	20-30	80	20	20-30	40	0	0
H	—	0	0	—	0	0	—	0	0
VH	—	0	0	—	0	0	—	0	0

Table 5. Application rates of nutrients for corn—three production levels

Soil test level	Expected yield ¹ 60-70 bu/A			Expected yield ² 90-100 bu/A			Expected yield ³ 120-130 bu/A		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
VL	80	50	50-70	120	80	80-100	160	120	100-120
L	60	40	30-50	100	70	60-80	140	100	80-100
M	40	30	20-30	80	60	40-60	120	80	60-80
H	—	0	0	—	30	30-40	—	40	30-60
VH	—	0	0	—	0	0	—	0	0

¹Plant population 8,000-10,000 plants per acre.

²Plant population 12,000-14,000 plants per acre.

³Plant population 16,000-18,000 plants per acre.

Calcium, Magnesium and Sulfur

Soils in the Coast Prairie vary in base status. However, soil test summary data in Table 1 show 52 percent samples tested below pH 6.5. Coastal Bend soils generally are less acid with only 3 percent of the samples below pH 6.5.

Sufficient samples have not been analyzed to indicate the extent of magnesium deficiency in

these regions, but it probably is greatest in the more acid sandy soils. Use of dolomitic limestone, containing at least 10 percent magnesium carbonate, is the most economical way to apply magnesium.

Sulfur needs and response in the Coastal region are being studied, but a general need is not anticipated because of industrial wastes and other means by which sulfur is added to the soils.

Table 6. Application rates of nutrients for ryegrass, oats and similar winter grasses (no legume)

Soil test level	2 tons/A ¹			3 tons/A ¹		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
VL	80	40	30-50	120	80	90-110
L	60	30	20-40	100	60	70-90
M	40	20	20-30	80	40	50-70
H	—	0	0	—	0	0
VH	—	0	0	—	0	0

¹Production from Nov. 15 through April 15. An additional 50 lb. of N can be used to extend ryegrass grazing into May.

Table 7. Application rates of nutrients for establishing and maintaining S₁ Louisiana white clover with grass

Soil test level	At or before seeding			Maintenance for grazing		
	N	P ₂ O ₅ ¹	K ₂ O ²	N ³	P ₂ O ₅	K ₂ O
VL	30	180	180-220	0	100	180-220
L	25	140	120-160	0	75	130-150
M	20	100	90-110	0	50	80-120
H	—	60	30-50	—	30	50-70
VH	—	0	0	—	0	0

¹If these rates are applied, the phosphorus level should be medium or above for maintenance.

²May apply half the spring following seeding.

³Nitrogen topdressing may be needed in mid-season, depending on management, grazing requirements and clover growth.

Table 8. Application rates of nutrients for common bermuda, Dallis and similar summer grasses—two grazing intensities (no legume)¹

Soil test level	2 tons/A ²			4 tons/A ²		
	N ²	P ₂ O ₅	K ₂ O	N ²	P ₂ O ₅	K ₂ O
VL	80	40	30-50	180	80	90-110
L	60	30	20-40	160	60	70-90
M	40	20	0-30	120	40	50-70
H	—	0	0	—	30	0
VH	—	0	0	—	0	0

¹With a legume such as vetch, peas or clover apply P₂O₅ and K₂O in the fall and delay nitrogen until that furnished by the legume has been used. Increase rates of P₂O₅ and K₂O 50% if clover is planted in fall.

²Production between April 15 and October 15.

Table 9. Application rates of nutrients for establishing coastal bermudagrass

Soil test level	At sprigging			First Summer		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
VL	40	80	70-90	40	0	40-60
L	30	60	60-80	30	0	30-50
M	30	40	50-70	30	0	0
H	—	0	0	—	0	0
VH	—	0	0	—	0	0

Table 10. Application rates of nutrients for 3 levels of coastal bermudagrass hay production (no legume)

Soil test level	Expected yield 6 tons/A			Expected yield 8 tons/A			Expected yield 10 tons/A		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
VL	280	80	180-220	400	100	280-320	550	120	380-420
L	220	60	120-160	320	80	220-260	480	100	320-360
M	180	40	70-90	280	60	160-200	360	80	240-280
H	—	0	0	—	40	80-120	—	60	180-220
VH	—	0	0	—	0	0	—	40	80-120

Table 11. Application rates of nutrients for annual summer forages—three production levels

Soil test level	Expected yield 3 to 4 tons/A			Expected yield 5 to 6 tons/A			Expected yield 9 to 10 tons/A		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
VL	120	60	80-120	220	80	130-150	300	100	160-200
L	100	40	60-100	200	60	100-140	280	80	120-160
M	80	30	40-80	180	40	60-100	260	60	100-120
H	—	0	0	—	20	30-50	—	40	40-80
VH	—	0	0	—	0	0	—	0	0

Table 12. Application rates for soybeans

Soil test level	For the production of from 35 to 40 bu/A		
	N ¹	P ₂ O ₅	K ₂ O
VL	0	60	90-110
L	0	40	60-80
M	0	20	30-40
H	—	0	0
VH	—	0	0

¹Inoculated soybeans are able to obtain nitrogen from the air. However, on new fields where soybeans have never been grown, up to 20 to 30 pounds of N per acre may be included in the fertilizer application to get the crop established until good inoculation is attained.

Micronutrients

The micronutrient group includes seven elements—iron, zinc, manganese, copper, boron, molybdenum and chlorine. Although general micronutrient deficiencies are not confirmed, localized problems with zinc and iron have been encountered. More information about iron and zinc is available in Extension Leaflets L-721 and L-723, available from your county agricultural agent.

On alkaline soils, rice may respond to applications of zinc or iron or both. Due to the tendency for flooding to affect soil pH, problems may be encountered under rice culture even though other crops are not affected. A soil test will provide information about pH and level of extractable iron and zinc. However, results must be interpreted to reflect the effects of flooding.

If a rice grower decides to use zinc he can apply about 10 lbs. of zinc sulfate or equivalent per acre (see L-721 and L-783). A possible iron application may be 250 lbs. of ferric (iron) sulfate per acre (see L-723 and L-783). Other sources of either zinc or iron can be used and have produced satisfactory results.

Conversion Factor

The principle involved in using micronutrients is the same as for other nutrients. That is, identify

and confirm the need, then apply amounts sufficient to meet the production requirement.

Fertilizers are labeled as percent P₂O₅ and K₂O, and soil test values are reported in these terms. However, plant analyses results are usually reported as percentages of the element. For this reason, the following factors are presented for use in converting from one form to the other.

From P ₂ O ₅ to P	multiply by .44
From P to P ₂ O ₅	multiply by 2.3
From K ₂ O to K	multiply by .83
From K to K ₂ O	multiply by 1.2

LIMING ACID SOILS

The soil pH should be known before liming acid soils, as well as the cropping system and soil properties.

Rates of limestone

pH level ¹		Rates in ton/acre ²		
High Ca crops	Low Ca crops	Sands	Sandy loams & loams	Clay & clay loams
6.0-6.3	5.8-6.0	1	1 1/2	2
5.6-5.9	5.4-5.7	1 1/2	2	2 1/2
5.0-5.5	5.0-5.3	2	3	4

¹High calcium crops are legumes and legume grass mixtures. The pH levels under low-calcium crops are for grasses and row crops.

²May be increased 1/2 ton per acre for soils high in montmorillonite.

Magnesium Soil Test

Magnesium is being measured is a routine soil test. The following recommendations are:

Lb/A magnesium ¹	Rating
0-75	Low ²
75-250	Medium
Above 250	High

¹Refers to the soil testing methods and calibrations used by Texas A&M University laboratories.

²Dolomitic limestone containing at least 10% magnesium carbonate should be used for liming soils that are low in this nutrient.

Educational programs conducted by the Texas Agricultural Extension Service serve people of all ages regardless of socio-economic levels, race, color, sex, religion or national origin.

Cooperative Extension Work in Agriculture and Home Economics, The Texas A&M University System and the United States Department of Agriculture cooperating. Distributed in furtherance of the Acts of Congress of May 8, 1914, as amended, and June 30, 1914.