

Applied Mathematical Sciences, Vol. 3, 2009, no. 4, 161 - 164

A New Proof of Watson's Theorem for the Series ${}_3F_2(1)$

Arjun K. Rathie

Manda Institute of Technology
Raisar NH-11, Jaipur Road, Bikaner 334001, India

R. B. Paris

Division of Complex Systems
University of Abertay Dundee, Dundee DD1 1HG, UK
R.Paris@abertay.ac.uk

Abstract

We give a new proof of the classical Watson theorem for the summation of a ${}_3F_2$ hypergeometric series of unit argument. The proof relies on the two well-known Gauss summation theorems for the ${}_2F_1$ function.

Mathematics Subject Classification: 33C20

Keywords: Watson's theorem, hypergeometric series

1. Introduction

The classical Watson theorem for the summation of a ${}_3F_2$ hypergeometric function of unit argument takes the form

$${}_3F_2 \left(\begin{matrix} a, b, c \\ \frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}, 2c \end{matrix}; 1 \right) = \frac{\Gamma(\frac{1}{2}) \Gamma(c + \frac{1}{2}) \Gamma(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}) \Gamma(c + \frac{1}{2} - \frac{1}{2}a - \frac{1}{2}b)}{\Gamma(\frac{1}{2}a + \frac{1}{2}) \Gamma(\frac{1}{2}b + \frac{1}{2}) \Gamma(c + \frac{1}{2} - \frac{1}{2}a) \Gamma(c + \frac{1}{2} - \frac{1}{2}b)} \quad (1.1)$$

provided $\operatorname{Re}(2c - a - b) > -1$ and the parameters are such that the series on the left is defined. The proof of this result when one of the parameters a or b is a negative integer was given by Watson in [7], and subsequently was established more generally in the non-terminating case by Whipple in [8].

The standard proof of the general case given in [2, p. 149; 6, p. 54] relies on the following transformation due to Thomae

$${}_3F_2 \left(\begin{matrix} a, b, c \\ d, e \end{matrix}; 1 \right) = \frac{\Gamma(d)\Gamma(e)\Gamma(s)}{\Gamma(a)\Gamma(b+s)\Gamma(c+s)} {}_3F_2 \left(\begin{matrix} d-a, e-a, s \\ b+s, c+s \end{matrix}; 1 \right),$$

where $s = d + e - a - b - c$ is the parametric excess, combined with Dixon's theorem for the evaluation of the sum on the right when $d = \frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}$ and $e = 2c$. An alternative and more involved proof [4, p. 363] exploits the quadratic transformations for the Gauss hypergeometric function. A third proof, due to Bhatt in [3], exploits a known relation between the F_2 and F_4 Appell functions combined with a comparison of the coefficients in their series expansions.

In this note, we give a simple proof of (1.1) that relies only on the well-known Gauss summation theorems for the ${}_2F_1$ function, namely [1, pp. 556, 557]

$${}_2F_1(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}, \quad \operatorname{Re}(c-a-b) > 0 \quad (1.2)$$

and

$${}_2F_1(a, b; \frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}; \frac{1}{2}) = \frac{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2})}{\Gamma(\frac{1}{2}a + \frac{1}{2})\Gamma(\frac{1}{2}b + \frac{1}{2})}. \quad (1.3)$$

We shall also require the following elementary identities for the Pochhammer symbol, or ascending factorial, $(a)_n = \Gamma(a+n)/\Gamma(a)$ given by

$$(a)_{2m} = 2^{2m}(\frac{1}{2}a)_m(\frac{1}{2}a + \frac{1}{2})_m, \quad (a)_{k+2m} = (a)_{2m}(a+2m)_k \quad (1.4)$$

for nonnegative integers m and k , together with

Lemma 1 *Let k be a nonnegative integer and c be (in general) a complex parameter satisfying $2c \neq -1, -2, \dots$. Then*

$$\frac{(c)_k}{(2c)_k} = \sum_{m=0}^{\lfloor k/2 \rfloor} \frac{2^{-k-2m}k!}{(c + \frac{1}{2})_m m! (k-2m)!}, \quad (1.5)$$

where $\lfloor k/2 \rfloor$ is the integer part of $k/2$.

The proof of this lemma uses (1.2) to express the ratio of Pochhammer symbols as a terminating Gauss hypergeometric function in the form

$$\begin{aligned} \frac{(c)_k}{(2c)_k} &= 2^{-k} {}_2F_1(-\frac{1}{2}k, \frac{1}{2} - \frac{1}{2}k; c + \frac{1}{2}; 1) \\ &= 2^{-k} \sum_{m=0}^{\lfloor k/2 \rfloor} \frac{(-\frac{1}{2}k)_m (\frac{1}{2} - \frac{1}{2}k)_m}{(c + \frac{1}{2})_m m!}. \end{aligned}$$

The result (1.5) then follows upon making use of the identity

$$\left(-\frac{1}{2}k\right)_m \left(\frac{1}{2} - \frac{1}{2}k\right)_m = \frac{2^{-2m} k!}{(k - 2m)!}.$$

2. Proof of Watson's theorem (1.1)

We denote the left-hand side of (1.1) by F and express the ${}_3F_2$ function as a series to find

$$\begin{aligned} F &= \sum_{k=0}^{\infty} \frac{(a)_k (b)_k (c)_k}{\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}\right)_k (2c)_k k!} \\ &= \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}\right)_k k!} \sum_{m=0}^{\lfloor k/2 \rfloor} \frac{2^{-k-2m} k!}{\left(c + \frac{1}{2}\right)_m m! (k - 2m)!} \end{aligned}$$

by Lemma 1. Upon reversal of the order of summation, making use of the easily established result [5, p. 57]

$$\sum_{k=0}^{\infty} \sum_{m=0}^{\lfloor k/2 \rfloor} A(m, k) = \sum_{m=0}^{\infty} \sum_{k=2m}^{\infty} A(m, k) = \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} A(m, k + 2m),$$

then

$$\begin{aligned} F &= \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \frac{(a)_{k+2m} (b)_{k+2m} 2^{-k-4m}}{\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}\right)_{k+2m} \left(c + \frac{1}{2}\right)_m m! k!} \\ &= \sum_{m=0}^{\infty} \frac{(a)_{2m} (b)_{2m} 2^{-4m}}{\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}\right)_{2m} \left(c + \frac{1}{2}\right)_m m!} \sum_{k=0}^{\infty} \frac{(a + 2m)_k (b + 2m)_k 2^{-k}}{\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2} + 2m\right)_k k!} \quad (2.1) \end{aligned}$$

by the second equation in (1.4).

The inner sum in (2.1) can be expressed as a ${}_2F_1$ function in the form

$${}_2F_1 \left(\begin{matrix} a + 2m, b + 2m \\ \frac{1}{2}a + \frac{1}{2}b + \frac{1}{2} + 2m \end{matrix} ; \frac{1}{2} \right) = \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}\right)}{\Gamma\left(\frac{1}{2}a + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}b + \frac{1}{2}\right)} \frac{\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}\right)_{2m}}{\left(\frac{1}{2}a + \frac{1}{2}\right)_m \left(\frac{1}{2}b + \frac{1}{2}\right)_m}$$

which has been summed by Gauss' second theorem in (1.3). Substitution of this summation into (2.1), combined with use of the first equation in (1.4), then yields

$$F = \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}\right)}{\Gamma\left(\frac{1}{2}a + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}b + \frac{1}{2}\right)} \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}a\right)_k \left(\frac{1}{2}b\right)_k}{\left(c + \frac{1}{2}\right)_k k!}.$$

This last sum can be summed by Gauss' first theorem in (1.2) when $\text{Re}(2c - a - b) > -1$, and the desired result in (1.1) follows. This completes the proof of Watson's theorem.

References

- [1] M. Abramowitz, I. Stegun (Eds.), *Handbook of Mathematical Functions*. Dover, New York, 1965.
- [2] G. E. Andrews, R. Askey, R. Roy, *Special Functions*, Cambridge University Press, Cambridge 1999.
- [3] R. C. Bhatt, Another proof of Watson's theorem for summing ${}_3F_2(1)$, J. London Math. Soc. **40** (1965), 47–48.
- [4] T. M. MacRobert, *Functions of a complex variable*, 5th edition, Macmillan, London, 1962.
- [5] E. D. Rainville, *Special Functions*, Macmillan, New York, 1960.
- [6] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.
- [7] G. N. Watson, A note on generalized hypergeometric series, Proc. London Math. Soc. (2), **23** (1925), xiii-xv.
- [8] F. J. W. Whipple, A group of generalized hypergeometric series: relations between 120 allied series of the type $F(a, b, c; e, f)$, Proc. London Math. Soc. (2), **23** (1925), 104–114.

Received: October 3, 2008