

RVC OPEN ACCESS REPOSITORY – COPYRIGHT NOTICE

This is the peer-reviewed, manuscript version of an article published in *THERIOGENOLOGY*. The version of record is available from the journal site: <https://doi.org/10.1016/j.theriogenology.2017.10.009>.

© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

The full details of the published version of the article are as follows:

TITLE: Effect of melatonin supplementation in the long-term preservation of the sheep ovaries at different temperatures and subsequent in vitro embryo production

AUTHORS: Goodarzi, A; Zare Shahneh, A; Kohram, H; Sadeghi, M; Moazenizadeh, M H; Fouladi-Nashta, A; Dadashpour Davachi, N

JOURNAL: THERIOGENOLOGY

PUBLISHER: Elsevier


PUBLICATION DATE: January 2018

DOI: 10.1016/j.theriogenology.2017.10.009

Accepted Manuscript

Effect of melatonin supplementation in the long-term preservation of the sheep ovaries at different temperatures and subsequent in vitro embryo production

Abbas Goodarzi, Ahmad Zare Shahneh, Hamid Kohram, Mostafa Sadeghi, Mohamed Hussein Moazeni zadeh, Ali Fouladi-Nashta, Navid Dadashpour Davachi



PII: S0093-691X(17)30483-1

DOI: [10.1016/j.theriogenology.2017.10.009](https://doi.org/10.1016/j.theriogenology.2017.10.009)

Reference: THE 14292

To appear in: *Theriogenology*

Received Date: 14 April 2017

Revised Date: 1 October 2017

Accepted Date: 7 October 2017

Please cite this article as: Goodarzi A, Zare Shahneh A, Kohram H, Sadeghi M, Moazeni zadeh MH, Fouladi-Nashta A, Dadashpour Davachi N, Effect of melatonin supplementation in the long-term preservation of the sheep ovaries at different temperatures and subsequent in vitro embryo production, *Theriogenology* (2017), doi: 10.1016/j.theriogenology.2017.10.009.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Effect of melatonin supplementation in the long-term preservation of the sheep ovaries at** 2 **different temperatures and subsequent in vitro embryo production.**

3 Abbas Goodarzi^a, Ahmad Zare Shahneh^{a*}, Hamid Kohram^a, Mostafa Sadeghi^a, Mohamed Hussein Moazeni zadeh^a,
4 Ali Fouladi-Nashta^b, Navid Dadashpour Davachi^c.

5 ^a Department of Animal Science, Faculty College of Agriculture and Natural Resources, University of Tehran,
6 Karaj, Iran

7 ^b Royal Veterinary College, Reproduction Research Group, Hawkshead Campus, Hatfield, UK.

8 ^c Department of Research, Breeding and Production of Laboratory Animals, Razi Vaccine and Serum Research
9 Institute, Agricultural Research, Education and Extension Organization (AREEO), Karaj, Iran

10 * Correspondence: azareh@ut.ac.ir

11 Abstract

12 Investigations in the past decades have shown that oocytes developmental competence following in vitro
13 fertilization is greatly influenced by an interval between isolation of the ovaries immediately after
14 death/slaughter and oocytes recovery from the visible follicles. In order to determine the optimal
15 conditions for long-term preservation of ovaries, an experiment was conducted with adding different
16 doses of melatonin (0 (C), 500 (M1), 600 (M2), 700 (M3) and 800 (M4) μM) as an antioxidant to sheep
17 ovaries preservation medium (PBS) maintained at 4 and 20°C for 24 hours. The effects on in vitro
18 embryo production (IVEP) parameters including maturation, fertilization, cleavage, and blastocyst rates
19 and the total number of blastomere were evaluated after the ovaries preservation. Melatonin reduced the
20 decline in fertilization rate as an indicator of success in vitro maturation ($P \leq 0.05$). Furthermore, ovarian
21 storage time had significant negative effect ($P \leq 0.05$) on IVEP parameters. Supplementation with
22 melatonin increased the total cell number of blastocysts as an indicator of embryo quality (i.e. mean
23 blastomeric cells in 4°C groups: 86.00 ± 3.00 , 98.50 ± 3.5 , 111.5 ± 1.5 , 125.5 ± 2.00 and 126.50 ± 5.5 for C,
24 M1, M2, M3 and M4. respectively). Overall, the results showed that the use of melatonin antioxidant in
25 the ovaries storage medium had beneficial effects on sheep oocytes development and embryos quality.

26 Key words: melatonin, ovary, preservation medium, sheep

27 1. Introduction

28 Sheep is an important species in animal husbandry industry, and a critical research model to development
29 of assisted reproductive technologies (ARTs) in human and endangered species [1, 2]. Cumulus-oocyte-
30 complexes (COCs) from both small and large antral follicles can be isolated directly from ovarian tissue
31 and matured in vitro to obtain mature gametes. However, collection of oocytes for ART from live animals
32 is costly. The use of ovaries from slaughtered animals has provided an affordable source of oocytes for
33 researchers and laboratories involved in animal ART [2, 3]. However, integrating ovarian transport into in
34 vitro embryo production (IVEP) protocols has been an important challenge in large countries and/or in
35 low resource settings where the slaughterhouse is located far from the laboratory [4]. The long distance
36 transportation of ovaries to the laboratory has adverse effects on oocyte quality in terms of nuclear
37 maturation and developmental competence after the in vitro maturation (IVM) and fertilization (IVF) [5].
38 Ovaries need to be collected and returned to the laboratory instantly after the slaughter in order to make

39 effective use of the oocytes contained within [6]. The type of transportation medium [3, 6-8], storage time
40 [3-16], as well as, its temperature during transportation of the ovaries [4, 5, 10-16] are among the factors
41 affecting subsequent follicular and oocyte survival, and oocyte developmental competence [7].
42 Consequently, many studies have been conducted in order to improve oocyte preservation by modifying
43 the transportation solutions. These studies have demonstrated that the transportation of ovaries is possible
44 without a considerable harm to oocyte and follicles [4, 7, 10, 17]. However, there seems to be differences
45 between animal species in sensitivity of the oocytes to the transportation conditions. For example in
46 bovine, when the ovaries were stored for 7 hours at 4°C, 20°C, and 39°C, none of the oocytes from the
47 ovaries stored at 4°C and 39°C developed to blastocyst stage compared to the other group [13]. Whereas
48 in mice, storage of ovaries at 4°C for up to 24 hours did not affect the number of mature gametes which
49 could be collected or fertilized post-orthotropic transplantation [10].

50 Normal cellular metabolism produces reactive oxygen species (ROS) and reactive nitrogen species
51 (RNS), which regulate diverse cell functions. Reactive species, nevertheless are highly reactive with lipid,
52 protein, and nucleic acid resulting in a loss of membrane integrity, structural, or functional changes in
53 proteins, and damage in nucleic acids referred to as oxidative stress [18]. During the ovary transportation
54 to laboratory, the stoppage of blood flow reduces oxygen and energy supply, and put ovaries under
55 ischemic conditions [19]. Ischemia damages follicles viability and luteal function in ovaries; oxygen free
56 radicals in particular, are major contributors to organ damage during preservation [20]. Furthermore, the
57 antioxidant system of ovary cells is compromised during the preservation. To prevent the damage due to
58 the reactive species, the cells possess a number of antioxidant enzymes, such as superoxide dismutase
59 (SOD), glutathione peroxidase (GPX) and catalase (CAT) [7]. The balance between ROS and
60 antioxidants within the follicle seems to be critical to the function of oocyte and granulosa cells [4, 7, 21].
61 During the ovary transportation, ROS and RNS generation in the follicular microenvironment by oocyte
62 and somatic cells accompanied by reduced levels of antioxidant enzymes may cause oxidative stress-
63 mediated apoptosis in follicles [22].

64 The main goal in ovaries preservation is to maintain the function of the oocyte enclosed in follicle for in
65 vitro maturation and fertilization. Accordingly, many studies investigated the effects of different
66 antioxidants to reduce the damage caused by free oxygen radicals during ovaries preservation [6-8, 15].
67 Several studies indicated that the imbalance between ROS production and antioxidant activities could
68 cause oxidative stress [7, 23]. In cow, oocyte meiotic competence could be maintained when ovaries
69 were stored in saline supplemented with epigallocatechingallate compared to glutathione [6].
70 Experimental data from the feline model showed that oocytes from ovaries preserved in PBS
71 supplemented with SOD reduced the percentage of apoptosis in COCs compared to control group [7].

72 The protective effects of melatonin as a powerful direct scavenger of free radicals are well documented in
73 the recent years [21, 23-29]. Melatonin, because of its amphiphilic nature, diffuses broadly in diverse
74 subcellular compartments barriers. It is also a highly effective antioxidant and anti-apoptotic agent, which
75 due to its direct scavenging of toxic oxygen derivatives and its ability to reduce of ROS and RNS,
76 prevents oxidative and nitrosative damages to all macromolecules in all compartments of a
77 cell[30]. Therefore, we hypothesized that the use of this antioxidant in preservation medium may also have
78 beneficial effects by reducing oxidative stress in ovaries during long term preservation. In the present
79 study, we used the sheep as an experimental model to examine the effects of supplementing the ovaries
80 transport medium with melatonin at two different temperatures on blastocyst rate and quality after IVEP.

81 2. Materials and methods

82 Except otherwise indicated, chemicals were purchased from Sigma Chemical Co. (Saint Louis, MO,
83 USA). Adult testes and ovaries (without distinguished corpus luteum) were obtained from slaughtered
84 Lory-Bakhtiary sheep aged 2 to 3 years without apparent abnormality in reproductive tract. Due to the
85 low solubility of melatonin in water, it was dissolved in ethanol before adding to PBS.

86 2.1 Experimental design

87 There were two experimental groups in the present study design. Two different temperatures (4
88 and 20 °C) and 4 concentrations of melatonin (M1: 500, M2: 600, M3: 700 and M4: 800 µM) were
89 considered as the treatments. In experiment 1 we analyzed the effect of maintaining ovaries in PBS
90 supplemented with four concentrations of melatonin (M1- M4) at 4°C for a period of 24 h prior to oocyte
91 retrieval. As the control group(C) ovaries were maintained in PBS without melatonin supplementation for
92 a period of 24 hours before oocyte collection. Fertilization was carried out using the freshly collected
93 sperm from ram testicles of the same breed at the time of ovaries collection.

94 In the experiment 2 we assessed the effect of maintaining ovaries at 20°C for a period of 24h in PBS
95 supplemented with (M1-M4) or without (C) melatonin.

96 To evaluate the accuracy of the laboratory procedure during this study we also had considered IVF cycles
97 with standard condition (SC) concurrent with experiment 1, 2.

98 2.2 Ovaries collection

99 Ovine ovaries were collected immediately after slaughter and transferred into PBS containing 100 µg/mL
100 penicillin–streptomycin (Gibco; Grand Island, NY, USA) as a primary repository. Afterwards, the
101 collected ovaries were washed three times in sterile PBS and then randomly divided between the
102 treatment conditions in thermos flask containing PBS (C) and PBS ± melatonin (M1-M4). The insulation
103 efficiency of the thermos flask in maintaining a constant temperature was tested prior to conducting the
104 experiments. After arrival in laboratory the SC group was instantly conformed for embryo production and
105 the other groups were preserved for 24 h in the experimental conditions.

106 2.3 In vitro maturation of oocyte

107 Oocyte collection and in vitro maturation method were carried out as previously described [31], with
108 some modifications. In brief, ovarian antral follicles 2-6 mm in diameter were aspirated (45 ovaries per
109 treatment group) using an aspiration pump (MEDAP Sekretrauger P7040; Tilburg, The Netherlands)
110 fitted with a disposable vacuum line (length-35 cm, the internal diameter of 3 mm). The flow rate was set
111 at 10 mL H₂O/min using an attached disposable 20-gauge needle. Next, oocytes with more than three
112 layers of cumulus cells and uniform ooplasm were selected for IVM [31-33]. The TCM199 medium used
113 for IVM was supplemented with 10% fetal bovine serum (FBS) (Cat. Number: A6003), 0.2 mM sodium
114 pyruvate, 5 µg/mL of gentamicin, 10 µg/mL of ovine follicle-stimulating hormone (oFSH), and 1 µg/mL
115 of estradiol. COCs were cultured for 24 hours in a 50 µl droplet of maturation medium (approximately 10
116 oocytes per drop) under mineral oil at 38.5 °C in an atmosphere of 5% CO₂ with maximum humidity.
117 After 24 hours oocyte with expanded cumulus considered as mature and selected for next stage.

118 2.4 Sperm preparation, in vitro fertilization (IVF) and in vitro culture (IVC), and staining 119 methods

120 In vitro fertilization was carried out as previously described [34] with some modifications. Briefly, fresh
121 spermatozoa were obtained from the slaughtered Lory-Bakhtiary rams (2-3 years old) in fertilization day.
122 After transport of the testicles to the laboratory in a cool box (5°C), all the blood and connective tissues
123 were removed aseptically in a cold room with the temperature of 5°C. For sperm recovery, 1 mL
124 tuberculin syringe attached with 22-gauge needle was inserted into the vas deference. The content of the
125 vas deference was aspirated gently and the recovered spermatozoa was diluted 1:100 in sperm-TALP and
126 storage in 4°C less dan one heure. Samples with more than 60% progressive motility, which had normal
127 appearance, were selected for IVF. Motile spermatozoa were separated by percoll gradient (45% over
128 90%). The fertilization medium included 12 mM KCL, 25 mM NaHCO₃, 90 mM NaCl, 0.5 mM
129 NaH₂PO₄, 0.5 mM MgSO₄, 10 mM sodium lactate, 3 mg/mL BSA (fatty acid free), 50 µg/mL gentamicin.
130 At least 15 min prior to insemination the sperm suspension was transferred into a droplet of fertilization
131 medium immediately prior to co-culture adjusted to a final concentration of 1×10^6 cells/ml. Co-
132 incubation of gametes (day 0; IVF) was carried out under 5% CO₂/5% O₂ in air for 18 hours at 38.5 °C.
133 After co-incubation, cumulus cells and attached spermatozoa to zona pellucida were removed by
134 repeatedly pipetting. The presumptive fertilized eggs were cultured at 38.5°C under 5% CO₂, 5% O₂, 90%
135 N₂ for the next 7 days in synthetic oviduct fluid (SOF), used as in vitro culture (IVC) medium[31]. Two
136 days after IVC the cleavage rate of the embryo were recorded. At day 8 after insemination, the rate of
137 embryo development to the blastocyst stage was recorded.

138 Examination of the IVF results carried out according to the previously described procedures [31]. Briefly,
139 24 hours after insemination one fourth of zygote were removed randomly from the culture medium and
140 washed twice in PBS-PVP (Polyvinylpyrrolidone, 1mg/mL). Then the zygote were fixed on a 100 µL
141 drop of paraformaldehyde solution [4% (w/v) in PBS, pH 7.4], at room temperature. In the final step, the
142 zygote were removed from the fixation drop and washed three times in PBS-PVP. Afterwards, zygotes
143 were transferred to 100µl drop of PBS-PVP containing 1mg/mL Hoechst 33342 for assessment of
144 fertilization rate. The slides were examined under a fluorescent microscope with a UV filter showing the
145 sperm head and pronuclei having a blue appearance. The same method was used for counting the total
146 cells in blastocysts day 8 after insemination.

147 2.5 Statistical analysis

148 The GLM procedure of the SAS (SAS, Inc., Cary, NC, USA) was used for the analysis of
149 variance (ANOVA). The Duncan test was taken into account for comparisons of mean values with a
150 significant main effect. P values <0.05 were considered statistically significant. Data are presented as
151 mean ± SEM, total number of blastomeric are presented as number.

152 3. Results

153 As expected, the standard condition group which was used as a control to assure the accuracy of
154 IVEP procedure, resulted in higher maturation, fertilization, blastocyst rates and mean
155 blastomeric cells (93.12±0.47, 84.62±0.75, 28.27±1.73 and 171.5±4.50, respectively). The

156 oocytes in this group were retrieved from ovaries which were transported to the laboratory in less
157 than three hours in PBS at 37.5°C,

158 3.1 Effects of storage condition at 4°C

159 The effect of melatonin supplementation in ovaries preservation for 24 h at 4°C is presented in Table 1.
160 The results show that oocyte maturation rate was affected by treatment ($P < 0.05$). Only M3 (700) and M4
161 (800) increased maturation rates compared to the control, and the rates were similar to M1 and M2
162 groups. Higher concentration of melatonin (M3 and M4) improved fertilization and cleavage rates
163 compared to M1 and control group ($p < 0.05$). Moreover, the highest melatonin supplementation (M4:
164 800) increased morula rates compared to all groups. Interestingly, some of the cleaved embryos in control
165 group developed to blastocyst stage. The percentages of zygotes undergoing blastocyst in oocytes
166 obtained from ovaries of the M3 and M4 groups were similar (16.95 ± 0.59 vs. 18.19 ± 0.33 , respectively)
167 but in all melatonin-treated groups were significantly higher ($P < 0.05$) than C group. The total cell number
168 was significantly greater for M3-M4 compared with M1 and C blastocysts (Figure 1. a-f).

169 3.2 Effects of storage condition at 20°C

170 As shown in Table 2, supplementation of different concentration of melatonin to ovaries preservation
171 medium led to significant increase in oocyte maturation rate evaluated based on cumulus expansion ($p <$
172 0.05). There were no significant differences between 500, 700, and 800 μM treatment groups. No
173 differences were observed in the maturation rate of oocytes from PBS alone (control) with 600 μM
174 melatonin (M2) group. Also, fertilization rate were not significant between M3 and M4 groups ($p < 0.05$).
175 Higher concentration of melatonin (M4; 800 μM) resulted in the best fertilization rate. Oocytes from
176 control group and low melatonin groups had inferior fertilization rate ($p < 0.05$). Similarly, the percent of
177 cleaved zygote in melatonin-treated groups was higher than the control group and the highest
178 concentration of melatonin resulted in higher cleavage rate ($p < 0.05$), except compared to M3 group.
179 Moreover, the highest melatonin concentration showed the highest morula rates compared to all groups
180 ($P < 0.05$). No blastocyst was formed from the control group. Melatonin supplementation resulted in
181 significant increase in blastocyst formation with the higher doses (700 and 800 μM) resulting in higher
182 blastocyst rate than the lower doses (500 and 600 μM melatonin). Similar results were obtained for
183 blastocyst quality as compared by the mean number of blastomeric cells being significantly higher in M3
184 and M4 groups than the low melatonin groups ($p < 0.05$) (Figure 1. B-F).

185 4. Discussion

186 Preservation of ovarian tissue is challenging because there are several cell types within the tissue that
187 need to be maintained in a normal state. During the ovaries transportation to the laboratory, the stoppage
188 of blood flow reduces energy supply, and put ovaries under ischemic conditions. As well as, storage of
189 ovary at atmospheric level of oxygen (21%) could generate higher amounts of free radicals in follicle
190 microenvironment, which is toxic to oocyte and other cells surrounding it. Most likely due to the
191 formation of superoxide anion ($\text{O}_2^{\cdot-}$), this leads to the ulterior formation of HO^{\cdot} , H_2O_2 , NO^{\cdot} , and ONOO^- ,
192 which results in oxidative stress [24, 35, 36]. The oocyte oxidative stress caused by ROS must be limited
193 in order for a good embryo to be produced [36]. Free radicals spreads within follicle microenvironment,
194 and react with proteins, lipids and DNA of oocyte and cumulus cells leading to cell membrane-lipid
195 peroxidation, DNA damage and apoptosis [27, 37]. The results reported here show that the

196 supplementation of ovaries transport medium (PBS) with melatonin significantly maintained the
197 developmental potential of the oocytes than PBS for 24 hours at 4 and 20°C. In particular, the oocyte
198 competence for development the blastocyst stage was significantly improved as compared to control.
199 Furthermore, high-dose melatonin (700 and 800 µM) during 24 h storage in both 4, 20°C resulted in
200 significant improvement in blastocyst quality as evidenced by greater numbers of blastomeric cells in
201 these groups (Tables 1 and 2).

202 The melatonin directly suppresses free radicals derived from superoxide anion. In addition, metabolites
203 that are formed during interaction with free radicals (i.e., cyclic 3-hydroxymelatonin, N1-acetyl-N2-
204 formyl-5-methoxykynuramine, and N1-acetyl-5-methoxykynuramine) are high potent scavengers of toxic
205 reactants [24]. Furthermore, it is well known that melatonin has an important indirect function in up
206 regulation of antioxidant and down regulation of prooxidant enzymes [30]. Brzezinski et al [38] reported
207 that melatonin concentration in human follicular fluid is significantly higher than serum samples.
208 Melatonin in follicular fluid besides their physiological role in steroidogenic mechanism, acts as free
209 radical scavenger [21]. The antioxidant activities of melatonin in oxidative stress has been reported in
210 different species. In the mouse oocyte, melatonin promote the development rate of mouse two-cell
211 embryos to blastocysts stage [26]. Moreover, melatonin treatment increases intrafollicular concentration
212 of melatonin and improved fertilization rate embryo transfer [39]. A recent study presented that
213 supplementation of IVM medium with melatonin-loaded lipid-core Nano capsules (Mel-LNC) during
214 IVEP improved cleavage and blastocyst rates of bovine embryo. Also, significantly decreased ROS
215 levels, and down-regulated the genes involved apoptosis caspase 3 (CASPA3) B and BAX [40].
216 Rodriguez-Osorio et al. [37] reported that, in IVC medium melatonin supplementation improved cleavage
217 rates in stressed conditions (H₂O₂ or 40°C) and increased blastocyst cell numbers in usual terms. Our
218 results showed a significant effect of melatonin on developmental potential of oocytes derived from
219 ovaries preserved in supplemented medium. In addition, our analysis demonstrated that the resultant
220 blastocysts in the M3 (700µM) and M4 (800µM) groups were better considering the healthy blastomeric
221 cells (Figure 1. E, e, F and f). These findings demonstrated that melatonin significantly reduced the toxic
222 effect of ROSs that generated in ovaries during storage period.

223 Although, improvement was noted when ovaries stored in melatonin supplemented medium, this positive
224 effects in low temperature (4°C) were more evident than that of high temperature (20°C). The oocytes
225 from the ovaries stored at 20°C without melatonin (control) lost their competence to develop to blastocyst
226 stages. Whereas, when ovaries stored at 4°C, obtained COCs in untreated group developed to blastocyst
227 stage, although, had significantly lower rate than in melatonin treated groups (Tables 1, 2). It is well
228 known that preservation of organs in hypothermia suppresses metabolic requirements. Consequently, it
229 results in reduced tissue damage caused by loss of ATP, increase in pH, lipid peroxidation and proteolysis
230 associated with ischemic condition [12, 20, 40]. Several studies investigated effects of temperature on
231 pool of follicles in farm animal ovaries. In the Iberian red deer, preservation of ovaries in hypothermia
232 condition significantly elevated cleavage rate [4]. Nakao et al. [13] reported that oocyte obtained from
233 ovaries stored at 20°C had higher competence than that of stored at 39°C or 4°C. Results showed that
234 temperature of preservation media affected the quality of oocytes, in which cytoplasmic membrane,
235 microtubule, cytoskeleton and zona pellucida might be sensitive to low temperatures [5, 12]. Canine
236 ovaries were transported for up to 4 hours at 4 and 35°C, obtained oocyte from the ovaries stored at 4°C
237 had higher MII maturation rate compared with the other group [16]. In the present study, the storage of

238 sheep ovaries in low temperature media exhibited better results in terms of the developmental competence
239 of oocytes and the cleavage and production of the morula and blastocysts in contrast to high temperature.

240 In conclusion, the present study showed that (i) supplementation of phosphate buffer saline as ovine
241 ovaries storage media with melatonin (500, 600, 700 and 800 μM), reduced the oocytes competence
242 decline to develop to blastocyst stage. (ii) for long preservation of ovaries (24 hours), low storage
243 temperature (4°C) can maintain oocyte competence but higher temperature (20°C) has a detrimental effect
244 on oocyte viability, maturation, fertilization and subsequent development after IVF; and (iii) storage of
245 ovaries in high concentration (700, 800 μM) for 24 hours significantly maintain the oocyte quality and
246 obtained blastocysts were better than other group, but this protective effect of melatonin were improved in
247 cool condition. Beneficial effect of melatonin in ovine ovaries storage medium may be employed to
248 reduce injuries mediated by ROS during ovaries preservation.

249 Acknowledgments

250 The authors would like to thank Helia Shamsi for her assistance during the laboratory procedure. We
251 would also like to express our appreciation to the University of Tehran for financial support of for this
252 project under grant number: 7108011/6/44. Similarly, the authors would like to acknowledge the financial
253 support of Iran National Science Foundation under grant number: 94810027.

254 Table 1: Effects of ovaries preservation for 24 h at 4°C in melatonin supplemented PBS on oocyte and embryo
255 development.

antioxidant dosage (μM)	Cumulus cells expansion% (n)	Fertilization% (n*)	Cleaved zygote on day 2 % (n)	Morula rate% (n)	Blast rate% (n)	Mean blastomere (n)
0	84.19 \pm 1.52 ^c (122/145)	72.12 \pm 2.88 ^c (18/25)	61.98 \pm 3.98 ^c (60/97)	34.09 \pm 2.09 ^c (33/97)	7.19 \pm 0.81 ^c (7/97)	86.00 \pm 3.00 ^c
500	86.83 \pm 2.21 ^{bc} (120/138)	71.67 \pm 1.66 ^c (18/25)	68.34 \pm 1.66 ^{dc} (65/95)	44.11 \pm 1.89 ^b (42/95)	10.56 \pm 0.55 ^b (10/95)	98.50 \pm 3.5 ^{bc}
600	90.18 \pm 2.68 ^{abc} (128/142)	76.79 \pm 1.78 ^{bc} (20/26)	72.28 \pm 0.27 ^{cd} (73/101)	44.57 \pm 1.43 ^b (45/101)	10.90 \pm 1.10 ^b (11/101)	111.5 \pm 1.5 ^b
700	91.20 \pm 1.28 ^{ab} (142/156)	80.00 \pm 0.75 ^{ab} (24/30)	75.9 \pm 0.46 ^{bc} (85/112)	49.99 \pm 0.89 ^b (56/112)	16.95 \pm 0.59 ^a (19/112)	125.00 \pm 2.00 ^a
800	91.22 \pm 0.50 ^{ab} (135/148)	80.13 \pm 3.20 ^{ab} (20/25)	80.89 \pm 1.26 ^{ab} (89/110)	56.35 \pm 0.79 ^a (62/110)	18.19 \pm 0.33 ^a (20/110)	126.50 \pm 5.5 ^a

256 Each value represents the means \pm SEM.

257 * randomly selected from zygotes for assessment IVF parameters

258 Table 2: . Effects of ovaries preservation for 24 h at 20°C in melatonin supplemented PBS on oocyte and embryo
259 development. No melatonin was added to the control group.

Antioxidant dosage (μM)	Cumulus cells expansion % (n)	Fertilization% (n)	Cleaved zygote on day 2 % (n)	Morula rate% (n)	Blast rate% (n)	Mean blastomere (n)
0	75.50 \pm 1.64 ^b (119/158)	53.57 \pm 3.57 ^c (14/26)	32.42 \pm 5.59 ^c (30/93)	21.53 \pm 5.74 ^c (20/93)	0.00 ^c (0/93)	-

500	83.13±1.88 ^a (133/160)	62.08±1.19 ^b (18/29)	52.53±3.47 ^b (54/104)	33.38±1.24 ^b (35/104)	5.83±0.17 ^b (6/104)	83.50±4.50 ^c
600	79.75±0.93 ^b (138/173)	65.48±109 ^b (19/29)	60.67±2.98 ^b (65/109)	33.81±0.48 ^b (37/109)	7.48±0.21 ^b (8/109)	89.00±4.00 ^{bc}
700	80.37±0.88 ^a (127/158)	75.13±1.80 ^{ab} (21/28)	70.82±1.18 ^a (75/99)	41.24±2.35 ^b (40/99)	10.36±0.36 ^a (11/99)	105.50±5.50 ^{ab}
800	80.29±1.04 ^a (130/158)	77.38±5.95 ^a (20/26)	68.66±1.34 ^a (70/104)	45.72±2.86 ^a (47/104)	12.72±0.73 ^a (13/104)	119.00±6.00 ^a


260 Each value represents the means ± SEM.

261 * Randomly selected from zygotes for assessment IVF parameters



262


263

SC



C





264 Figure 1. Example of blastocysts 7 days. Stained with Hoechst fluorescent dye. SC: Blastocyst obtained in
265 conventional method (up to 3h storage in 37.5°C). C: from control group (ovaries stored without melatonin

266 supplementation) at 4°C [B] or 20°C [b]. M1-M4: Blastocysts from ovaries stored with 500, 600, 700 or 800 µM
 267 melatonin at 4 [C, D, E and F] or 20°C [c, d, e and f]. A high-resolution version of panels A, B, b, C, c, D, d, E, e, F
 268 and f of this image are available as Virtual Microscope eSlides: VM04430, VM04431, VM04432, VM04433,
 269 VM04434, VM04435, VM04436, VM04437, VM04438, VM04439 and VM04440, respectively.

270

271 References

272 [1] Menchaca A, Anegón I, Whitelaw CBA, Baldassarre H, Crispo M. New insights and current tools for genetically
 273 engineered (GE) sheep and goats. *Theriogenology*. 2016;86:160-9.

274 [2] Wani NA. In vitro maturation and in vitro fertilization of sheep oocytes. *Small Ruminant Research*. 2002;44:89-
 275 95.

276 [3] Gonçalves R, Cavalcante A, Gouveia B, Lins T, Barberino R, Menezes V, et al. Lower apoptosis rate in ovine
 277 preantral follicles from ovaries stored in supplemented preservation media. *Zygote*. 2015;23:943-50.

278 [4] García-Álvarez O, Maroto-Morales A, Berlinguer F, Fernández-Santos M, Esteso M, Mermillod P, et al. Effect
 279 of storage temperature during transport of ovaries on in vitro embryo production in Iberian red deer (*Cervus elaphus*
 280 *hispanicus*). *Theriogenology*. 2011;75:65-72.

281 [5] Wongsrikeao P, OTOI T, KARJA NWK, AGUNG B, NII M, NAGAI T. Effects of ovary storage time and
 282 temperature on DNA fragmentation and development of porcine oocytes. *Journal of Reproduction and*
 283 *Development*. 2005;51:87-97.

284 [6] Nagao Y, Harada Y, Yamaguchi M, Igarashi A, Ooshima Y, Kato Y. Antioxidant treatment during preservation
 285 of bovine ovaries increased the development potential of embryos. *Zygote*. 2010;18:315-21.

286 [7] Cocchia N, Corteggio A, Altamura G, Tafuri S, Rea S, Rosapane I, et al. The effects of superoxide dismutase
 287 addition to the transport medium on cumulus-oocyte complex apoptosis and IVF outcome in cats (*Felis catus*).
 288 *Reproductive biology*. 2015;15:56-64.

289 [8] Gomes R, Andrade E, Lisboa L, Ciquini A, Barreiros T, Fonseca N, et al. Effect of holding medium, temperature
 290 and time on structural integrity of equine ovarian follicles during the non-breeding season. *Theriogenology*.
 291 2012;78:731-6.

292 [9] Hornick J, Duncan F, Shea L, Woodruff T. Isolated primate primordial follicles require a rigid physical
 293 environment to survive and grow in vitro. *Human reproduction*. 2012;27:1801-10.

294 [10] Kamoshita K, Okamoto N, Nakajima M, Haino T, Sugimoto K, Okamoto A, et al. Investigation of in vitro
 295 parameters and fertility of mouse ovary after storage at an optimal temperature and duration for transportation.
 296 *Human Reproduction*. 2016:dew023.

297 [11] Lucci CM, Kacinskis MA, Rumpf R, Bão SN. Effects of lowered temperatures and media on short-term
 298 preservation of zebu (*Bos indicus*) preantral ovarian follicles. *Theriogenology*. 2004;61:461-72.

299 [12] Matsushita S, Tani T, Kato Y, Tsunoda Y. Effect of low-temperature bovine ovary storage on the maturation
 300 rate and developmental potential of follicular oocytes after in vitro fertilization, parthenogenetic activation, or
 301 somatic cell nucleus transfer. *Animal reproduction science*. 2004;84:293-301.

302 [13] Nakao H, Nakatsuji N. Effects of storage conditions of bovine ovaries and oocytes on the success rate of in
 303 vitro fertilization and culture. *Journal of Reproduction and Development*. 1992;38:11-3.

304 [14] Naoi H, Otoi T, Shimamura T, Karja NWK, Agung B, Shimizu R, et al. Developmental competence of cat
 305 oocytes from ovaries stored at various temperature for 24 h. *Journal of Reproduction and Development*.
 306 2007;53:271-7.

- 307 [15] Silva J, Lucci C, Carvalho F, Bao S, Costa S, Santos R, et al. Effect of coconut water and Braun-Collins
308 solutions at different temperatures and incubation times on the morphology of goat preantral follicles preserved in
309 vitro. *Theriogenology*. 2000;54:809-22.
- 310 [16] Taş M, Evecen M, ozdaş OB, Cirit U, Demir K, Birler S, et al. Effect of transport and storage temperature of
311 ovaries on in vitro maturation of bitch oocytes. *Animal reproduction science*. 2006;96:30-4.
- 312 [17] Onions V, Mitchell M, Campbell B, Webb R. Ovarian tissue viability following whole ovine ovary
313 cryopreservation: assessing the effects of sphingosine-1-phosphate inclusion. *Human Reproduction*. 2008;23:606-
314 18.
- 315 [18] Devine PJ, Perreault SD, Luderer U. Roles of Reactive Oxygen Species and Antioxidants in Ovarian Toxicity
316 1. *Biology of reproduction*. 2011;86:Article 27, 1-10.
- 317 [19] Iwata H, Hayashi T, Sato H, Kimura K, Kuwayama T, Monji Y. Modification of ovary stock solution with
318 magnesium and raffinose improves the developmental competence of oocytes after long preservation. *Zygote*.
319 2005;13:303-8.
- 320 [20] Guibert EE, Petrenko AY, Balaban CL, Somov AY, Rodriguez JV, Fuller BJ. Organ preservation: current
321 concepts and new strategies for the next decade. *Transfusion Medicine and Hemotherapy*. 2011;38:125-42.
- 322 [21] Adriaens I, Jacquet P, Cortvrindt R, Janssen K, Smits J. Melatonin has dose-dependent effects on
323 folliculogenesis, oocyte maturation capacity and steroidogenesis. *Toxicology*. 2006;228:333-43.
- 324 [22] Cruz M, Leal C, Cruz J, Tan D, Reiter R. Essential actions of melatonin in protecting the ovary from oxidative
325 damage. *Theriogenology*. 2014;82:925-32.
- 326 [23] Jiang J, Cheung C, Wang Y, Tsang B. Regulation of cell death and cell survival gene expression during ovarian
327 follicular development and atresia. *Frontiers in bioscience: a journal and virtual library*. 2003;8:d222-37.
- 328 [24] Bonnefont-Rousselot D, Collin F, Jore D, Gardès-Albert M. Reaction mechanism of melatonin oxidation by
329 reactive oxygen species in vitro. *Journal of pineal research*. 2011;50:328-35.
- 330 [25] Galano A, Tan DX, Reiter RJ. On the free radical scavenging activities of melatonin's metabolites, AFMK and
331 AMK. *Journal of pineal research*. 2013;54:245-57.
- 332 [26] Ishizuka B, Kuribayashi Y, Murai K, Amemiya A, Itoh MT. The effect of melatonin on in vitro fertilization and
333 embryo development in mice. *Journal of pineal research*. 2000;28:48-51.
- 334 [27] Izadpanah G, Zare-Shahneh A, Zhandi M, Yousefian I, Emamverdi M. Melatonin has a beneficial effect on
335 stallion sperm quality in cool condition. *Journal of Equine Veterinary Science*. 2015;35:555-9.
- 336 [28] Mondal P, Hasan KN, Pal PK, Maitra SK. Influences of exogenous melatonin on the oocyte growth and
337 oxidative status of ovary during different reproductive phases of an annual cycle in carp *Catla catla*.
338 *Theriogenology*. 87:349-59.
- 339 [29] Tamura H, Nakamura Y, Korkmaz A, Manchester LC, Tan D-X, Sugino N, et al. Melatonin and the ovary:
340 physiological and pathophysiological implications. *Fertility and sterility*. 2009;92:328-43.
- 341 [30] R Ramis M, Esteban S, Miralles A, Tan D-X, J Reiter R. Protective effects of melatonin and mitochondria-
342 targeted antioxidants against oxidative stress: a review. *Current medicinal chemistry*. 2015;22:2690-711.
- 343 [31] Dadashpour Davachi N, Zare Shahneh A, Kohram H, Zhandi M, Shamsi H, Hajiyavand AM, et al. Differential
344 influence of ampullary and isthmic derived epithelial cells on zona pellucida hardening and in vitro fertilization in
345 ovine. *Reproductive Biology*. 2016;16:61-9.
- 346 [32] Dadashpour Davachi N, Kohram H, Zainoaldini S. Cumulus cell layers as a critical factor in meiotic
347 competence and cumulus expansion of ovine oocytes. *Small Ruminant Research*. 2012;102:37-42.

- 348 [33] Dadashpour Davachi N, Kohram H, Zare Shahneh A, Zhandi M, Goudarzi A, Fallahi R, et al. The effect of
349 conspecific ampulla oviductal epithelial cells during in vitro maturation on oocyte developmental competence and
350 maturation-promoting factor (MPF) activity in sheep. *Theriogenology*. 2017;88:207-14.
- 351 [34] Dashti S, Zare Shahneh A, Kohram H, Zhandi M, Dadashpour Davachi N. Differential influence of ovine
352 oviduct ampullary and isthmic derived epithelial cells on in vitro early embryo development and kinetic. *Small*
353 *Ruminant Research*. 2016;136:197-201.
- 354 [35] Fischer B, Bavister B. Oxygen tension in the oviduct and uterus of rhesus monkeys, hamsters and rabbits.
355 *Journal of reproduction and fertility*. 1993;99:673-9.
- 356 [36] Thompson J, Simpson A, Pugh P, Donnelly P, Tervit H. Effect of oxygen concentration on in-vitro
357 development of preimplantation sheep and cattle embryos. *Journal of reproduction and fertility*. 1990;89:573-8.
- 358 [37] Rodriguez Osorio N, Kim I, Wang H, Kaya A, Memili E. Melatonin increases cleavage rate of porcine
359 preimplantation embryos in vitro. *Journal of pineal research*. 2007;43:283-8.
- 360 [38] Brzezinski A, Seibel MM, Lynch HJ, Deng M-H, Wurtman RJ. Melatonin in human preovulatory follicular
361 fluid. *The Journal of Clinical Endocrinology & Metabolism*. 1987;64:865-7.
- 362 [39] Tamura H, Takasaki A, Miwa I, Taniguchi K, Maekawa R, Asada H, et al. Oxidative stress impairs oocyte
363 quality and melatonin protects oocytes from free radical damage and improves fertilization rate. *Journal of pineal*
364 *research*. 2008;44:280-7.
- 365 [40] Remião MH, Lucas CG, Domingues WB, Silveira T, Barther NN, Komninou ER, et al. Melatonin delivery by
366 nanocapsules during in vitro bovine oocyte maturation decreased the reactive oxygen species of oocytes and
367 embryos. *Reproductive Toxicology*. 2016;63:70-81.

368

369

1 **Highlight**

2

3 Effect of melatonin supplementation in ovaries storage medium for up 24 hours in 4 and 20°C
4 was studied on oocyte quality and embryo production in sheep.

5 Oocyte competence, fertilization rate, morula and blastocyst improved by melatonin (500, 600,
6 700 and 800 μM) supplementation in ovaries storage medium.

7 Storage of ovaries in high concentration (700, 800 μM) for 24 hours significantly maintain the
8 oocyte quality and obtained blastocysts were better than other groups.