



Review

Sagittal joint spaces of the temporomandibular joint: Systematic review and meta-analysis



Eugénio Martins^a, Joana C. Silva^{a,*}, Carlos A. Pires^b, Maria J. Ponces^a, Jorge D. Lopes^a

^a Faculdade de Medicina da Dentária da Universidade do Porto, Portugal

^b Departamento de matemática, Universidade de Trás-os-montes e Alto Douro, Portugal

ARTICLE INFO

Article history:

Received 9 February 2015

Accepted 12 April 2015

Available online 26 May 2015

Keywords:

Temporomandibular joint

Review

Meta-analysis

ABSTRACT

The aim of this study was to perform a systematic review and meta-analysis on the sagittal joint spaces measurements of the temporomandibular joint. An electronic database search was performed with the terms “condylar position”; “joint space” AND “TMJ”. The risk of bias of each study was assessed with “Cochrane risk of bias tool”. The values used in the meta-analysis were the joint space measurements and their differences between the right and left joint.

From the initial search 2706 articles were retrieved. Eighteen articles classified for final review. Only one study was classified as having high level of evidence. Seventeen of the reviewed studies were included in the meta-analysis concluding that the mean sagittal joint space values were: anterior joint space 1.86 mm, superior 2.36 mm and posterior 2.22 mm. However, the analysis also showed high levels of heterogeneity. Right and left comparison has shown statistically significant differences.

© 2015 Sociedade Portuguesa de Estomatologia e Medicina Dentária. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Espaços articulares sagitais da articulação temporomandibular: revisão sistemática e meta-análise

RESUMO

O objetivo deste estudo foi realizar uma revisão sistemática e meta-análise sobre os espaços articulares sagitais da articulação temporomandibular. Foi realizada uma pesquisa eletrónica com os termos “condylar position”, “joint space” AND “TMJ”. O nível de evidência de cada estudo foi avaliado com “Cochrane risk of bias tool”. Os valores sumariados na meta-análise foram os espaços articulares e a diferença entre a articulação direita e esquerda.

Palavras chave:

Articulação temporomandibular

Revisão

Meta-análise

* Corresponding author.

E-mail address: joanacristinapsilva@gmail.com (J.C. Silva).

<http://dx.doi.org/10.1016/j.rpemd.2015.04.002>

1646-2890/© 2015 Sociedade Portuguesa de Estomatologia e Medicina Dentária. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Da pesquisa inicial resultaram 2076 artigos dos quais 18 foram selecionados para a revisão. Apenas um estudo foi considerado de elevado nível de evidência. Foram incluídos na meta-análise 17 dos artigos da revisão concluindo-se que, os valores médios para os espaços articulares sagitais foram: 1.86 mm para o anterior, 2.36 mm para o superior e 2.22 mm para o posterior. No entanto, a análise revelou ainda grande heterogeneidade nos resultados dos estudos avaliados. Verificaram-se diferenças estatisticamente significativas entre as articulações esquerda e direita.

© 2015 Sociedade Portuguesa de Estomatologia e Medicina Dentária. Publicado por Elsevier España, S.L.U. Este é um artigo Open Access sob a licença de CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The mandibular condyle position has been at the centre of a long lasting controversy among gnathologists and orthodontists. The ideal concept of the mandibular condyle position has changed from the most retruded position of the condyle in the glenoid fossa to the most superior position of the condyle. Nowadays it is accepted as the most anterosuperior position of the mandibular condyle in the glenoid fossa with the articular disk placed in between.¹⁻⁴ The literature also shows a great confusion concerning the relationship between dental occlusion and the temporomandibular joints. It is possible to find articles proving the relationship between these two variables, while others achieved contrary results with no relationship being suggested.⁵⁻⁸ The major focus of the discussion usually is the ideal mandibular condyle position and the effects of its variation.^{4,9,10} With the evolution of radiographic exams like computerized tomographies (CT), including the new 3D cone-beam computed tomography (CBCT) and magnetic resonance imaging (MRI) it is now possible to radiographically examine the position of the condyle.¹¹⁻¹³ The most common method found in the literature to determine this position is the assessment of the joint space measurements, which are the radiographic space found between the condyle and the glenoid fossa where the articular disk is placed.¹⁴ A variation on the values of these measurements suggest a displacement of the condyle and so, the determination of the "gold standard" for these values would be a very important tool to determine any variation to the condyle position. The aim of this study is to perform a systematic review and meta-analysis on the sagittal joint space measurements of the temporomandibular joint to assess the mean values for these measurements.

Methods

Search strategy

A comprehensive electronic database search to identify relevant publications was conducted, and the reference lists in relevant articles were searched manually for additional literature. No language restrictions were set, although no attempt was made to explore the informally published literature, like conference proceedings and abstracts of researches presented at conferences and dissertations. The research extended to the following databases: Medline (Pubmed), Lilacs, Scopus, Ebsco

(Host by University of Porto) and Cochrane Central Register of Controlled Clinical Trials.

The search terms were "condylar position" and "joint space" AND "TMJ" with no year of publication restriction in order to include the highest number of articles (to 22 April 2014). No restriction to study design was applied.

Faculty of Dental Medicine of University of Porto and Portuguese Society of Dentofacial Orthopedics' libraries were also consulted for printed articles not available online.

Critical evaluation

At the first stage, two reviewers screened independently the titles of the retrieved records, and only the titles related to TMJ joint space assessment were included. Joint space was defined as the radiographic image between the mandibular condyle and the glenoid fossa where the disk is interposed. Next, the abstracts of the retrieved publications were read by the two reviewers and categorized according to the radiographic procedure used to assess the joint space. An article had only to be justified by one reviewer to be included in the second selection phase. Eligibility of the retrieved articles was determined by applying the following inclusion criteria: (1) tomographic examination of the TMJ (2) determination of sagittal joint space measurements at least on two different points (anterior and posterior). The main reasons for exclusion were: mandible fractures, studies not performed in living humans, surgical interventions, studies with patients with syndromes or chronic diseases (including degenerative pathology of the TMJ), samples containing patients only in the primary or mixed/early permanent dentition, clinical only evaluation of the mandibular condyle position, 2D radiograph or magnetic resonance imaging, previous orthodontic or splint therapy, case reports, discussion or debate articles. All not published studies were also excluded.

The analysis was based on primary materials. When an abstract was considered by at least one author to be relevant, it was read in full text. At the second stage, the full texts were retrieved and critically examined. Reference lists from the articles selected on the second stage were screened and articles related to joint space measurements were hand-searched.

Data gathering

The following data were extracted from the selected articles: year of publication, study type, study method, sample description, joint space measurements on the sagittal plane,

error analysis method, statistical analysis and author's conclusion. This method was pilot-tested on ten randomly selected included articles and then refined. One reviewer author then extracted the mentioned data from the included articles and the second author checked. Any disagreement was resolved with discussion between the two authors until a consensus was reached. The risk of bias was assessed according to the "Cochrane risk of bias tool"¹⁵ as suggested by the "PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration".¹⁶

Meta-analysis

The values studied in this meta-analysis were the sagittal joint space measurements (anterior, superior and posterior joint space) and the differences between the right and left joints. As not all the included articles presented the values for all the spaces from the right and left joints, the analysis were performed including all the data presented in each selected study. For the comparative analysis between the right and

left joints, mean and standard deviation values from the samples of each article were used. For global joint space assessment, mean and standard deviation of the total sample (including both the values from the right and left joints) were used.

The restricted maximum-likelihood (REML) method was used to estimate de variability between the studies. Inverse variance method was used to assess the weight of each study.

Heterogeneity was determined using the Q Cochran test and the I^2 statistics by Higgins and Thompson.¹⁷

Statistical analysis was performed using "R", version 2.15.2 from "The R Project for Statistical Computing", available from <http://www.r-project.org>.

Results

Search results

The initial search strategy allowed retrieving 916 articles from Medline (Pubmed), 1114 from Scopus, 158 from EBSCOhost,




Fig. 1 – Flow diagram illustrating the search strategy results.

Source: Adapted from: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(6): e1000097. doi:10.1371/journal.pmed1000097.

19 from Lilacs and none from the Cochrane Central Register of Controlled Clinical Trials. The number of articles reviewed in each phase of this systematic review is presented in the PRISMA flow diagram (Fig. 1). After excluding 978 duplicates, 1230 articles remained for review. In the first phase selection, the observers screened the articles by reading titles and abstracts. Articles that were not eligible because of irrelevant aims and were not directly related to this systematic review were excluded, thus 61 articles remained for further reading. 28 articles were assessed for eligibility. After screening all the articles full text according to the inclusion/exclusion criteria, 18 articles classified for final review.

Type of study and method used for joint space assessment

A Randomized Clinical Trial (RCT)¹⁸ was found. Additionally, six prospective¹⁹⁻²⁴ and eleven retrospective²⁵⁻³⁵ studies were found that fulfilled the eligibility criteria defined for this review. Six of the selected articles^{22,25,27,32-34} used CBCT to assess the TMJ joint space while nine^{18-21,23,26,28,30,35} used conventional CT and three^{24,29,31} used linear tomography.

Twelve of the selected studies^{19-21,23,24,26,28-30,32,33,35} assessed the joint space measurements by determining the closest distance between the mandibular condyle and the glenoid fossa surface.

Table 1 – Summary of the quality analysis of the 18 included studies.

	Estimation of sample size	Sample description	Error analysis	Normality tests	Adequate statistics provided	Randomization	Statistical analysis	Level of evidence
1 ²⁵	No/Not known	Yes	No	Yes	Yes	No	Adequate	Low
2 ³²	No/Not known	Yes	No	No information N > 30	Yes	No	Adequate (correlation)	Low
3 ²⁶	No/Not known	Yes	Yes	No information	Yes	No	Adequate	Low
4 ¹⁹	No/Not known	Incomplete	No	No information N > 30	Yes	No	Adequate (correlation)	Low
5 ²⁷	No/Not known	Yes	Yes	No information	Yes	No	Adequate	Low
6 ¹⁸	No/Not known	Yes	Yes	Yes	Yes	Yes	Adequate	Moderate
7 ²⁸	No/Not known	Incomplete	Yes	No information	Yes	No	Adequate	Low
8 ²⁹	No/Not known	Yes	No	No information N > 30	Incomplete	No	Adequate (correlation)	Low
9 ³³	Yes	Yes	Yes	No information Non-parametric tests	Yes	No	Adequate	Low
10 ²⁰	No/Not known	Incomplete	Yes	No information N > 30	Yes	No	Adequate	Low
11 ²¹	No/Not known	Incomplete	Yes	No information	Yes	No	Adequate (correlation)	Low
12 ²²	No/Not known	No	Yes	No information	Yes	No	Adequate	Low
13 ³¹	No/Not known	Yes	No	No information N > 30	Yes	No	Adequate	Low
14 ²³	No/Not known	Incomplete	No	No information	Yes	No	Adequate	Low
15 ³⁴	No/Not known	Yes	No	No information	Incomplete	No	Adequate	Low
16 ²⁴	No/Not known	Yes	No	No information	Incomplete	No	Adequate (correlation)	Low
17 ³⁰	No/Not known	Yes	No	No information	Incomplete	No	Adequate	Low
18 ³⁵	No/Not known	Incomplete	No	No information N = 30	Yes	No	Adequate (correlation)	Low




Fig. 2 – Mean anterior joint space in each study and globally.

On the other hand, four articles^{25,27,31,34} used a geometrical construction to assess the joint space measurements. Different authors used the Frankfurt horizontal line or the true horizontal line as a reference plane to determine the most superior point of the glenoid fossa. Following, the distance between this point and the highest point of the condyle (determined by the same method) was measured, resulting the value of the sagittal joint space. After this, starting from the most superior point of the glenoid fossa, two tangent lines were traced to the most anterior and posterior point of the condyle respectively. The distance between each of these points and the point where a perpendicular line to the tangents crosses the surface of the glenoid fossa was the anterior and posterior joint space respectively.

Two studies^{18,22} used a similar method to the one described above, but used the centre of the mandibular condyle as the reference point.

Quality assessment

The summary of the quality analysis of the selected articles is presented in Table 1.

In general, the statistical analysis performed was adequate to the objectives defined on each study and the statistical data are adequately presented in most cases. Apart from this, nine of the included articles^{21-24,26-28,30,34} used parametric tests (T Student or ANOVA) in small samples (less than 30), with no information on the normality of the data.




Fig. 3 – Mean superior joint space in each study and globally.




Fig. 4 – Mean posterior joint space in each study and globally.

Pearson correlation was used by six authors,^{19,21,24,29,32,35} with no mention to the linear relation between the variables, which is necessary to the use and interpretation of this coefficient. Seven studies^{19-23,28,35} did not present an adequate sample description, with no information about the age or gender. The other eleven^{18,24-27,29-34} showed at least, the number of patients from each gender and data concerning the age of the included sample (mean, standard deviation, minimum and maximum values). The error analysis was presented in seven of the articles.^{18,21,22,26-28,33} Only one study presented the estimation of sample size.³³ Randomization was used by Tsuruta et al.¹⁸ and none of the retrieved articles presented blinding in measurements. Only one¹⁸ of the studies was classified as moderate level of evidence, as it presents randomization and adequate statistics but fails to presents blinding in measurements. All the other articles were classified as low level of evidence.

Meta-analysis results

Seventeen of the studies presented on the review were included in this meta-analysis. One study³⁴ did not present standard deviation values and so, statistical comparison with other studies was not possible.

The mean anterior joint space from the 17 considered studies was 1.86 mm (95% CI: 1.62-2.10), although high levels of heterogeneity were found among the studies (Q(16) = 1032.11; P < 0.001; I² = 98.49%) (Fig. 2).

The superior joint space presented a mean value of 2.36 mm (95% CI: 1.99-2.72), also with high levels of heterogeneity between the 14 articles that presented this value (Q(13) = 1117.08; P < 0.001; I² = 98.64%) (Fig. 3).

The posterior joint space also presented high heterogeneity among the 16 included samples (Q(15) = 440.70; P < 0.001; I² = 97.37%) with a mean value of 2.22 mm (95% CI: 2.00-2.45) (Fig. 4).




Fig. 5 – Mean difference between the anterior joint space on the right and left joints for each study and globally.




Fig. 6 – Mean difference between the superior joint space on the right and left joints for each study and globally.

The mean global differences and the differences among the studies between the right and left anterior, superior and posterior joint spaces are shown in Figs. 5–7 respectively. In all of the comparisons, the difference is close to zero.

For the anterior joint space, the mean difference is -0.10 mm (95% CI: -0.17; -0.03), without heterogeneity between the included samples (Q(9) = 9.94; P = 0.355; I² = 0.02%) (Fig. 5).

As for the mean difference between the right and left superior joint spaces, the value was 0.04 mm (95% CI: -0.07; 0.15), presenting low heterogeneity among the differences found on the different studies considered (Q(8) = 8.24; P = 0.410; I² = 19.31%) (Fig. 6).

The posterior joint space of the right and left joints showed moderate heterogeneity between the samples (Q(9) = 21.07; P = 0.012; I² = 56.48%) with a mean global difference of -0.04 mm (95% CI: -0.17; 0.10) (Fig. 7).

Discussion

Joint space measurements have been used to assess the mandibular condyle position radiographically since this method was used in laminographies.³⁶ Since then the technology has evolved so much that it is now possible to assess the joint space in 3D radiographic imaging with CT, CBCT and MRI. Therefore, a systematic review to assess the relevance of these methods and their scientific evidence is necessary. In the present study, all the articles about joint space assessment on 2D radiographic examination of the TMJ were excluded as these methods have proven lower accuracy both in the image acquisition process and in measurements, than 3D radiographic methods.³⁷ MRI was also excluded because this exam is not indicated to assess hard structures and, as both the mandibular condyle and the glenoid fossa joint space limits are mainly bone and cartilage, this is not the best exam for




Fig. 7 – Mean difference between the posterior joint space on the right and left joints for each study and globally.

accurately determine joint space measurements.³⁸ Furthermore, all the articles including extensive treatment that could significantly influence the joint space, like orthodontic treatment and splint therapy, have been excluded. Finally, studies with samples exclusively on the mixed and early permanent dentition were excluded as, according to some authors, the mandibular condyle is not completely formed before the end of the growth, in other words until 15–16 years old.³⁹ The exclusion of studies that only assessed the joint space in less than two separate points of the TMJ was due to the definition of the position of an object in space depending on three coordinates. According to this, joint space analysis done only on one point does not provide enough information to determine the position of the mandibular condyle in the glenoid fossa.

The meta-analysis showed high levels of heterogeneity among the selected studies what is possibly explained by the high heterogeneity between the selected studies: the sample sizes varies between 24²⁷ and 464 patients,³¹ there are different samples from carefully selected normal joints^{19,27} to patients presenting malocclusions^{21,26,31–33,35} or temporomandibular disorders^{23,24} and different methods^{19,25,27} were used to determine the joint spaces (as showed during the systematic review). All these factors may have contributed to high heterogeneity between the samples.

The results from this investigation also showed statistically significant differences between the right and left joints for the anterior joint space (AJS), while the opposite was found for the posterior joint space (PJS). However, the mean difference was -0.10 mm (95% CI: -0.17 ; -0.03) and, therefore, it is very close to 0. Moreover, comparison of the AJS and the PJS between the right and left joints was performed on 10 of the retrieved articles, and only two showed statistically significant differences for the AJS. The fact that one of the studies had the biggest sample ($n = 232$) with a power of 16.30% on the meta-analysis might explain the statistically significant result for the AJS. A previous study suggested that this asymmetry would be related to normally asymmetric cranial bases or side preferences during mastication.³¹ Moreover, most patients show a centric relation-centric occlusion discrepancy, usually caused by a posterior interference that is unilateral in most cases.^{40–43} As an adaptation, the condyles might move asymmetrically and while the contra-lateral condyle moves sagittally, the ipsilateral rotates to establish a more balanced dental occlusion. The absence of statistically significant differences between right and left joints for the PJS may be explained by stabilization of the TMJ posteriorly by the articular disk.

Conclusion

There is insufficient scientific evidence concerning sagittal joint spaces of the TMJ, as there are no articles with high level of evidence and only one study presents moderate level of evidence.

Although no high level of evidence studies were found, the authors decided to perform a meta-analysis of the mean sagittal joint spaces of the TMJ and the differences between the right and left joints. The mean anterior joint space was 1.86 mm (CI 95%: 1.62–2.10), the superior joint space was 2.36 mm (CI 95%: 1.99–2.72) and the posterior was 2.22 mm

(CI 95%: 2.00–2.45). However, a high level of heterogeneity was found, meaning that these results must be considered with care. However, the results of the meta-analysis suggest that the posterior joint space is larger than the anterior joint space. This result is in accordance to the concept in use that the mandibular condyle must be on the most superior-anterior position in the glenoid fossa.^{5,8–10}

The analysis of the difference between the right and left sagittal joint spaces showed statistically significant differences between the two joints in the anterior joint space, but not in the superior and posterior joint spaces.

More research with more solid methodology is needed on this topic.

Conflicts of interest

The authors have no conflicts of interest to declare.

REFERENCES

1. Posselt U. Studies in the mobility of the human mandible. *Acta Odontol Scand.* 1952;10 Suppl. 10:19–27.
2. Dawson PE. Evaluation diagnosis and treatment of occlusal problems. St Louis: Mosby; 1989.
3. Van Blarcom C, Campbell S, Can A. The glossary of prosthodontic terms. 7th ed. St Louis: Mosby; 1999.
4. Rinchuse DJ, Kandasamy S. Centric relation: a historical and contemporary orthodontic perspective. *J Am Dent Assoc.* 2006;137:494–501.
5. Dawson PE. A classification system for occlusions that relates maximal intercuspation to the position and condition of the temporomandibular joints. *J Prosthet Dent.* 1996;75:60–6.
6. Roth RH. Occlusion and condylar position. *Am J Orthod Dentofac Orthop.* 1995;107:315–8.
7. Rinchuse DJ. A three-dimensional comparison of condylar change between centric relation and centric occlusion using the mandibular position indicator. *Am J Orthod Dentofac Orthop.* 1995;107:319–28.
8. Okeson J. Management of temporomandibular disorders and occlusion. 6th ed. Mosby Elsevier; 2008.
9. Dawson PE. Centric relation. Its effect on occluso-muscle harmony. *Dent Clin N Am.* 1979;23:169–80.
10. Dawson PE. New definition for relating occlusion to varying conditions of the temporomandibular joint. *J Prosthet Dent.* 1995;74:619–27.
11. Kandasamy S, Boeddinghaus R, Kruger E. Condylar position assessed by magnetic resonance imaging after various bite position registrations. *Am J Orthod Dentofac Orthop.* 2013;144:512–7.
12. Honey OB, Scarfe WC, Hilgers MJ, Klueber K, Silveira AM, Haskell BS, et al. Accuracy of cone-beam computed tomography imaging of the temporomandibular joint: comparisons with panoramic radiology and linear tomography. *Am J Orthod Dentofac Orthop.* 2007;132:429–38.
13. Hilgers ML, Scarfe WC, Scheetz JP, Farman AG. Accuracy of linear temporomandibular joint measurements with cone beam computed tomography and digital cephalometric radiography. *Am J Orthod Dentofac Orthop.* 2005;128:803–11.
14. White S, Pharoah M. Oral radiology principles and interpretation. 5th ed. Los Angeles: Mosby; 2009.
15. Collaboration TC. Cochrane reviewers' handbook; 2004. Available from: <http://www.cochrane.org/resources/handbook>
16. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting

- systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med.* 2009;6:e1001000.
17. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327:557-60.
 18. Tsuruta A, Yamada K, Hanada K, Hosogai A, Kohno S, Koyama J, et al. The relationship between morphological changes of the condyle and condylar position in the glenoid fossa. *J Orofac Pain.* 2004;18:148-55.
 19. Vitral RW, da Silva Campos MJ, Rodrigues AF, Fraga MR. Temporomandibular joint and normal occlusion: is there anything singular about it? A computed tomographic evaluation. *Am J Orthod Dentofac Orthop.* 2011;140:18-24.
 20. Rodrigues AF, Fraga MR, Vitral RW. Computed tomography evaluation of the temporomandibular joint in Class I malocclusion patients: condylar symmetry and condyle-fossa relationship. *Am J Orthod Dentofac Orthop.* 2009;136:192-8.
 21. Rodrigues AF, Fraga MR, Vitral RW. Computed tomography evaluation of the temporomandibular joint in Class II Division 1 and Class III malocclusion patients: condylar symmetry and condyle-fossa relationship. *Am J Orthod Dentofac Orthop.* 2009;136:199-206.
 22. Henriques JC, Fernandes Neto AJ, Almeida Gde A, Machado NA, Lelis ER. Cone-beam tomography assessment of condylar position discrepancy between centric relation and maximal intercuspation. *Braz Oral Res.* 2012;26:29-35.
 23. Okur A, Ozkiris M, Kapusuz Z, Karacavus S, Saydam L. Characteristics of articular fossa and condyle in patients with temporomandibular joint complaint. *Eur Rev Med Pharmacol Sci.* 2012;16:2131-5.
 24. Pereira LJ, Gavião MB. Tomographic evaluation of TMJ in adolescents with temporomandibular disorders. *Braz Oral Res.* 2004;18:208-14.
 25. Dalili Z, Khaki N, Kia SJ, Salamat F. Assessing joint space and condylar position in the people with normal function of temporomandibular joint with cone-beam computed tomography. *Dent Res J (Isfahan).* 2012;9:607-12.
 26. Prabhat K, Kumar V, Maheshwari S, Ahmad I, Tariq M. Computed tomography evaluation of craniomandibular articulation in Class II Division 1 malocclusion and Class I normal occlusion subjects in North Indian Population. *ISRN Dent.* 2012;2012:312031.
 27. Ikeda K, Kawamura A. Assessment of optimal condylar position with limited cone-beam computed tomography. *Am J Orthod Dentofac Orthop.* 2009;135:495-501.
 28. Seren E, Akan H, Toller MO, Akyar S. An evaluation of the condylar position of the temporomandibular joint by computerized tomography in Class III malocclusions: a preliminary study. *Am J Orthod Dentofac Orthop.* 1994;105:483-8.
 29. Gianelly AA, Petras JC, Boffa J. Condylar position and Class II deep-bite, no-overjet malocclusions. *Am J Orthod Dentofac Orthop.* 1989;96:428-32.
 30. Christiansen EL, Thompson JR, Zimmerman G, Roberts D, Hasso AN, Hinshaw DB Jr, et al. Computed tomography of condylar and articular disk positions within the temporomandibular joint. *Oral Surg Oral Med Oral Pathol.* 1987;64:757-67.
 31. Cohlmlia J, Ghosh J, Sinha P, Nanda R, Currier GF. Tomographic assessment of temporomandibular joints in patients with malocclusion. *Angle Orthod.* 1996;66:27-36.
 32. Uzel A, Özyürek Y, Öztunç H. Condyle position in Class II Division 1 malocclusion patients: correlation between MPI records and CBCT images. *J World Fed Orthodont.* 2013;2:e65-70.
 33. Arieta-Miranda JM, Silva-Valencia M, Flores-Mir C, Paredes-Sampen NA, Arriola-Guillen LE. Spatial analysis of condyle position according to sagittal skeletal relationship, assessed by cone beam computed tomography. *Prog Orthod.* 2013;14:36.
 34. Alves N, Deana NF, Schilling QA, González VA, Schilling LJ, Pastenes RC. Assessment of TMJ condylar position and joint space in Chilean individuals with temporomandibular disorders. *Int J Morphol.* 2014;32:32-5.
 35. Vitral RW, Telles Cde S, Fraga MR, de Oliveira RS, Tanaka OM. Computed tomography evaluation of temporomandibular joint alterations in patients with class II division 1 subdivision malocclusions: condyle-fossa relationship. *Am J Orthod Dentofac Orthop.* 2004;126:48-52.
 36. Ricketts RM. Present status of laminagraphy as related to dentistry. *J Am Dent Assoc.* 1962;65:56-64.
 37. Quintero JC, Trosien A, Hatcher D, Kapila S. Craniofacial imaging in orthodontics: historical perspective, current status, and future developments. *Angle Orthod.* 1999;69:491-506.
 38. Commission E. Radiation protection n° 172. In: Protection D-GfEDDNEUDR, editor. Cone beam CT for dental and maxillofacial radiology: evidence-based guidelines. 2012. Luxembourg.
 39. Enlow DH. Facial growth. 3rd ed. Philadelphia: Saunders; 1990.
 40. Crawford SD. Condylar axis position, as determined by the occlusion and measured by the CPI instrument, and signs and symptoms of temporomandibular dysfunction. *Angle Orthod.* 1999;69:103-15, discussion 15-6.
 41. Cordray FE. Three-dimensional analysis of models articulated in the seated condylar position from a deprogrammed asymptomatic population: a prospective study. Part 1. *Am J Orthod Dentofac Orthop.* 2006;129:619-30.
 42. Cordray FE. The importance of the seated condylar position in orthodontic correction. *Quintessence Int.* 2002;33:284-93.
 43. Ponces MJ, Tavares JP, Lopes JD, Ferreira AP. Comparison of condylar displacement between three biotypological facial groups by using mounted models and a mandibular position indicator. *Korean J Orthod.* 2014;44:312-9.