

CLUSTERING OF EXTREME EVENTS CREATED BY MULTIPLE CORRELATED MAXIMA

DAVIDE AZEVEDO, ANA CRISTINA MOREIRA FREITAS, JORGE MILHAZES FREITAS,
AND FAGNER B. RODRIGUES

ABSTRACT. We consider stochastic processes arising from dynamical systems by evaluating an observable function along the orbits of the system. The novelty is that we will consider observables achieving a global maximum value (possibly infinite) at multiple points with special emphasis for the case where these maximal points are correlated or bound by belonging to the same orbit of a certain chosen point. These multiple correlated maxima can be seen as a new mechanism creating clustering of extreme observations, *i.e.*, the occurrence of several extreme observations concentrated in the time frame. We recall that clustering was intimately connected with periodicity when the maximum was achieved at a single point. We will study this mechanism for creating clustering and will address the existence of limiting Extreme Value Laws, the repercussions on the value of the Extremal Index, the impact on the limit of Rare Events Points Processes, the influence on clustering patterns and the competition of domains of attraction. We also consider briefly and for comparison purposes multiple uncorrelated maxima. The systems considered include expanding maps of the interval such as Rychlik maps but also maps with an indifferent fixed point such as Manneville-Pommeau maps.

1. INTRODUCTION

In the past few years, the study of Extreme Value Theory (EVT) for dynamical systems has been a subject of much interest. We mention in particular the inspiring paper of Collet [Col01] and refer to [Fre13] for a review on further developments and references. Extreme events (which occur with small probability and for that reason are also called rare events) are characterised by abnormally high observations that are identified as exceedances of high thresholds.

In this dynamical setting the observation data comes from starting the system at a certain initial state and evaluate a certain observable function along the subsequent states through which the system goes while time goes by. Then, exceedances of a high threshold correspond

Date: April 25, 2016.

2010 *Mathematics Subject Classification.* 37A50, 60G55, 60G70, 37B20, 60G10, 37C25.

Corresponding author: JMF; Tel.: +351 220402158; Fax: +351 220402108; E-mail: jmfreita@fc.up.pt .

ACMF was partially supported by FCT grant SFRH/BPD/66040/2009. JMF was partially supported by FCT grant SFRH/BPD/66174/2009. Both these grants are financially supported by the program POPH/FSE. DA, ACMF, JMF are partially supported by FCT (Portugal) projects PTDC/MAT/120346/2010 and FAPESP/19805/2014, which are funded by national and European structural funds through the programs FEDER and COMPETE. FBR was supported by BREUDS, a Brazilian-European partnership of the FP7-PEOPLE-2012-IRSES program, with project number 318999, which is supported by an FP7 International International Research Staff Exchange Scheme (IRSES) grant of the European Union. All authors were partially supported by CMUP (UID/MAT/00144/2013), which is funded by FCT (Portugal) with national (MEC) and European structural funds through the programs FEDER, under the partnership agreement PT2020. All authors thank Mike Todd for careful reading and suggestions.

to entrances or hits of the orbits of the system to some designated target sets on the phase space. As the threshold increases, the respective target sets shrink. This fact explains the connection between the existence of distributional limits, called Extreme Value Laws (EVL), for the partial maximum of observations when the thresholds increase to the possible maximum value, and the existence of distributional limits, called Hitting Times Statistics (HTS), for the waiting time before hitting the corresponding target sets as they shrink. This connection hinted in [Col01] was formally established in [FFT10, FFT11].

In the study of HTS, in most cases, the neighbourhoods are either cylinder sets or metric balls and shrink to a point ζ in the phase space. In the study of EVL the observable φ has a global maximum at ζ and, in almost all cases, the exceedances correspond to metric balls around ζ . The limiting laws for HTS and EVL were proved to be the same in [FFT10, FFT11] and typically one obtains a standard exponential distribution $H(\tau) = 1 - e^{-\tau}$, with $\tau \geq 0$. The word typically is used in the sense that for almost every point ζ (with respect to the invariant measure) one obtains a standard exponential HTS and EVL. See for example [HSV99, Col01, BSTV03, HNT12, CC13, PS14] and [Sau09] for an excellent review. In this case, the rule is the absence of clustering of exceedances or hits to the target sets. This is perceived by realising that the exceedances (hits) appear as isolated occurrences scattered along the time frame.

On the other hand, since the work of Hirata [Hir93] on Axiom A diffeomorphisms, it is known that for cylinder target sets based on periodic points, *i.e.*, when ζ is a periodic point, one observes clustering of hits and a parameter $0 \leq \theta \leq 1$ appears in the asymptotic distribution: $H(\tau) = 1 - e^{-\theta\tau}$. This observation was further developed in the paper [HV09], where a compound Poisson distribution for the number of hits to target sets composed of unions of dynamical cylinders was obtained for ψ mixing measures. Then in [FFT12], using the relation between HTS and EVL, a new technique based on suitably adjusted dependent conditions allowed to show the existence of EVL (and consequently HTS) in the case ζ is a periodic point and the target sets are metric balls for systems with a strong form of decay of correlations. Due to the connection with EVL the parameter $0 \leq \theta \leq 1$ appearing in the limit distribution $H(\tau) = 1 - e^{-\theta\tau}$ was identified to be the Extremal Index (EI), a parameter already appearing in the context of classical EVT. In the presence of clustering, long time series reveal several separated clusters or groups of exceedances, *i.e.*, instead of the exceedances appearing as isolated events they appear concentrates in bulks of nearly consecutive abnormally high observations. The EI ‘measures’ the intensity of clustering. An EI equal to 1 means no clustering and an EI close to 0 means intense clustering. In fact, in most cases the EI can be seen as the inverse of the average cluster size. We remark that, in some less frequent situations, lack of tightness can lead to counter examples (see [Smi88]). Nevertheless, throughout this paper, the EI will always coincide with inverse of average cluster size.

Moreover, in [FFT12], for uniformly expanding systems such as the doubling map, a dichotomy was shown which states that either ζ is periodic and we have an EI with a very precise formula depending on the expansion rate at ζ , or for every non-periodic ζ , we have an EI equal to 1 (which means no clustering). The dichotomy was then obtained for more general systems such as: conformal repellers [FP12], systems with spectral gaps for the Perron-Frobenius operator [Kel12] and systems with strong decay of correlations [AFV14].

The study of EVL and HTS can be enhanced by considering Rare Events Point Processes (REPP). These point processes keep record of the exceedances of the high thresholds by counting the number of such exceedances on a rescaled time interval. At typical points it was known that the REPP converge in distribution to a standard Poisson process. From [HV09], at

periodic points, we expect that the REPP converge to compound Poisson process instead. In [FFT13], a study regarding the convergence of REPP was performed for metric balls as target sets around periodic points ζ and for non-uniformly hyperbolic systems. The limit process obtained for the convergence in distribution of the REPP in [FFT13] was a compound Poisson process which could be described as combination of Poisson process ruling the positions of the clusters in the time line being that these positions are marked by the cluster sizes ruled by a geometric multiplicity distribution. Both components are determined by the value of EI θ .

Again, as proved in [AFV14], for continuous systems with strong decay of correlations a dichotomy regarding the convergence of REPP holds: either it converges to a compound Poisson distribution at repelling periodic points ζ , with a geometric multiplicity distribution, or it converges to a standard Poisson process at every other non-periodic point ζ . Moreover, as already seen in [FFT13], the clustering pattern observed at repelling periodic points ζ obeys a very rigid pattern: it consists of ‘bulk’ of strictly decreasing exceedances observed at precise fixed times, corresponding to the period.

Contrary to the customary case studies, in which the set of maximising points \mathcal{M} of the observable φ is reduced to a single point ζ , in this paper we will consider the case of multiple maxima, with special emphasis for the case of correlated maxima, where \mathcal{M} consists of finite number of points bound together by belonging to the same orbit. This binding gives a mechanism of creating clustering of exceedances, which we will study and explore below. We will also consider the case of uncorrelated maxima but the focus will be turned to the correlated case.

We remark that in the literature not all examples regarding the study of rare events are reduced to the cases in which the target sets shrink to a point. For example, in [CCC09] the authors consider a sets generated by a proper subshift of finite type of a one-sided irreducible and aperiodic shift of finite type and in [KL09] the authors consider target sets shrinking to the diagonal of the phase space of a system obtained by coupling two expanding interval maps. The setting which probably most resembles ours is that studied in [HNT12], where the authors, in the EVL approach, consider observables with multiple maxima that are chosen independently as typical points for the invariant probability measure. In this case the maxima are uncorrelated and they obtained the standard exponential distribution as limiting EVL, which means that there is no clustering of exceedances.

In classical Extreme Value Theory, it is usual to use normalising thresholds $(u_n)_{n \in \mathbb{N}}$ that are linear sequences depending on a parameter y , say $u_n = y/a_n + b_n$. Then the folklore theorem (see [LLR83]) asserts that there are only three types of limiting distributions for the partial maxima linearly normalised, namely: Gumbel, Fréchet or Weibull. Moreover, there is relation between the tail of the distribution function that governs the random variables and the type of EVL that applies. This relation can be summarised, in rough words, as follows: heavy tails fall in the domain of attraction of the Fréchet type distributions; light tails fall are linked to the Gumbel distribution; finite tails are typically associated with Weibull type distributions.

This connection between the tails and the domains of attraction for maxima reveals of great importance in statistics of extremes where the estimation of the behaviour of the tail is crucial for several applications such as to Risk Theory, Insurance, Finance, etc. (see [BGTS04]). Hence, EVL techniques allow to recover such information by performing, for example, a distributional fitting to the sampled maxima.

As explained in [FFT10], the behaviour of the observable φ as a function of the distance to ζ is responsible for determining the type of EVL that applies. Hence, an interesting question

that immediately arises when considering multiple maxima is the competition between the different types.

By taking multiple maxima over the same orbit we will see the following consequences:

- (1) appearance of clustering not caused necessarily by periodic orbits;
- (2) the possibility of creating different clustering patterns (not reduced to a bulk of strictly decreasing observations over the threshold);
- (3) the possibility of affecting the EI when we already consider periodic points;
- (4) different multiplicity distributions for the limit of REPP;
- (5) competition between different types of distributions.

The systems considered include systems with decay of correlations against L^1 observables such as Rychlik maps [Ryc83] and piecewise uniformly expanding maps in higher dimensions studied in [Sau00] but also non-uniformly expanding maps such as intermittent maps.

We believe that this study will be a precursor of further developments regarding more sophisticated maximal sets \mathcal{M} , that could be, for example, submanifolds of codimension 1. In fact, some work in that direction is already being done by some of the authors and we believe it carries a large potential of applications since it will more easily accommodate physical observables in applications like to meteorology, where the set \mathcal{M} plays the role of a critical set, which is the source of abnormal and possibly catastrophic events. Understanding the geometry and the recurrence properties of \mathcal{M} , as we will see here in a much simpler context, will provide knowledge about the extremal behaviour like for example the clustering patterns that could help devise early warning systems.

An example of possible application is to structural failures. The anticipated study of the type of the clustering patterns of a certain natural phenomenon is of crucial importance, on one side to predict the likelihood of such failure and on, another side, to help designing the material and structures to stand stronger against the natural causes they have to face.

The unfolding of such possibilities will give deeper understanding of the possible outputs of EVT for dynamical systems, as it will open a door for modelling physical phenomena with some underlying periodic effect, which is often difficult to perceive. Moreover, again, these short recurrence mechanisms have an enormous potential as a source of examples for the classical EVT of stochastic processes and serving as a model for several practical situations.

2. THE SETTING AND BACKGROUND

Take a system $(\mathcal{X}, \mathcal{B}, \mu, f)$, where \mathcal{X} is a Riemannian manifold, \mathcal{B} is the Borel σ -algebra, $f : \mathcal{X} \rightarrow \mathcal{X}$ is a measurable map and μ an f -invariant probability measure. Suppose that the time series X_0, X_1, \dots arises from such a system simply by evaluating a given observable $\varphi : \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ along the orbits of the system, or in other words, the time evolution given by successive iterations by f :

$$X_n = \varphi \circ f^n, \quad \text{for each } n \in \mathbb{N}. \tag{1}$$

Clearly, X_0, X_1, \dots defined in this way is not necessarily an independent sequence. However, f -invariance of μ guarantees that this stochastic process is stationary.

We suppose that the r.v. $\varphi : \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ has N global maxima $\xi_1, \dots, \xi_N \in \mathcal{X}$ (we allow $\varphi(\xi_1) = +\infty$) and assume they all belong to the orbit of the point ζ . (See (18) below). We are particularly interested in the events $\{X_j > u\}$ corresponding to exceedances of high

thresholds u and for that reason we suppose φ and μ are sufficiently regular in the following sense:

(R1) for u sufficiently close to $u_F := \varphi(\xi_i)$ ($i \in \{1, \dots, N\}$),

$$U(u) := \{x \in \mathcal{X} : \varphi(x) > u\} = \{X_0 > u\}$$

corresponds to a union of balls centered at the points ξ_i , i.e., $U(u) = \bigcup_{i=1}^N B_{\varepsilon_i}(\xi_i)$ with $\varepsilon_i = \varepsilon_i(u)$. Moreover, the quantity $\mu(U(u))$, as a function of u , varies continuously on a neighbourhood of u_F .

Remark 2.1. In the vicinity of each maximal point ξ_i , the observable φ can be thought as $\varphi(x) = h_i(\text{dist}(x, \xi_i))$, where h_i is a strictly decreasing bijection defined on a neighbourhood of 0 (see Section 3.1 for more details). We remark that there is plenty of freedom in the choice of h_i and there is no restriction in the choice of the metric used, which of course needs not to be the usual Euclidean metric. This means that the connected components of the level sets $\{X_0 > u\}$ will be diffeomorphic to an Euclidean ball but may assume many different shapes.

2.1. Extreme Value Laws. In this paper, we will use an extreme value approach rather than an hitting times approach, which we have already mentioned to be two sides of the same coin as can be fully appreciated in [FFT10, FFT11].

We are interested in studying the extremal behaviour of the stochastic process X_0, X_1, \dots which is tied to the occurrence of exceedances of high levels u . The occurrence of an exceedance at time $j \in \mathbb{N}_0$ means that the event $\{X_j > u\}$ occurs, where u is close to u_F . Observe that a realisation of the stochastic process X_0, X_1, \dots is achieved if we pick, at random and according to the measure μ , a point $x \in \mathcal{X}$, compute its orbit and evaluate φ along it. Then, saying that an exceedance occurs at time j means that the orbit of the point x hits one of the balls in $U(u)$ at time j , i.e., $f^j(x) \in B_{\varepsilon_i}(\xi_i)$ for some $i \in \{1, \dots, N\}$.

Given a stochastic process X_0, X_1, \dots we define a new sequence of random variables M_1, M_2, \dots given by

$$M_n = \max\{X_0, X_1, \dots, X_{n-1}\}. \tag{2}$$

On the independent context, i.e., when the stochastic process X_0, X_1, \dots is a sequence of independent and identically distributed (i.i.d.) r.v., the first statement regarding M_n is that M_n converges almost surely (a.s.) to u_F . Then, the next natural question is whether we can find a distributional limit for M_n , when conveniently normalised.

Definition 2.2. We say that we have an *Extreme Value Law* (EVL) for M_n if there is a non-degenerate d.f. $H : \mathbb{R} \rightarrow [0, 1]$ with $H(0) = 0$ and, for every $\tau > 0$, there exists a sequence of levels $u_n = u_n(\tau)$, $n = 1, 2, \dots$, such that

$$n\mu(X_0 > u_n) \rightarrow \tau, \text{ as } n \rightarrow \infty, \tag{3}$$

and for which the following holds:

$$\mu(M_n \leq u_n) \rightarrow \bar{H}(\tau), \text{ as } n \rightarrow \infty. \tag{4}$$

where the convergence is meant at the continuity points of $H(\tau)$, and $\bar{H}(\tau) = 1 - H(\tau)$.

The motivation for using a normalising sequence $(u_n)_{n \in \mathbb{N}}$ satisfying (3) comes from the case when X_0, X_1, \dots are independent and identically distributed (i.i.d.). In this setting, it is clear that $\mu(M_n \leq u) = (F(u))^n$, where F is the d.f. of X_0 . Hence, condition (3) implies that

$$\mu(M_n \leq u_n) = (1 - \mu(X_0 > u_n))^n \sim \left(1 - \frac{\tau}{n}\right)^n \rightarrow e^{-\tau}, \tag{5}$$

as $n \rightarrow \infty$. Moreover, the reciprocal is also true (see [LLR83, Theorem 1.5.1] for more details). Note that in this case $H(\tau) = 1 - e^{-\tau}$ is the standard exponential d.f.

2.2. The existence of Extreme Value Laws. In [FFT15], the authors synthesised the conditions in [FF08] and in [FFT12] to obtain a couple of general conditions that apply both in the presence and absence of clustering. These conditions are particularly tailored to the application of dynamical systems and follow from a strong form of decay of correlations (against L^1 observables) to be defined below. Since the original conditions of Leadbetter were denoted by a capital D and in the literature there is now a profusion of upper and lower indices appended to D and Δ for adaptations of the original Leadbetter's conditions, in [FFT15] the authors chose a Cyrillic D , *i.e.*, the symbol \mathbb{D} , to denote the new conditions that we recall below.

In what follows for every $A \in \mathcal{B}$, we denote the complement of A as $A^c := \mathcal{X} \setminus A$.

For some $u \in \mathbb{R}$, $q \in \mathbb{N}$, we define the events:

$$\begin{aligned} U(u) &:= \{X_0 > u\}, \\ \mathcal{A}_q(u) &:= U(u) \cap \bigcap_{i=1}^q f^{-i}(U(u)^c) = \{X_0 > u, X_1 \leq u, \dots, X_q \leq u\}. \end{aligned} \quad (6)$$

where $\mathcal{A}_q(u)$ corresponds to the case where we have an extreme event at time zero that is not followed by another one up to time $t = q$. This is a condition clearly pointing to the absence of clustering. In fact, the occurrence of $f^{-i}(\mathcal{A}_q(u))$ marks the end of a cluster of exceedances at time i because, for a good choice of q , we have that, within a cluster, no exceedance can be separated from another exceedance by more than q units of time. The notation f^{-i} is used for the preimage by f^i . We refer to the events $f^{-i}(\mathcal{A}_q(u))$ as the occurrence of an *escape* at time i , since the orbits that after i iterates fall into $\mathcal{A}_q(u)$ produce an exceedance at time i that escapes the underlying periodic effect and is not followed by any other exceedance in the same cluster. We also set $\mathcal{A}_0(u) := U(u)$ and for the given sequence of levels $u_n = u_n(\tau)$, $\mathcal{A}_{q,n} := \mathcal{A}_q(u_n)$, for all $n \in \mathbb{N}$ and $q \in \mathbb{N}_0$. Let

$$\theta_n := \frac{\mu(\mathcal{A}_{q,n})}{\mu(U(u_n))}. \quad (7)$$

Note that θ_n ‘measures’ the proportion of points from U_n that escape the underlying periodic effect and do not produce another exceedance in the same cluster. Let $B \in \mathcal{B}$ be an event. For some $s \geq 0$ and $\ell \geq 0$, we define:

$$\mathcal{W}_{s,\ell}(B) = \bigcap_{i=\lfloor s \rfloor}^{\lfloor s \rfloor + \max\{\lfloor \ell \rfloor - 1, 0\}} f^{-i}(B^c), \quad (8)$$

which is a compact notation for the events that correspond to the non occurrence of B during the time frame $[s, s + \ell)$. We will write $\mathcal{W}_{s,\ell}^c(B) := (\mathcal{W}_{s,\ell}(B))^c$. Whenever is clear or unimportant which event $B \in \mathcal{B}$ applies, we will drop the B and write just $\mathcal{W}_{s,\ell}$ or $\mathcal{W}_{s,\ell}^c$. For a set $B \in \mathcal{B}$ we define, the *first hitting time* to B denoted by $r_B : \mathcal{X} \rightarrow \mathbb{N}$, where

$$r_B(x) = \min \{j \in \mathbb{N} \cup \{+\infty\} : f^j(x) \in B\}, \quad (9)$$

which corresponds to the first entrance of the orbit of x into B . Note that

$$\mathcal{W}_{0,n}(U(u)) = \{M_n \leq u\} \quad \text{and} \quad f^{-1}(\mathcal{W}_{0,n}(B)) = \{r_B > n\}. \quad (10)$$

We now give the first condition \mathbb{D} which is a special type of mixing condition imposing an asymptotic independence of events related to $\mathcal{A}_{q,n}$.

Condition ($\mathbb{D}_q(u_n)$). We say that $\mathbb{D}(u_n)$ holds for the sequence X_0, X_1, \dots if for every $\ell, t, n \in \mathbb{N}$,

$$|\mu(\mathcal{A}_{q,n} \cap \mathcal{W}_{t,\ell}(\mathcal{A}_{q,n})) - \mu(\mathcal{A}_{q,n})\mu(\mathcal{W}_{0,\ell}(\mathcal{A}_{q,n}))| \leq \gamma(q, n, t), \quad (11)$$

where $\gamma(q, n, t)$ is decreasing in t for each n and, there exists a sequence $(t_n)_{n \in \mathbb{N}}$ such that $t_n = o(n)$ and $n\gamma(q, n, t_n) \rightarrow 0$ when $n \rightarrow \infty$.

The next condition precludes the occurrence of events of the type $\mathcal{A}_{q,n}$ concentrated in the time frame, *i.e.*, it forces the occurrence of such events to appear scattered in time. Note that when $q = 0$, we have $\mathcal{A}_{q,n} = U_n$, which means that, in this case, the next condition can be seen as an anti-clustering condition, which forbids the occurrence of clustering of exceedances.

For some fixed $q \in \mathbb{N}_0$, consider the sequence $(t_n)_{n \in \mathbb{N}}$, given by condition $\mathbb{D}(u_n)$ and let $(k_n)_{n \in \mathbb{N}}$ be another sequence of integers such that

$$k_n \rightarrow \infty \quad \text{and} \quad k_n t_n = o(n). \quad (12)$$

Condition ($\mathbb{D}'_q(u_n)$). We say that $\mathbb{D}'_q(u_n)$ holds for the sequence X_0, X_1, X_2, \dots if there exists a sequence $(k_n)_{n \in \mathbb{N}}$ satisfying (12) and such that

$$\lim_{n \rightarrow \infty} n \sum_{j=q+1}^{\lfloor n/k_n \rfloor - 1} \mu(\mathcal{A}_{q,n} \cap f^{-j}(\mathcal{A}_{q,n})) = 0. \quad (13)$$

From [FFT15, Corollary 2.4] follows that if the stochastic process satisfies both conditions $\mathbb{D}_q(u_n)$ and $\mathbb{D}'_q(u_n)$ for some $q \in \mathbb{N}_0$, where $(u_n)_{n \in \mathbb{N}}$ satisfies (3), then $\lim_{n \rightarrow \infty} \mu(M_n \leq u_n) = e^{-\theta\tau}$, whenever the limit $\theta = \lim_{n \rightarrow \infty} \theta_n$ exists.

The role of the conditions \mathbb{D}_q and \mathbb{D}'_q above in the proof of the last statement is connected to a blocking argument that splits the random variables of the process into different blocks. Then condition \mathbb{D}_q is used to show that the occurrence of escapes in a certain block is nearly independent from the same occurrence in a different block. Finally, condition \mathbb{D}'_q is used to estimate the probability of having no escapes in each block, which ultimately leads to the desired limit distribution. See [FFT15] for more details.

In this approach, it is rather important to observe the prominent role played by condition $\mathbb{D}'_q(u_n)$. In particular, note that if condition $\mathbb{D}'_q(u_n)$ holds for some particular $q = q_0 \in \mathbb{N}_0$, then condition $\mathbb{D}'_q(u_n)$ holds for all $q \geq q_0$, which also implies that if the limit of θ_n in (7) exists for such $q = q_0$ it will also exist for all $q \geq q_0$ and takes always the same value. This suggests that in trying to find the existence of EVL, one should try the values $q = q_0$ until we find the smallest one that makes $\mathbb{D}'_q(u_n)$ hold. With that purpose, as in [FFT15, Theorem 3.7], we consider the following. Let $A \in \mathcal{B}$. Recall the definition of the first hitting time to A , presented in (9), $r_A(x) = \min \{j \in \mathbb{N} \cup \{+\infty\} : f^j(x) \in A\}$. The restriction of r_A to A is called the *first return time function* to A . We define the *first return time* to A , which we denote by $R(A)$, as the infimum of the return time function to A , *i.e.*,

$$R(A) = \inf_{x \in A} r_A(x). \quad (14)$$

Assume that there exists $q \in \mathbb{N}_0$ such that

$$q := \min \left\{ j \in \mathbb{N}_0 : \lim_{n \rightarrow \infty} R(\mathcal{A}_{j,n}) = \infty \right\}. \quad (15)$$

Then such q is the natural candidate to try to show the validity of $\mathbb{D}'_q(u_n)$.

2.3. Rare Events Point Processes. A more sophisticated way of studying rare events consists in studying Rare Events Point Processes (REPP). These point processes keep record of the exceedances of the high thresholds u_n by counting the number of such exceedances on a rescaled time interval. For every $A \subset \mathbb{R}$ we define

$$\mathcal{N}_{u_n}(A) := \sum_{i \in A \cap \mathbb{N}_0} \mathbf{1}_{X_i > u_n}.$$

In order to provide a proper framework of the problem we introduce next the necessary formalism to state the results regarding the convergence of point processes. We recommend the book of Kallenberg [Kal86] for further reading.

Consider the interval $[0, \infty)$ and its Borel σ -algebra $\mathcal{B}_{[0, \infty)}$. Let $x_1, x_2, \dots \in [0, \infty)$ and define

$$\nu = \sum_{i=1}^{\infty} \delta_{x_i},$$

where δ_{x_i} is the Dirac measure at x_i , *i.e.*, for every $A \in \mathcal{B}_{[0, \infty)}$, we have that $\delta_{x_i}(A) = 1$ if $x_i \in A$ or $\delta_{x_i}(A) = 0$, otherwise. The measure ν is said to be a counting measure on $[0, \infty)$. Let $\mathcal{M}_p([0, \infty))$ be the space of counting measures on $([0, \infty), \mathcal{B}_{[0, \infty)})$. We equip this space with the vague topology, *i.e.*, $\nu_n \rightarrow \nu$ in $\mathcal{M}_p([0, \infty))$ whenever $\nu_n(\psi) \rightarrow \nu(\psi)$ for any continuous function $\psi : [0, \infty) \rightarrow \mathbb{R}$ with compact support. A *point process* N on $[0, \infty)$ is a random element of $\mathcal{M}_p([0, \infty))$, *i.e.*, let (X, \mathcal{B}_X, μ) be a probability space, then any measurable map $N : X \rightarrow \mathcal{M}_p([0, \infty))$ is a point process on $[0, \infty)$.

To give a concrete example of a point process, which in particular will appear as the limit of the REPP, we consider:

Definition 2.3. Let T_1, T_2, \dots be an i.i.d. sequence of r.v. with common exponential distribution of mean $1/\theta$. Let D_1, D_2, \dots be another i.i.d. sequence of r.v., independent of the previous one, and with d.f. π . Given these sequences, for $J \in \mathcal{B}_{[0, \infty)}$, set

$$N(J) = \int \mathbf{1}_J d \left(\sum_{i=1}^{\infty} D_i \delta_{T_1 + \dots + T_i} \right),$$

where δ_t denotes the Dirac measure at $t > 0$. Let X denote the space of all possible realisations of T_1, T_2, \dots and D_1, D_2, \dots , equipped with a product σ -algebra and measure, then $N : X \rightarrow \mathcal{M}_p([0, \infty))$ is a point process which we call a compound Poisson process of intensity θ and multiplicity d.f. π .

Remark 2.4. Throughout the paper, the multiplicity will always be integer valued which means that π is completely defined by the values $\pi_k = \mathbb{P}(D_1 = k)$, for every $k \in \mathbb{N}_0$. Note that, if $\pi_1 = 1$ and $\theta = 1$, then N is the standard Poisson process and, for every $t > 0$, the random variable $N([0, t])$ has a Poisson distribution of mean t .

In order to define the REPP we need to rescale time. We do it using the factor $v_n := 1/\mathbb{P}(X_0 > u_n)$ given by Kac's Theorem. However, before we give the definition, we need some more formalism. Let \mathcal{S} denote the semi-ring of subsets of \mathbb{R}_0^+ whose elements are intervals of the type $[a, b)$, for $a, b \in \mathbb{R}_0^+$. Let \mathcal{R} denote the ring generated by \mathcal{S} . Recall that for every $J \in \mathcal{R}$ there are $k \in \mathbb{N}$ and k intervals $I_1, \dots, I_k \in \mathcal{S}$ such that $J = \cup_{i=1}^k I_i$. In order to fix notation, let $a_j, b_j \in \mathbb{R}_0^+$ be such that $I_j = [a_j, b_j) \in \mathcal{S}$. For $I = [a, b) \in \mathcal{S}$ and $\alpha \in \mathbb{R}$, we denote $\alpha I := [\alpha a, \alpha b)$ and $I + \alpha := [a + \alpha, b + \alpha)$. Similarly, for $J \in \mathcal{R}$ define $\alpha J := \alpha I_1 \cup \dots \cup \alpha I_k$ and $J + \alpha := (I_1 + \alpha) \cup \dots \cup (I_k + \alpha)$.

Definition 2.5. We define the *rare event point process* (REPP) by counting the number of exceedances during the (rescaled) time period $v_n J \in \mathcal{R}$, where $J \in \mathcal{R}$. To be more precise, for every $J \in \mathcal{R}$, set

$$N_n(J) := \mathcal{N}_{u_n}(v_n J) = \sum_{j \in v_n J \cap \mathbb{N}_0} \mathbf{1}_{X_j > u_n}. \quad (16)$$

We will see that the REPP considered here converge to a compound Poisson process. For completeness, we define here what we mean by convergence of point processes (see [Kal86] for more details).

Definition 2.6. Let $(N_n)_{n \in \mathbb{N}} : X \rightarrow \mathcal{M}_p([0, \infty))$ be a sequence of point processes defined on a probability space (X, \mathcal{B}_X, μ) and let $N : Y \rightarrow \mathcal{M}_p([0, \infty))$ be another point process defined on a possibly distinct probability space (Y, \mathcal{B}_Y, ν) . We say that N_n converges in distribution to N if $\mu \circ N_n^{-1}$ converges weakly to $\nu \circ N^{-1}$, *i.e.*, for every continuous function φ defined on $\mathcal{M}_p([0, \infty))$, we have $\lim_{n \rightarrow \infty} \int \varphi d\mu \circ N_n^{-1} = \int \varphi d\mu \circ N^{-1}$. We write $N_n \xrightarrow{\mu} N$.

Remark 2.7. It can be shown that $(N_n)_{n \in \mathbb{N}}$ converges in distribution to N if the sequence of vector r.v. $(N_n(J_1), \dots, N_n(J_k))$ converges in distribution to $(N(J_1), \dots, N(J_k))$, for every $k \in \mathbb{N}$ and all $J_1, \dots, J_k \in \mathcal{S}$ such that $N(\partial J_i) = 0$ a.s., for $i = 1, \dots, k$.

Note that

$$\{\mathcal{N}_{u_n}([0, n]) = 0\} = \{M_n \leq u_n\}, \quad (17)$$

hence the limit distribution of M_n can be easily recovered from the convergence of the REPP.

2.4. The convergence of REPP. In [FFR], the authors are preparing a synthesised version of the conditions in [FFT10] and in [FFT13] in order to obtain a couple of general conditions that apply both in the presence and absence of clustering. Moreover, these conditions, as the previous $\mathbb{D}_q(u_n)$ and $\mathbb{D}'(u_n)$, are particularly tailored to the application of dynamical systems and follow from a strong form of decay of correlations (against L^1 observables) to be defined below.

Before we state the next condition we introduce some notation which will be useful in the rest of the paper. Let $(U^{(\kappa)}(u))_{\kappa \geq 0}$ be a sequence of nested open sets defined by:

$$U^{(0)}(u) := U(u)$$

$$\mathcal{A}_q^{(0)}(u) := \{X_0 > u, X_1 \leq u, \dots, X_q \leq u\} = U^{(0)}(u) \cap \bigcap_{i=1}^q f^{-i} \left((U^{(0)}(u))^c \right).$$

$$U^{(\kappa)}(u) := U^{(\kappa-1)}(u) - \mathcal{A}_q^{(\kappa-1)}(u) \quad \text{and} \quad \mathcal{A}_q^{(\kappa)}(u) := U^{(\kappa)}(u) \cap \bigcap_{i=1}^q f^{-i} \left((U^{(\kappa)}(u))^c \right).$$

Let

$$\theta_n := \frac{\mu(\mathcal{A}_{q,n}^{(0)})}{\mu(U(u_n))} \quad \text{and} \quad \pi_n(\kappa) = \frac{\mu(\mathcal{A}_{q,n}^{(\kappa)}) - \mu(\mathcal{A}_{q,n}^{(\kappa+1)})}{\mu(\mathcal{A}_{q,n}^{(0)})}.$$

We notice that the set $\mathcal{A}_{q,n}^{(0)}$ was defined in Section 2.2 and named $\mathcal{A}_{q,n}$. We note that condition $\mathbb{D}_q(u_n)$ was optimised to impose the necessary memory loss in order to prove the existence of an EVL. However, when dealing with point processes we need to have mixing for more sophisticated events and hence introduce the adapted condition:

Condition $(\mathbb{D}_q(u_n)^*)$. We say that $\mathbb{D}_q(u_n)^*$ holds for the sequence X_0, X_1, X_2, \dots if for any integers $t, \kappa_1, \dots, \kappa_\zeta, n$ and any $J = \cup_{i=2}^\zeta I_j \in \mathcal{R}$ with $\inf\{x : x \in J\} \geq t$,

$$\left| \mu \left(\mathcal{A}_{q,n}^{(\kappa_1)} \cap \left(\cap_{j=2}^\zeta \mathcal{N}_{u_n}(I_j) = \kappa_j \right) \right) - \mu \left(\mathcal{A}_{q,n}^{(\kappa_1)} \right) \mu \left(\cap_{j=2}^\zeta \mathcal{N}_{u_n}(I_j) = \kappa_j \right) \right| \leq \gamma(n, t),$$

where for each n we have that $\gamma(n, t)$ is nonincreasing in t and $n\gamma(n, t_n) \rightarrow 0$ as $n \rightarrow \infty$, for some sequence $t_n = o(n)$.

In [FFR] the authors show that under conditions $\mathbb{D}_q(u_n)^*$ and $\mathbb{D}'_q(u_n)$, where $(u_n)_n$ satisfies (3), the REPP $(N_n)_{n \in \mathbb{N}}$ converges in distribution to a point process N whenever the limits $\theta = \lim_{n \rightarrow \infty} \theta_n$ and $\pi(\kappa) = \lim_{n \rightarrow \infty} \pi_n(\kappa)$ exist (for all $\kappa \in \mathbb{N}$).

If one assumes that the maximum of f occurs in a repelling periodic point ζ and the probability μ is sufficiently regular then

(R2) the fact that ζ is repelling means that we have backward contraction implying that there exists $0 < \theta < 1$ so that $\cap_{j=0}^i f^{-jp}(X_0 > u)$ is another ball of smaller radius around ζ with

$$\mu \left(\bigcap_{j=0}^i f^{-jp}(X_0 > u) \right) \sim (1 - \theta)^i \mu(X_0 > u),$$

for all u sufficiently large.

Condition (R2) guarantees that $\theta_n = \frac{\mu(\mathcal{A}_{q,n}^{(0)})}{\mu(U(u_n))}$ converges to θ . Moreover, by [FFT13, Theorem 1] the REPP $(N_n)_{n \in \mathbb{N}}$ converges in distribution to a compound Poisson process N with intensity θ and multiplicity distribution given by $\pi(\kappa) = \theta(1 - \theta)^{\kappa-1}$.

3. MULTIPLE CORRELATED MAXIMA

In this paper the novelty of the approach resides in the fact that instead of considering observables $\varphi : \mathcal{X} \rightarrow \mathbb{R}$ achieving a global maximum at a single point $\zeta \in \mathcal{X}$, we assume that the maximum is achieved at N points denoted by $\xi_1, \xi_2, \dots, \xi_N$, where in almost all cases (except in Sections 4.4 and 5.4), all these points are correlated by belonging to the orbit of a certain point $\zeta \in \mathcal{X}$, *i.e.*, there exist $m_1 < m_2 < \dots < m_N \in \mathbb{N}_0$ such that

$$\xi_i = f^{m_i}(\zeta), \tag{18}$$

after possible relabelling of the maximal points. Let $\mathcal{M} := \{\xi_1, \xi_2, \dots, \xi_N\}$ be the set of maximal points of φ . We will see that the binding between the maximal points expressed in (18) is responsible for a fake periodic behaviour [FFT15, p.679] which creates clustering of exceedances. Moreover, this mechanism of creating clustering turns out to be more flexible and allows to obtain easily different multiplicity distributions for the limit of the REPP (we recall that at repelling periodic points one obtains a Geometric multiplicity distribution) and different clustering patterns (we recall that at repelling periodic points the clusters consist of a strictly decreasing bulk of exceedances occurring at precise periodic intervals).

3.1. Further assumptions on the observable. Besides having N global maxima ξ_1, \dots, ξ_N , we assume that the observable φ also satisfies

$$\varphi(x) = h_i(\text{dist}(x, \xi_i)), \quad \forall x \in B_{\varepsilon_i}(\xi_i), \quad i \in \{1, \dots, N\} \tag{19}$$

where $B_{\varepsilon_i}(\xi_i) \cap B_{\varepsilon_j}(\xi_j) = \emptyset$ and the function $h_i : [0, +\infty) \rightarrow \mathbb{R} \cup \{\infty\}$ is such that 0 is a global maximum ($h_i(0)$ may be $+\infty$), h_i is a strictly decreasing bijection $h_i : V \rightarrow W$ in a neighbourhood V of 0; and has one of the following three types of behaviour:

- (1) Type 1: there exists some strictly positive function $g : W \rightarrow \mathbb{R}$ such that for all $y \in \mathbb{R}$

$$\lim_{s \rightarrow h_i(0)} \frac{h_i^{-1}(s + yg(s))}{h_i^{-1}(s)} = e^{-y}; \tag{20}$$

- (2) Type 2: $h_i(0) = +\infty$ and there exists $\beta > 0$ such that for all $y > 0$

$$\lim_{s \rightarrow \infty} \frac{h_i^{-1}(sy)}{h_i^{-1}(s)} = y^{-\beta}; \tag{21}$$

- (3) Type 3: $h_i(0) = D < +\infty$ and there exists $\gamma > 0$ such that for all $y > 0$

$$\lim_{s \rightarrow 0} \frac{h_i^{-1}(D - sy)}{h_i^{-1}(D - s)} = y^\gamma. \tag{22}$$

Remark 3.1. Note that as long as the invariant measure has no atoms, then under the assumptions above on the observable we have that condition (R1) is easily satisfied.

3.2. Assumptions on the system and examples of application. We assume that the system admits a first return time induced map with decay of correlations against L^1 observables. In order to clarify what is meant by the latter we define:

Definition 3.2 (Decay of correlations). Let $\mathcal{C}_1, \mathcal{C}_2$ denote Banach spaces of real valued measurable functions defined on \mathcal{X} . We denote the *correlation* of non-zero functions $\phi \in \mathcal{C}_1$ and $\psi \in \mathcal{C}_2$ w.r.t. a measure \mathbb{P} as

$$\text{Cor}_{\mathbb{P}}(\phi, \psi, n) := \frac{1}{\|\phi\|_{\mathcal{C}_1} \|\psi\|_{\mathcal{C}_2}} \left| \int \phi(\psi \circ f^n) d\mathbb{P} - \int \phi d\mathbb{P} \int \psi d\mathbb{P} \right|.$$

We say that we have *decay of correlations*, w.r.t. the measure \mathbb{P} , for observables in \mathcal{C}_1 against observables in \mathcal{C}_2 if, for every $\phi \in \mathcal{C}_1$ and every $\psi \in \mathcal{C}_2$ we have

$$\text{Cor}_{\mathbb{P}}(\phi, \psi, n) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

We say that we have *decay of correlations against L^1 observables* whenever this holds for $\mathcal{C}_2 = L^1(\mathbb{P})$ and $\|\psi\|_{\mathcal{C}_2} = \|\psi\|_1 = \int |\psi| d\mathbb{P}$.

If a system already has decay of correlations against L^1 observables, then by taking the whole set \mathcal{X} as the base for the first return time induced map, which coincides with the original system, then the assumption we impose on the system is trivially satisfied. Examples of systems with such property include:

- Uniformly expanding maps on the circle/interval (see [BG97]);
- Markov maps (see [BG97]);
- Piecewise expanding maps of the interval with countably many branches like Rychlik maps (see [Ryc83]);
- Higher dimensional piecewise expanding maps studied by Saussol in [Sau00].

Remark 3.3. In the first three examples above the Banach space \mathcal{C}_1 for the decay of correlations can be taken as the space of functions of bounded variation. In the fourth example the Banach space \mathcal{C}_1 is the space of functions with finite quasi-Hölder norm studied in [Sau00]. We refer the readers to [BG97, Sau00] or [AFV14] for precise definitions but mention that

if $J \subset \mathbb{R}$ is an interval then $\mathbf{1}_J$ is of bounded variation and its BV-norm is equal to 2, *i.e.*, $\|\mathbf{1}_J\|_{BV} = 2$ and if A denotes a ball or an annulus then $\mathbf{1}_A$ has a finite quasi-Hölder norm.

Although the examples above are all in some sense uniformly hyperbolic, we can consider non-uniformly hyperbolic systems, such as intermittent maps, which admit a ‘nice’ first return time induced map over some subset $Y \subset \mathcal{X}$, called the base of the induced map. To be more precise, consider the usual original system as $f : \mathcal{X} \rightarrow \mathcal{X}$ with an ergodic f -invariant probability measure μ , choose a subset $Y \subset \mathcal{X}$ and consider $F_Y : Y \rightarrow Y$ to be the first return map f^{r_Y} to Y (note that F may be undefined at a zero Lebesgue measure set of points which do not return to Y , but most of these points are not important, so we will abuse notation here). Let $\mu_Y(\cdot) = \frac{\mu(\cdot \cap Y)}{\mu(Y)}$ be the conditional measure on Y . By Kac’s Theorem μ_Y is F_Y -invariant.

In [BSTV03], the authors show that for typical points $\zeta \in Y$ the HTS to balls around ζ is the same both for the original and induced maps. This means, in particular, that the existence of EVL for the induced map implies the existence of an EVL for the original system, at typical points, and the limit is actually the same. In [FFT13], a similar statement was obtained but this time for the convergence of the REPP at periodic points having the same compound Poisson limit. In [HWZ14] the authors show that regardless of the point $\zeta \in Y$ taken, the existence of a limit for the HTS (or EVL) for the induced first return time map implies the existence of a limit for the HTS (or EVL) of the original system, at ζ , and the limits coincide. This was generalised for the convergence of REPP in [FFTV15]. Namely, setting $v_n^Y = 1/\mu_Y(X > u_n)$, for the induced process X_i^Y ,

$$N_n^Y(J) := \mathcal{N}_{u_n}^Y(v_n^Y J) = \sum_{j \in v_n^Y J \cap \mathbb{N}_0} \mathbf{1}_{X_j^Y > u_n}.$$

Denote the speeded up return time r_A by $r_{A,Y}$ and the induced measure on Y by μ_Y .

Theorem 3.4 ([FFTV15, Theorem 3]). *For $\eta > 0$, setting $J_\eta := \cup_{s \in J} B_\eta(s)$, we assume that $N(J_\eta)$ is continuous in η , for all small η .*

$$N_n^Y \xrightarrow{\mu_Y} N \text{ as } n \rightarrow \infty \text{ implies } N_n \xrightarrow{\mu} N \text{ as } n \rightarrow \infty.$$

This means that if we have a system that admits a first return time induced map with decay of correlations against L^1 , then as long as the points $\xi_1, \dots, \xi_N \in Y$, the convergence of the REPP (which implies the existence of EVL or HTS) for the original system is determined by the convergence of the corresponding REPP for the induced map. Hence, we are reduced to proving the convergence of the REPP for systems with decay of correlations against L^1 observables. This fact motivates the following:

Assumption A *Let $f : \mathcal{X} \rightarrow \mathcal{X}$ be a systems with summable decay of correlations against L^1 observables, *i.e.*, for all $\varphi \in \mathcal{C}_1$ and $\psi \in L^1$, then $\text{Cor}(\varphi, \psi, n) \leq \rho_n$, with $\sum_{n \geq \mathbb{N}} \rho_n < \infty$. Moreover, we assume that there exists $C > 0$ such that for all $n \in \mathbb{N}$, we have $\mathbf{1}_{A_n^{(q)}}, \mathbf{1}_{A_{q,n}^{(\kappa)}} \in \mathcal{C}_1$ and $\|\mathbf{1}_{A_n^{(q)}}\|_{\mathcal{C}_1}, \|\mathbf{1}_{A_{q,n}^{(\kappa)}}\|_{\mathcal{C}_1} \leq C$.*

Among the examples of systems with these ‘nice’ induced maps we mention the *Manneville-Pomeau* (MP) map equipped with an absolutely continuous invariant probability measure.

The form for such maps given in [LSV99, BSTV03] is, for $\alpha \in (0, 1)$,

$$f = f_\alpha(x) = \begin{cases} x(1 + 2^\alpha x^\alpha) & \text{for } x \in [0, 1/2) \\ 2x - 1 & \text{for } x \in [1/2, 1] \end{cases}$$

Members of this family of maps are often referred to as Liverani-Saussol-Vaianti maps since their actual equation was first introduced in [LSV99]. Let \mathcal{P} be the *renewal partition*, that is the partition defined inductively by $Z \in \mathcal{P}$ if $Z = [1/2, 1)$ or $f(Z) \in \mathcal{P}$. Now let $Y \in \mathcal{P}$ and let F_Y be the first return map to Y and μ_Y be the conditional measure on Y . It is well-known that (Y, F_Y, μ_Y) is a Bernoulli map and hence, in particular, a Rychlik system (see [Ryc83] or [AFV14, Section 3.2.1] for the essential information about such systems).

3.3. Existence of distributional limits under Assumption A. As explained in [Fre13, Section 5.1], conditions $\mathbb{D}_q(u_n)$ and $\mathbb{D}_q(u_n)^*$ are designed to follow easily from decay of correlations. In fact, if we choose $\phi = \mathbf{1}_{\mathcal{A}_{q,n}}$ and $\psi = \mathbf{1}_{\mathcal{W}_{0,\ell}(\mathcal{A}_{q,n})}$, in the case of $\mathbb{D}_q(u_n)$, and $\phi = \mathbf{1}_{\mathcal{A}_{q,n}^{(\kappa)}}$ and $\psi = \mathbf{1}_{\cap_{j=2}^{\kappa} \mathcal{N}_{u_n}(I_j) = \kappa_j}$, in the case of $\mathbb{D}_q(u_n)^*$, we have that we can take $\gamma(n, t) = C\rho t$. Hence, by taking $(t_n)_n$ such that $\lim_{n \rightarrow \infty} n\rho t_n = 0$, we get that conditions $\mathbb{D}_q(u_n)$ and $\mathbb{D}_q(u_n)^*$ are trivially satisfied.

Now, we turn to condition $\mathbb{D}'_q(u_n)$. Taking $\phi = \psi = \mathbf{1}_{\mathcal{A}_{q,n}}$ and since $\|\mathbf{1}_{\mathcal{A}_{q,n}}\|_{\mathcal{C}_1} \leq C$ we easily get

$$\mu(U(u_n) \cap f^{-j}(U(u_n))) \leq (\mu(\mathcal{A}_{q,n}))^2 + \|\mathbf{1}_{\mathcal{A}_{q,n}}\|_{\mathcal{C}_1} \|\mathbf{1}_{\mathcal{A}_{q,n}}\|_{L^1(\mu)} \rho_j \leq (\mu(\mathcal{A}_{q,n}))^2 + C\mu(\mathcal{A}_{q,n})\rho_j. \quad (23)$$

Let $R_n := R(\mathcal{A}_{q,n})$. Using estimate (23) and $n\mu(U(u_n)) \rightarrow \tau$ as $n \rightarrow \infty$ it follows that there exists some constant $D > 0$ such that

$$\begin{aligned} n \sum_{j=q+1}^{\lfloor n/k_n \rfloor} \mu(\mathcal{A}_{q,n} \cap f^{-j}(\mathcal{A}_{q,n})) &= n \sum_{j=R_n}^{\lfloor n/k_n \rfloor} \mu(\mathcal{A}_{q,n} \cap f^{-j}(\mathcal{A}_{q,n})) \\ &\leq n \lfloor \frac{n}{k_n} \rfloor \mu(\mathcal{A}_{q,n})^2 + n C \mu(\mathcal{A}_{q,n}) \sum_{j=R_n}^{\lfloor n/k_n \rfloor} \rho_j \leq \frac{(n\mu(\mathcal{A}_{q,n}))^2}{k_n} + n C \mu(\mathcal{A}_{q,n}) \sum_{j=R_n}^{\infty} \rho_j \\ &\leq D \left(\frac{\tau^2}{k_n} + \tau \sum_{j=R_n}^{\infty} \rho_j \right) \xrightarrow{n \rightarrow \infty} 0, \end{aligned}$$

if we check that $\lim_{n \rightarrow \infty} R_n = +\infty$.

4. MAXIMA ALONG A NON-PERIODIC ORBIT

In this section we consider the case where ζ is a non-periodic point. Our goal is to see what are the effects on the REPP when we compare it with the process obtained by an observable which attains its maximum at a unique point. As we will see, despite ζ being non-periodic, the existence of multiple maxima along its orbit affects, for example, the extremal index and the multiplicity distribution.

Note that by (R1), we know that, for n sufficiently large, $U(u_n)$ is the union of N disjoint balls around each ξ_i that we denote by $U_n(\xi_i) = U_n^{(0)}(\xi_i)$, for $i = 1, \dots, N$, so that

$$U(u_n) = \bigcup_{i=1}^N U_n(\xi_i). \quad (24)$$

We begin by choosing a value of q for which condition $\mathcal{D}'(u_n)$ can be checked. From the computations in Section 3.3 we need to verify that $R(\mathcal{A}_{q,n}) \rightarrow \infty$, as $n \rightarrow \infty$ for such choice of q .

Lemma 4.1. *Assume that f satisfies Assumption A and that ζ is a non-periodic point and that f is continuous in every point of the orbit of ζ , namely $\zeta, f(\zeta), f^2(\zeta), \dots$. Let ξ_1, \dots, ξ_N be as in (18). Let $q = m_N - m_1$, then $\lim_{n \rightarrow \infty} R(\mathcal{A}_{q,n}) = \infty$.*

Proof. Let $q = m_N - m_1$. By definition of $\mathcal{A}_{q,n}$ we have that $f^j(\mathcal{A}_{q,n}) \cap \mathcal{A}_{q,n} \subset f^j(\mathcal{A}_{q,n}) \cap U(u_n) = \emptyset$, for all $j = 1, \dots, q$. Hence, we only need to check that the same holds true for all $j = q + 1, \dots, q + L$, where L is a given arbitrarily large integer. For that we will use the continuity of f over the orbit of ζ . Let

$$\epsilon := \min_{1 \leq i \leq L, 1 \leq j \leq N} \text{dist}(f^i(\xi_N), \xi_j).$$

Using the continuity over the orbit of ζ and in particular of ξ_N , for every $1 \leq i \leq L$, consider the balls $B_{\delta_i}(\xi_N)$ such that $f^i(B_{\delta_i}(\xi_N)) \subset B_{\epsilon/2}(f^i(\xi_N))$ and consider the open set $D_N = \bigcap_{i=1}^L B_{\delta_i}(\xi_N)$. Now, for each $j = 1, \dots, N-1$ let D_j be a ball around ξ_j so that $f^{m_N - m_j}(D_j) \subset D_N$.

Finally, recalling (R1) take n sufficiently large so that each $U_n(\xi_i) \subset D_i \cap B_{\epsilon/2}(\xi_i)$ so that $U(u_n) \subset \bigcup_{i=1}^N D_i \cap B_{\epsilon/2}(\xi_i)$. Recall that $\mathcal{A}_{q,n} \subset U(u_n)$ and then by construction for every $j = 1, \dots, L$ we have that $f^{q+j}(\mathcal{A}_{q,n}) \subset \bigcup_{i=1}^N B_{\epsilon/2}(f^{q+j}(\xi_i))$ and since by definition of ϵ , we have $B_{\epsilon/2}(f^{q+j}(\xi_i)) \cap U(u_n) = \emptyset$, then the result follows. \square

4.1. The clustering effect and estimates for the extremal index. When the point ζ is an absolute maximum and ζ is non-periodic we know that the extremal index θ is equal to 1. The situation is completely different when the maximum occurs at more than one point along the orbit of ζ .

Proposition 4.2. *Assume that f satisfies Assumption A. Let X_0, X_1, \dots be given by (1), where the observable φ has N maximal points ξ_1, \dots, ξ_N related by (18) and satisfies (19), with $\zeta \in \mathcal{X}$ a non-periodic point. In this case the extremal index θ is given by*

$$\theta = \lim_{n \rightarrow \infty} \frac{\sum_{i=1}^N \mu \left(U_n^{(0)}(\xi_i) - \bigcup_{j=i+1}^N f^{-(m_j - m_i)}(U_n^{(0)}(\xi_j)) \right)}{\sum_{i=1}^N \mu \left(U_n^{(0)}(\xi_i) \right)},$$

whenever the limit exists.

Proof. In this case we have that $q = m_N - m_1$ and $\mathcal{A}_{q,n}^{(0)} = \bigcup_{i=1}^N \mathcal{A}_{q,n}^{(0)}(\xi_i)$, where

$$\mathcal{A}_{q,n}^{(0)}(\xi_i) = U_n^{(0)}(\xi_i) - \bigcup_{j=i+1}^N f^{-(m_j - m_i)}(U_n^{(0)}(\xi_j)).$$

By the definition of $\theta = \lim_{n \rightarrow \infty} \frac{\mu(\mathcal{A}_{q,n}^{(0)})}{\mu(U(u_n)^{(0)})}$, the conclusion follows. \square

In order to illustrate the richness of scenarios that the formula for the EI of the previous proposition encompasses we consider below some examples. By condition (R1), we know that, for n sufficiently large, $U(u_n) = U(u_n)^{(0)}$ is the union of N disjoint balls around each ξ_i that we denote by $U_n^{(0)}(\xi_i)$, for $i = 1, \dots, N$, so that $U(u_n)^{(0)} = \cup_{i=1}^N U_n^{(0)}(\xi_i)$. Depending on the different shapes of h_i in (19), the balls $U_n^{(0)}(\xi_i)$ may have different sizes and shapes. Moreover, if f is continuous along the orbit of ζ , then for $1 \leq i < j \leq N$ we have $f^{m_j - m_i}(U_n^{(0)}(\xi_j)) \cap U_n^{(0)}(\xi_i) \neq \emptyset$ but, depending on the expansion rate and geometric distortion of $f^{m_j - m_i}$, it is not a priori clear if the one of the sets contains the other or not. Hence, in order to simplify, we assume that for $1 \leq i < j \leq N$

$$(R3) \quad U_n^{(0)}(\xi_i) \subset f^{-(m_j - m_i)}(U_n^{(0)}(\xi_j)) \quad \text{or} \quad f^{-(m_j - m_i)}(U_n^{(0)}(\xi_j)) \cap \partial U_n^{(0)}(\xi_i) = \emptyset,$$

where the notation ∂B is used to denote the boundary of the set B .

Remark 4.3. Note that for all $1 \leq i < j \leq N$, we have $f^{-(m_j - m_i)}(U_n^{(0)}(\xi_j)) \cap U_n^{(0)}(\xi_i) \neq \emptyset$. Hence, $f^{-(m_j - m_i)}(U_n^{(0)}(\xi_j)) \cap \partial U_n^{(0)}(\xi_i) = \emptyset$ means that the connected component of $f^{-(m_j - m_i)}(U_n^{(0)}(\xi_j))$ that intersects $U_n^{(0)}(\xi_i)$ is completely contained in $U_n^{(0)}(\xi_i)$.

In the next result we treat the case where the condition (R3) is valid. Before we assert the corollary we introduce some notation.

Let I_1 be the set of indices $i \in \{1, \dots, N - 1\}$ such that

$$\bigcup_{\ell=i+1}^N f^{-(m_\ell - m_i)}(U_n^{(0)}(\xi_\ell)) \cap \partial U_n^{(0)}(\xi_i) = \emptyset. \quad (25)$$

Under (R3), for every $i \in I_1$, there exists $j_i \in \{i + 1, \dots, N\}$ such that

$$\bigcup_{\ell=i+1}^N f^{-(m_\ell - m_i)}(U_n^{(0)}(\xi_\ell)) \cap U_n^{(0)}(\xi_i) = f^{-(m_{j_i} - m_i)}(U_n^{(0)}(\xi_{j_i})) \cap U_n^{(0)}(\xi_i).$$

Corollary 4.4. *If we are under the conditions of Proposition 4.2 and the condition (R3) is satisfied, then the extremal index θ is given by*

$$\theta = \lim_{n \rightarrow \infty} \frac{\sum_{i \in I_1} \left[\mu(U_n^{(0)}(\xi_i)) - \mu(f^{-(m_{j_i} - m_i)}(U_n^{(0)}(\xi_{j_i})) \cap U_n^{(0)}(\xi_i)) \right] + \mu(U_n^{(0)}(\xi_N))}{\sum_{i=1}^N \mu(U_n^{(0)}(\xi_i))},$$

whenever the limit exists.

Let us now assume we are in the particular case where μ is absolutely continuous with respect to the Lebesgue measure and its Radon-Nikodym density is sufficiently regular so that for all $x \in \mathcal{X}$ we have

$$\lim_{\varepsilon \rightarrow 0} \frac{\mu(B_\varepsilon(x))}{\text{Leb}(B_\varepsilon(x))} = \frac{d\mu}{d\text{Leb}}(x). \quad (26)$$

Note that if f is one dimensional smooth map and $\log(Df)$ is Hölder continuous then, as seen in [FFT15, Section 7.2], formula (26) holds.

Corollary 4.5. *If we are under the hypothesis of Corollary 4.4 and μ is absolutely continuous and (26) holds then the extremal index θ is given by*

$$\theta = \lim_{n \rightarrow \infty} \frac{\sum_{i \in I_1} \left[\mu \left(U_n^{(0)}(\xi_i) \right) - \frac{1}{|\det Df^{m_{j_i} - m_i}(\xi_i)|} \mu \left(U_n^{(0)}(\xi_{j_i}) \right) \right] + \mu \left(U_n^{(0)}(\xi_N) \right)}{\sum_{i=1}^N \mu \left(U_n^{(0)}(\xi_i) \right)},$$

whenever the limit exists.

Proof. We only need to notice that, by the mean value theorem and (26)

$$\mu \left(f^{-(m_{j_i} - m_i)}(U_n^{(0)}(\xi_{j_i})) \cap U_n^{(0)}(\xi_i) \right) \sim \frac{1}{|\det Df^{m_{j_i} - m_i}(\xi_i)|} \mu \left(U_n^{(0)}(\xi_{j_i}) \right).$$

□

4.2. Estimating the multiplicity distribution. By Proposition 2.10 of [FFR] we have that the multiplicity distribution of the limiting compound Poisson process for the REPP is given by the formula

$$\pi(k) = \lim_{n \rightarrow \infty} \frac{\mu \left(\mathcal{A}_{q,n}^{(k-1)} \right) - \mu \left(\mathcal{A}_{q,n}^{(k)} \right)}{\mu \left(\mathcal{A}_{q,n}^{(0)} \right)}, \quad (27)$$

whenever the limit exists.

Assume that (R3) holds.

Let I_1 be defined as in (25). For every $i \in \{1, \dots, N\}$, we define $j_{i,0} = i$. Then, for $i \in \{1, \dots, N - k\}$ we inductively define I_k in the following way.

Assuming that I_k is defined as well as $j_{1,k-1}, \dots, j_{N-k+2,k-1}$, then for every $i = 1, \dots, N - k + 1$, we define $j_{i,k}$ to the index such that

$$\bigcup_{\ell=i+1}^N f^{-(m_{j_{\ell,k-1}} - m_i)}(U_n^{(0)}(\xi_{j_{\ell,k-1}})) \cap U_n^{(0)}(\xi_i) = f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i), \quad (28)$$

when $i \in I_k$ and $j_{i,k} = i$, otherwise.

Now, we define I_{k+1} as the set of indices $i \in \{1, \dots, N - k\}$ such that

$$\bigcup_{\ell=i+1}^N f^{-(m_{j_{\ell,k}} - m_i)}(U_n^{(0)}(\xi_{j_{\ell,k}})) \cap \partial U_n^{(0)}(\xi_i) = \emptyset.$$

Proposition 4.6. *Assume that f satisfies Assumption A. Let X_0, X_1, \dots be given by (1), where the observable φ has N maximal points ξ_1, \dots, ξ_N related by (18) and satisfies (19), with $\zeta \in \mathcal{X}$ a non-periodic point. Assume that condition (R3) holds. The multiplicity distribution of the point process is given by equation (27) (whenever the limit exists), where, for $k < N$,*

$$\begin{aligned} \mu(\mathcal{A}_{q,n}^{(k)}) &= \mu \left(f^{-(m_{j_{N-k,k}} - m_{N-k})}(U_n^{(0)}(\xi_{j_{N-k,k}})) \cap U_n^{(0)}(\xi_{N-k}) \right) \\ &+ \sum_{i \in I_{k+1}} \left[\mu \left(f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i) \right) - \mu \left(f^{-(m_{j_{i,k+1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}})) \cap U_n^{(0)}(\xi_i) \right) \right] \end{aligned}$$

and, for $k \geq N$, $\mu(\mathcal{A}_{q,n}^{(k)}) = 0$.

Proof. We will prove by induction that, for $i \in I_{k+1}$,

$$\mathcal{A}_{q,n}^{(k)}(\xi_i) = U_n^{(0)}(\xi_i) \cap f^{-(m_{j_i,k}-m_i)}(U_n^{(0)}(\xi_{j_i,k})) - f^{-(m_{j_i,k+1}-m_i)}(U_n^{(0)}(\xi_{j_i,k+1}))$$

and

$$U_n^{(k+1)}(\xi_i) = f^{-(m_{j_i,k+1}-m_i)}(U_n^{(0)}(\xi_{j_i,k+1})) \cap U_n^{(0)}(\xi_i),$$

and, for $i \notin I_{k+1}$, $\mu\left(\mathcal{A}_{q,n}^{(k)}(\xi_i)\right) = 0$ since $U_n^{(k)}(\xi_i) = U_n^{(k-1)}(\xi_i)$.

For $i \in I_1$ since $j_{\ell,0} = \ell$, we get

$$\begin{aligned} \mathcal{A}_{q,n}^{(0)}(\xi_i) &= U_n^{(0)}(\xi_i) - \bigcup_{\ell=i+1}^N f^{-(m_\ell-m_i)}(U_n^{(0)}(\xi_\ell)) \\ &= U_n^{(0)}(\xi_i) - \bigcup_{\ell=i+1}^N f^{-(m_{j_{\ell,0}}-m_i)}(U_n^{(0)}(\xi_{j_{\ell,0}})) \\ &= U_n^{(0)}(\xi_i) \cap f^{-(m_{j_{i,0}}-m_i)}(U_n^{(0)}(\xi_{j_{i,0}})) - f^{-(m_{j_{i,1}}-m_i)}(U_n^{(0)}(\xi_{j_{i,1}})), \end{aligned}$$

by definition of $j_{i,1}$, and, moreover, we have $U_n^{(1)}(\xi_i) = f^{-(m_{j_{i,1}}-m_i)}(U_n^{(0)}(\xi_{j_{i,1}})) \cap U_n^{(0)}(\xi_i)$.

Note that, for $i = N$, $\bigcup_{\ell=i+1}^N f^{-(m_\ell-m_i)}(U_n^{(0)}(\xi_\ell)) = \emptyset$ and so

$$\mathcal{A}_{q,n}^{(0)}(\xi_N) = U_n^{(0)}(\xi_N) = f^{-(m_{j_{N,0}}-m_i)}(U_n^{(0)}(\xi_{j_{N,0}})).$$

For $i \notin I_1$, $i \neq N$, $U_n^{(0)}(\xi_i) - \bigcup_{\ell=i+1}^{N+i} f^{-(m_\ell-m_i)}(U_n^{(0)}(\xi_\ell)) = \emptyset$, so $\mu\left(\mathcal{A}_{q,n}^{(0)}(\xi_i)\right) = 0$ and $\mu\left(U_n^{(1)}(\xi_i)\right) = \mu\left(U_n^{(0)}(\xi_i)\right)$.

By induction, for $i \in I_{k+1}$, we obtain

$$\begin{aligned} \mathcal{A}_{q,n}^{(k)}(\xi_i) &= U_n^{(k)}(\xi_i) - \bigcup_{\ell=i+1}^N f^{-(m_\ell-m_i)}(U_n^{(k)}(\xi_\ell)) \\ &= U_n^{(k)}(\xi_i) - \bigcup_{\ell=i+1}^N f^{-(m_\ell-m_i)}\left(f^{-(m_{j_{\ell,k}}-m_i)}(U_n^{(0)}(\xi_{j_{\ell,k}})) \cap U_n^{(0)}(\xi_i)\right), \text{ by induction} \\ &= f^{-(m_{j_{i,k}}-m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i) - \bigcup_{\ell=i+1}^N f^{-(m_{j_{\ell,k}}-m_i)}(U_n^{(0)}(\xi_{j_{\ell,k}})), \text{ by induction} \\ &= f^{-(m_{j_{i,k}}-m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i) - f^{-(m_{j_{i,k+1}}-m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}})), \text{ by definition of } j_{i,k+1}. \end{aligned}$$

and in particular we also have $U_n^{(k+1)}(\xi_i) = f^{-(m_{j_{i,k+1}}-m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}})) \cap U_n^{(0)}(\xi_i)$.

For $i = N - k + 1$, $\bigcup_{\ell=i+1}^N f^{-(m_\ell-m_i)}(U_n^{(k)}(\xi_\ell)) = \emptyset$ and so

$$\mathcal{A}_{q,n}^{(k)}(\xi_{N-k}) = f^{-(m_{j_{N-k+1,k}}-m_{N-k})}(U_n^{(0)}(\xi_{j_{N-k+1,k}})) \cap U_n^{(0)}(\xi_{N-k}).$$

For $i \notin I_{k+1}$, $i \neq N - k + 1$ $\mu\left(\mathcal{A}_{q,n}^{(k)}(\xi_i)\right) = 0$ and

$$U_n^{(k+1)}(\xi_i) = f^{-(m_{j_{i,k}}-m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i).$$

For $k > N + 1$, $\mu\left(\mathcal{A}_{q,n}^{(k)}(\xi_i)\right) = 0$.

This concludes the proof. \square

Corollary 4.7. *Under the assumptions of Proposition 4.6, assume that μ is absolutely continuous with respect to the Lebesgue measure and (26) holds. Then the multiplicity distribution of the point process is given by formula (27), where, for $k < N$,*

$$\mu(\mathcal{A}_{q,n}^{(k)}) = \sum_{i \in I_{k+1}} \left(\alpha_n(m_{j_{i,k}}, m_i) - \alpha_n(m_{j_{i,k+1}}, m_i) \right) + \alpha_n(m_{j_{N-k}}, m_{N-k})$$

and

$$\alpha_n(a, b) = \frac{1}{|\det Df^{a-b}(f^b(\zeta))|} \mu\left(U_n^{(0)}(f^a(\zeta))\right).$$

Proof. Since μ is absolutely continuous with respect to the Lebesgue measure and (26) holds, then

$$\mu\left(f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i)\right) = \frac{1}{|\det Df^{m_{j_{i,k}} - m_i}(\xi_i)|} \mu\left(U_n^{(0)}(\xi_{j_{i,k}})\right) = \alpha_n(m_{j_{i,k}}, m_i).$$

Replacing this in the previous proposition we obtain the conclusion \square

Corollary 4.8. *Under the assumptions of Proposition 4.6, assume further that μ is absolutely continuous with respect to the Lebesgue measure and (26) holds and*

$$f^{-(m_{i+1} - m_i)}\left(U_n^{(0)}(\xi_{i+1})\right) \subset U_n^{(0)}(\xi_i).$$

Then the multiplicity distribution of the point process is given by formula (27) where, for $k < N$,

$$\mu(\mathcal{A}_{q,n}^{(k)}) = \sum_{i \in I_{k+1}} \left(\alpha_n(m_{i+k}, m_i) - \alpha_n(m_{i+k+1}, m_i) \right) + \alpha_n(m_N, m_{N-k})$$

and $\alpha_n(a, b)$ is as in Corollary 4.7.

Proof. Using the notation of Proposition 4.6, note that in this case $j_{i,k} = i + k$. \square

4.3. An example. Let $(f, \mathbb{S}^1, Leb, \mathcal{B})$ be the dynamical system where $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ is the map given by $f(x) = 3x \pmod{1}$. Take $\zeta = \frac{\sqrt{2}}{16}$ and an observable $\varphi : \mathbb{S}^1 \rightarrow \mathbb{R} \cup \{\infty\}$ such that

$$\varphi(x) = \begin{cases} -\log|x - \zeta|, & \text{if } x \in B_\epsilon(\zeta), \\ |x - f(\zeta)|^{-\frac{1}{2}}, & \text{if } x \in B_\epsilon(f(\zeta)) \\ |x - f^3(\zeta)|^{-\frac{1}{2}}, & \text{if } x \in B_\epsilon(f^3(\zeta)) \\ 0, & \text{otherwise.} \end{cases}$$

where we choose $\epsilon > 0$, such that $B_\epsilon(f^i(\zeta)) \cap B_\epsilon(f^j(\zeta)) = \emptyset$ for $i, j \in \{0, 1, 3\}$ and $i \neq j$. In this case we have that $u_F = \sup_{x \in \mathbb{S}^1} \varphi(x) = \infty$ and it occurs at $\zeta, f(\zeta)$ and $f^3(\zeta)$.

Given $\tau > 0$ and a sequence $(u_n)_{n \in \mathbb{N}}$ satisfying (3) we have that

$$U_n = (\zeta - e^{-u_n}, \zeta + e^{-u_n}) \cup (f(\zeta) - u_n^{-2}, f(\zeta) + u_n^{-2}) \cup (f^3(\zeta) - u_n^{-2}, f^3(\zeta) + u_n^{-2}).$$

As before we let $U_n(\zeta) = (\zeta - e^{-u_n}, \zeta + e^{-u_n})$, $U_n(f(\zeta)) = (f(\zeta) - u_n^{-2}, f(\zeta) + u_n^{-2})$ and $U_n(f^3(\zeta)) = (f^3(\zeta) - u_n^{-2}, f^3(\zeta) + u_n^{-2})$. We notice that for n sufficiently large we have that

$$(\zeta - e^{-u_n}, \zeta + e^{-u_n}) \not\subseteq f^{-1}\left((f(\zeta) - u_n^{-2}, f(\zeta) + u_n^{-2})\right),$$

and

$$f^{-2}((f^3(\zeta) - u_n^{-2}, f^3(\zeta) + u_n^{-2})) \cap U_n(f(\zeta)) \not\subseteq (f(\zeta) - u_n^{-2}, f(\zeta) + u_n^{-2}).$$

The extremal index: Noting that $m_1 = 0, m_2 = 1, m_3 = 3$, then $I_1 = \{2\}$ and $j_2 = 3$. So, by Corollary 4.7 we have that

$$\begin{aligned} \theta &= \lim_{n \rightarrow \infty} \frac{\mu(U_n^{(0)}(f^{m_2}(\zeta))) - \mu(f^{-(m_3-m_2)}(U_n^{(0)}(f^{m_3}(\zeta)))) + \mu(U_n^{(0)}(f^{m_3}(\zeta)))}{\sum_{i=1}^3 \mu(U_n^{(0)}(\xi_i))} \\ &= \lim_{n \rightarrow \infty} \frac{u_n^{-2} - \frac{1}{Df^2(f^3(\zeta))} u_n^{-2} + u_n^{-2}}{e^{-u_n} + u_n^{-2} + u_n^{-2}} = \frac{2 - (\frac{1}{3})^2}{2} = \frac{17}{18}. \end{aligned}$$

The multiplicity distribution: To compute the multiplicity distribution we need to determine the sets I_k , for all $k \in \mathbb{N}$. It is easy to see that, in this case, $I_1 = \{2\}$ and $I_k = \emptyset$, for all $k \geq 2$. We now need to determine the value of $j_{i,k}$, for all $k \in \mathbb{N}$ and all $i \in I_k$. By definition $j_{i,0} = i$, for all $i \in \mathbb{N}$. In addition, we can see that $j_{2,1} = 3$. Since $m_1 = 0, m_2 = 1, m_3 = 3$, by Corollary 4.7, we have

$$\begin{aligned} \mu(\mathcal{A}_{q,n}^{(0)}) &= (\alpha(m_{j_{2,0},2}) - \alpha(m_{j_{2,1},2}) + \alpha_n(m_{j_{3,0},m_3}))2u_n^{-2} \\ &= (\alpha(m_2, m_2) - \alpha(m_3, m_2) + \alpha_n(m_3, m_3))2u_n^{-2} \\ &= 2 \left(u_n^{-2} - \left(\frac{1}{3}\right)^2 u_n^{-2} + u_n^{-2} \right) = \frac{34}{9} u_n^{-2}, \\ \mu(\mathcal{A}_{q,n}^{(1)}) &= \alpha(m_{j_{2,1},m_2}) = \alpha(m_3, m_2) = 2 \left(\frac{1}{3}\right)^2 u_n^{-2} = \frac{2}{9} u_n^{-2}, \\ \mu(\mathcal{A}_{q,n}^{(2)}) &= \alpha(m_{j_{1,2},m_1}) = \alpha(m_1, m_1) = 2e^{-u_n}, \\ \mu(\mathcal{A}_{q,n}^{(k)}) &= 0 \text{ for } k \geq 3. \end{aligned}$$

Therefore,

$$\begin{aligned} \pi(1) &= \lim_{n \rightarrow \infty} \frac{\mu(\mathcal{A}_{q,n}^{(0)}) - \mu(\mathcal{A}_{q,n}^{(1)})}{\mu(\mathcal{A}_{q,n}^{(0)})} = \lim_{n \rightarrow \infty} \frac{\frac{34}{9}u_n^{-2} - \frac{2}{9}u_n^{-2}}{\frac{34}{9}u_n^{-2}} = \frac{32}{34}, \\ \pi(2) &= \lim_{n \rightarrow \infty} \frac{\mu(\mathcal{A}_{q,n}^{(1)}) - \mu(\mathcal{A}_{q,n}^{(2)})}{\mu(\mathcal{A}_{q,n}^{(0)})} = \lim_{n \rightarrow \infty} \frac{\frac{2}{9}u_n^{-2} - e^{-u_n}}{\frac{34}{9}u_n^{-2}} = \frac{2}{34}, \\ \pi(3) &= \lim_{n \rightarrow \infty} \frac{\mu(\mathcal{A}_{q,n}^{(2)}) - \mu(\mathcal{A}_{q,n}^{(3)})}{\mu(\mathcal{A}_{q,n}^{(0)})} = \lim_{n \rightarrow \infty} \frac{e^{-u_n}}{\frac{34}{9}u_n^{-2}} = 0, \\ \pi(k) &= 0 \text{ for } k \geq 4. \end{aligned}$$

For this particular example, we performed a simulation study and computed a finite sample multiplicity distribution. In this simulation we chose at random an initial condition $x_0 = 0.887806951194545$ that was iterated 500 000 times by the system. The threshold taken was $u = 20.00000412229153$ which corresponds to the 0.99-quantile of the distribution of X_0 . The

results are presented in Figure 1 and reveal a good agreement with the theoretical estimates provided by π computed above.

4.4. Non correlated maxima along non periodic orbits. Although the main purpose of the paper is to study the effect of multiple maxima when they are bound by belonging to the same orbit, we note that if we suppose that φ achieves a global maximum value at the points ξ_1, \dots, ξ_n but these maximal points have no intersecting orbits then one can also obtain a limiting EVL, which in the case of the maximal points being non-periodic, has no clustering and hence the EI equals 1.

Let $\mathcal{O}(x)$ denote the sequence of points that form the orbit of the point x , *i.e.*, $\mathcal{O}(x) = \{x, f(x), f^2(x), \dots\}$.

Proposition 4.9. *Assume that f satisfies Assumption A. Let X_0, X_1, \dots be given by (1), where the observable φ has N maximal points ξ_1, \dots, ξ_N . Assume that all the maximal points are non-periodic, that f is continuous on $\cup_{i=1}^N \mathcal{O}(\xi_i)$ and that*

$$\bigcap_{i=1}^N \mathcal{O}(\xi_i) = \emptyset. \quad (29)$$

Then X_0, X_1, \dots satisfies conditions $\mathbb{D}_0(u_n)$ and $\mathbb{D}'_0(u_n)$ and consequently for $(u_n)_{n \in \mathbb{N}}$ satisfying (3) then $\lim_{n \rightarrow \infty} \mu(M_n \leq u_n) = e^{-\tau}$.

Proof. In this case we take $q = 0$, meaning that $\mathcal{A}_{q,n} = U(u_n)$. Conditions $\mathbb{D}_0(u_n)$ and $\mathbb{D}'_0(u_n)$ follow from the computations in Section 3.3 as long as one check that $\lim_{n \rightarrow \infty} R(U(u_n)) = +\infty$. This last statement follows from a continuity argument very similar to the proof of Lemma 4.1. \square

Remark 4.10. This is in agreement with the result obtained in [HNT12] for multiple uncorrelated maxima chosen as independent typical points of the invariant measure of non-uniformly hyperbolic systems admitting a Young tower with summable tails. Here, if the system is continuous and satisfies Assumption A (which is strictly contained in the class of systems considered in [HNT12]), the statement can be reinforced since it applies to all non-periodic points chosen for maxima as long as they have non-intersecting orbits.

5. MULTIPLE MAXIMA LYING ON PERIODIC ORBIT

In this section we consider the case where ζ is a repelling periodic point of prime period p , meaning that there exists $p \in \mathbb{N}$ such that $f^p(\zeta) = \zeta$, being that p is the smallest integer for which that happens, and the derivative $Df^p(\zeta)$ is defined and all its eigenvalues are strictly larger than 1. Note that in this case, if for example μ is absolutely continuous with respect to Lebesgue measure and (26) holds then condition (R_2) is satisfied. From [FFT12] we know that if the global maximum of φ is attained only at ζ then we expect the existence of an EI equals to

$$\theta = 1 - \frac{1}{|\det Df^p(\zeta)|},$$

in the case μ is absolutely continuous with respect to Lebesgue and (26) holds. Moreover, by [FFT13], we know that the REPP converges to a compound Poisson process with a geometric multiplicity distribution given by

$$\pi(\kappa) = \theta(1 - \theta)^{\kappa-1},$$

for all $\kappa \geq 1$.

In this section, our goal is to study the effect of having multiple maxima along the orbit of the periodic point ζ . We will see the EI is affected (it decreases) and asymptotics of the REPP is also affected.

We assume without loss of generality that φ achieves a maximum at the points ξ_1, \dots, ξ_N , where each ξ_i is given by equation (18), where $0 = m_1 < m_2 < \dots < m_N \leq p - 1$, so that, in particular $\xi_1 = \zeta$.

We begin by choosing a value of q for which condition $\mathbb{D}'_q(u_n)$ can be checked. One easily anticipates that a suitable choice is taking $q = p$. From the computations in Section 3.3 we need to verify that $R(\mathcal{A}_{q,n}) \rightarrow \infty$, as $n \rightarrow \infty$ for such choice of q .

Lemma 5.1. *Assume that ζ is a repelling periodic point of prime period p . Let ξ_1, \dots, ξ_N be as in (18), where $0 = m_1 < m_2 < \dots < m_N \leq p - 1$. Let $q = p$, then $\lim_{n \rightarrow \infty} R(\mathcal{A}_{q,n}) = \infty$.*

Proof. Let $g := f^p$, and so every ξ_i is a fixed point of g . As ζ is a repelling periodic point, we have that there is a neighbourhood of ξ_i that we denote by $V(\xi_i)$ where $g|_{V(\xi_i)} : V(\xi_i) \rightarrow g(V(\xi_i)) \subset \mathcal{X}$ is a diffeomorphism. By the Hartman-Grobman theorem there is a neighbourhood $W(0)$ of $0 \in T_\zeta \mathcal{X}$ and a homeomorphism $h : W(0) \rightarrow V(\zeta)$ such that $h \circ Dg(\zeta) = g \circ h$.

As $W(0)$ is an open set, there is $r > 0$ so that the ball of radius r , denoted by $B_r(0)$ is contained in $W(0)$. Let $x \in B_r(\xi_i)$ and let $\epsilon = \text{dist}(x, \xi_i)$. By hypothesis, the point ξ_i is repelling periodic and it implies that there exists $\lambda > 1$ such that the diameter of $(Dg(\zeta))^\kappa(B_\epsilon(\xi_i))$ is at least $\lambda^\kappa \epsilon$. So, in order for $\lambda^\kappa \epsilon > r$, we must have that $\kappa > \frac{\log r - \log \epsilon}{\log \lambda}$.

Hence, $(Dg(\zeta))^\kappa(B_\epsilon(\xi_i)) \subset B_r(\xi_i)$, for $\kappa \leq \frac{\log r - \log \epsilon}{\log \lambda}$.

Now, for each $j \in \{1, \dots, N\}$, let $V(\xi_j)$ be an open neighbourhood of ξ_j where $f|_{V(\xi_j)} : V(\xi_j) \rightarrow V(f^{m_{j+1}}(\zeta))$ is a diffeomorphism and the Hartman-Grobman theorem holds. Let $n \in \mathbb{N}$ such that $U_n(\xi_j) \subset V(\xi_j)$. Given $x \in \mathcal{A}_{q,n}$, by the definition of $\mathcal{A}_{q,n}$ we know that $g^\kappa(x) \notin \mathcal{A}_{q,n}$ while $g^\kappa(x) \in B_r(\xi_i)$. Let ϵ_n be the diameter of $\mathcal{A}_{q,n}$. Then while $\kappa \leq \frac{\log r - \log \epsilon_n}{\log \lambda}$ we have that $g^\kappa(x) \notin \mathcal{A}_{q,n}$. Since $\frac{\log r - \log \epsilon_n}{\log \lambda} \rightarrow \infty$, as $n \rightarrow \infty$, then it follows that $\lim_{n \rightarrow \infty} R(\mathcal{A}_{q,n}) = \infty$. \square

5.1. The extremal index. In what follows we need some notation, which we introduce now: We define $m_{i+N} := m_i + p$, so that $f^{m_{i+N}}(\zeta) = f^{m_i}(\zeta)$.

Proposition 5.2. *Let X_0, X_1, \dots be given by (1), where the observable φ is given by (19), with $\zeta \in \mathcal{X}$ a repelling periodic point. In this case the extremal index θ is given by*

$$\theta = \lim_{n \rightarrow \infty} \frac{\sum_{i=1}^N \mu \left(U_n^{(0)}(\xi_i) - \bigcup_{j=i+1}^{N+i} f^{-(m_j - m_i)}(U_n^{(0)}(\xi_j)) \right)}{\sum_{i=1}^N \mu \left(U_n^{(0)}(\xi_i) \right)},$$

whenever the limit exists.

Proof. Note that $\mathcal{A}_{p,n}^{(0)} = \bigcup_{i=1}^N \mathcal{A}_{p,n}^{(0)}(\xi_i)$, where

$$\mathcal{A}_{p,n}^{(0)}(\xi_i) = U_n^{(0)}(\xi_i) - \bigcup_{j=i+1}^{N+i} f^{-(m_j - m_i)}(U_n^{(0)}(\xi_j)).$$

Since $\theta = \lim_{n \rightarrow \infty} \frac{\mu(\mathcal{A}_{p,n}^{(0)})}{\mu(U(u_n)^{(0)})}$, the conclusion follows. \square

Just like in the non-periodic case, we now consider some particular examples that illustrate the richness of scenarios that the formula for the EI of the previous proposition encompasses. Hence, as before, we assume that condition (R3) holds.

First, we introduce some notation.

Let I_1 be the set of indices $i \in \{1, \dots, N\}$ such that

$$\bigcup_{\ell=i+1}^{N+i} f^{-(m_\ell - m_i)}(U_n^{(0)}(\xi_\ell)) \cap \partial U_n^{(0)}(\xi_i) = \emptyset. \quad (30)$$

Under (R3), for every $i \in I_1$, there exists $j_i \in \{i+1, \dots, N+i\}$ such that

$$\bigcup_{\ell=i+1}^{N+i} f^{-(m_\ell - m_i)}(U_n^{(0)}(\xi_\ell)) \cap U_n^{(0)}(\xi_i) = f^{-(m_{j_i} - m_i)}(U_n^{(0)}(\xi_{j_i})) \cap U_n^{(0)}(\xi_i).$$

Corollary 5.3. *If we are under the conditions of Proposition 5.2 and the condition (R3) is satisfied, then the extremal index θ is given by*

$$\theta = \lim_{n \rightarrow \infty} \frac{\sum_{i \in I_1} \left[\mu(U_n^{(0)}(\xi_i)) - \mu(f^{-(m_{j_i} - m_i)}(U_n^{(0)}(\xi_{j_i})) \cap U_n^{(0)}(\xi_i)) \right]}{\sum_{i=1}^N \mu(U_n^{(0)}(\xi_i))},$$

whenever the limit exists.

Let us now assume we are in the particular case where μ is absolutely continuous with respect to the Lebesgue measure and (26) holds.

Corollary 5.4. *If we are under the hypotheses of Corollary 5.3 and μ is absolutely continuous and (26) holds then the extremal index θ is given by*

$$\theta = \lim_{n \rightarrow \infty} \frac{\sum_{i \in I_1} \left[\mu(U_n^{(0)}(\xi_i)) - \frac{1}{|\det Df^{m_{j_i} - m_i}(\xi_i)|} \mu(U_n^{(0)}(\xi_{j_i})) \right]}{\sum_{i=1}^N \mu(U_n^{(0)}(\xi_i))}.$$

Proof. The proof is analogous to the proof of the non-periodic case presented in Corollary 4.5. \square

5.2. The multiplicity distribution. As mentioned before, by Proposition 2.10 of [FFR] we have that the multiplicity distribution of the limiting compound Poisson process for the REPP is given by the formula (27), whenever the limit exists.

Recall that we are assuming that (R3) holds.

Let I_1 be defined as in (30). For every $i \in \{1, \dots, N\}$, we define $j_{i,0} = i$. Then, for $i \in \{1, \dots, N\}$ we inductively define I_k in the following way.

Assuming that I_k is defined as well as $j_{1,k-1}, \dots, j_{N,k-1}$, then for every $i = 1, \dots, N$, we define $j_{i,k}$ to the index such that

$$\bigcup_{\ell=i+1}^{N+i} f^{-(m_{j_{\ell,k-1}} - m_i)}(U_n^{(0)}(\xi_{j_{\ell,k-1}})) \cap U_n^{(0)}(\xi_i) = f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i), \quad (31)$$

when $i \in I_k$ and $j_{i,k} = i$, otherwise.

Now, we define I_{k+1} as the set of indices $i \in \{1, \dots, N - k\}$ such that

$$\bigcup_{\ell=i+1}^{N+i} f^{-(m_{j_{\ell,k}} - m_i)}(U_n^{(0)}(\xi_{j_{\ell,k}})) \cap \partial U_n^{(0)}(\xi_i) = \emptyset.$$

Proposition 5.5. *The multiplicity distribution of the point process is given by equation (27) (whenever the limit exists), where*

$$\begin{aligned} \mu(\mathcal{A}_{q,n}^{(k)}) &= \sum_{i \in I_{k+1}} \left[\mu \left(f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i) \right) \right. \\ &\quad \left. - \mu \left(f^{-(m_{j_{i,k+1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}})) \cap U_n^{(0)}(\xi_i) \right) \right]. \end{aligned}$$

Proof. We will prove by induction that, for $i \in I_{k+1}$,

$$\mathcal{A}_{q,n}^{(k)}(\xi_i) = U_n^{(0)}(\xi_i) \cap f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) - f^{-(m_{j_{i,k+1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}}))$$

and

$$U_n^{(k+1)}(\xi_i) = f^{-(m_{j_{i,k+1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}})) \cap U_n^{(0)}(\xi_i),$$

and, for $i \notin I_{k+1}$, $\mu(\mathcal{A}_{q,n}^{(k)}(\xi_i)) = 0$ since $U_n^{(k)}(\xi_i) = U_n^{(k-1)}(\xi_i)$.

For $i \in I_1$ since $j_{\ell,0} = \ell$, we get

$$\begin{aligned} \mathcal{A}_{q,n}^{(0)}(\xi_i) &= U_n^{(0)}(\xi_i) - \bigcup_{\ell=i+1}^{N+i} f^{-(m_{\ell} - m_i)}(U_n^{(0)}(\xi_{\ell})) \\ &= U_n^{(0)}(\xi_i) - \bigcup_{\ell=i+1}^{N+i} f^{-(m_{j_{\ell,0}} - m_i)}(U_n^{(0)}(\xi_{j_{\ell,0}})) \\ &= U_n^{(0)}(\xi_i) \cap f^{-(m_{j_{i,0}} - m_i)}(U_n^{(0)}(\xi_{j_{i,0}})) - f^{-(m_{j_{i,1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,1}})), \end{aligned}$$

by definition of $j_{i,1}$, and, moreover, we have $U_n^{(1)}(\xi_i) = f^{-(m_{j_{i,1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,1}})) \cap U_n^{(0)}(\xi_i)$.

For $i \notin I_1$, $i \neq N$, $U_n^{(0)}(\xi_i) - \bigcup_{\ell=i+1}^{N+i} f^{-(m_{\ell} - m_i)}(U_n^{(0)}(\xi_{\ell})) = \emptyset$, so $\mu(\mathcal{A}_{q,n}^{(0)}(\xi_i)) = 0$ and $\mu(U_n^{(1)}(\xi_i)) = \mu(U_n^{(0)}(\xi_i))$.

By induction, for $i \in I_{k+1}$, we obtain

$$\begin{aligned} \mathcal{A}_{q,n}^{(k)}(\xi_i) &= U_n^{(k)}(\xi_i) - \bigcup_{\ell=i+1}^{N+i} f^{-(m_{\ell} - m_i)}(U_n^{(k)}(\xi_{\ell})) \\ &= U_n^{(k)}(\xi_i) - \bigcup_{\ell=i+1}^{N+i} f^{-(m_{\ell} - m_i)} \left(f^{-(m_{j_{\ell,k}} - m_i)}(U_n^{(0)}(\xi_{j_{\ell,k}})) \cap U_n^{(0)}(\xi_{\ell}) \right), \text{ by induction} \\ &= f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i) - \mu \left(\bigcup_{\ell=i+1}^{N+i} f^{-(m_{j_{\ell,k}} - m_i)}(U_n^{(0)}(\xi_{j_{\ell,k}})) \right), \text{ by induction} \\ &= f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i) - f^{-(m_{j_{i,k+1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}})), \text{ by definition of } j_{i,k+1}. \end{aligned}$$

and in particular we also have $U_n^{(k+1)}(\xi_i) = f^{-(m_{j_{i,k+1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}})) \cap U_n^{(0)}(\xi_i)$.

By induction, for $i \in I_{k+1}$, we obtain

$$\begin{aligned}
\mathcal{A}_{q,n}^{(k)}(\xi_i) &= U_n^{(k)}(\xi_i) - \bigcup_{\ell=i+1}^{N+i} f^{-(m_\ell - m_i)}(U_n^{(k)}(\xi_\ell)) \\
&= U_n^{(k)}(\xi_i) - \bigcup_{\ell=i+1}^{N+i} f^{-(m_\ell - m_i)} \left(f^{-(m_{j_{\ell,k}} - m_i)}(U_n^{(0)}(\xi_{j_{\ell,k}})) \cap U_n^{(0)}(\xi_i) \right), \text{ by induction} \\
&= f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i) - \mu \left(\bigcup_{\ell=i+1}^{N+i} f^{-(m_{j_{\ell,k}} - m_i)}(U_n^{(0)}(\xi_{j_{\ell,k}})) \right), \text{ by induction} \\
&= f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i) - f^{-(m_{j_{i,k+1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}})), \text{ by definition of } j_{i,k+1}.
\end{aligned}$$

and in particular we also have $U_n^{(k+1)}(\xi_i) = f^{-(m_{j_{i,k+1}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k+1}})) \cap U_n^{(0)}(\xi_i)$.

For $i \notin I_{k+1}$, $i \neq N - k + 1$ $\mu \left(\mathcal{A}_{q,n}^{(k)}(\xi_i) \right) = 0$ and

$$U_n^{(k+1)}(\xi_i) = f^{-(m_{j_{i,k}} - m_i)}(U_n^{(0)}(\xi_{j_{i,k}})) \cap U_n^{(0)}(\xi_i).$$

This concludes the proof. \square

Corollary 5.6. *Assume that μ is absolutely continuous with respect to the Lebesgue measure and (26) holds. Then the multiplicity distribution of the point process is given by formula (27), where*

$$\mu(\mathcal{A}_{q,n}^{(k)}) = \sum_{i \in I_{k+1}} \left(\alpha_n(m_{j_{i,k}}, m_i) - \alpha_n(m_{j_{i,k+1}}, m_i) \right) + \alpha_n(m_{j_{N-k,k}}, m_{N-k})$$

and

$$\alpha_n(a, b) = \frac{1}{|\det Df^{a-b}(f^b(\zeta))|} \mu \left(U_n^{(0)}(f^a(\zeta)) \right).$$

Proof. The proof is analogous to the proof of the non-periodic case presented in Corollary 4.7. \square

Corollary 5.7. *Assume that μ is absolutely continuous with respect to the Lebesgue measure and (26) holds and*

$$f^{-(m_{i+1} - m_i)} \left(U_n^{(0)}(\xi_{i+1}) \right) \subset U_n^{(0)}(\xi_i).$$

Then the multiplicity distribution of the point process is given by formula (27) where

$$\mu(\mathcal{A}_{q,n}^{(k)}) = \sum_{i \in I_{k+1}} \left(\alpha_n(m_{i+k}, m_i) - \alpha_n(m_{i+k+1}, m_i) \right) + \alpha_n(m_N, m_{N-k})$$

and $\alpha_n(a, b)$ is as in Corollary 5.6.

Proof. Using the notation of Proposition 5.5, note that in this case $j_{i,k} = i + k$. \square

5.3. Example. Let $(f, \mathbb{S}^1, Leb, \mathcal{B})$ be the dynamical system where $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ is the map given by $f(x) = 3x \pmod{1}$. Take the periodic point of period 5, $\zeta = \frac{81}{242}$, and an observable $\varphi : \mathbb{S}^1 \rightarrow \mathbb{R} \cup \{\infty\}$ such that

$$\varphi(x) = \begin{cases} -\log|x - \zeta|, & \text{if } x \in B_\epsilon(\zeta), \\ |x - f(\zeta)|^{-\frac{1}{2}}, & \text{if } x \in B_\epsilon(f(\zeta)) \\ |x - f^3(\zeta)|^{-\frac{1}{2}}, & \text{if } x \in B_\epsilon(f^3(\zeta)) \\ 0, & \text{otherwie,} \end{cases}$$

where we choose $\epsilon > 0$, such that $B_\epsilon(f^i(\zeta)) \cap B_\epsilon(f^j(\zeta)) = \emptyset$ for $i, j \in \{0, 1, 3\}$ and $i \neq j$. In this case we have that $u_F = \sup_{x \in \mathbb{S}^1} \varphi(x) = \infty$ and it occurs in $\zeta, f(\zeta)$ and $f^3(\zeta)$.

Given $\tau > 0$ and a sequence $(u_n)_{n \in \mathbb{N}}$ satisfying 3 we have that

$$U_n = (\zeta - e^{-u_n}, \zeta + e^{-u_n}) \cup (f(\zeta) - u_n^{-2}, f(\zeta) + u_n^{-2}) \cup (f^3(\zeta) - u_n^{-2}, f^3(\zeta) + u_n^{-2}).$$

We notice that for τ sufficiently large we have that

$$(\zeta - e^{-u_n}, \zeta + e^{-u_n}) \not\subseteq f^{-1}((f(\zeta) - u_n^{-2}, f(\zeta) + u_n^{-2})),$$

$$f^{-4}(\zeta - e^{-u_n}, \zeta + e^{-u_n}) \not\subseteq f^{-2}((f^3(\zeta) - u_n^{-2}, f^3(\zeta) + u_n^{-2})) \not\subseteq (f(\zeta) - u_n^{-2}, f(\zeta) + u_n^{-2}),$$

and

$$f^{-2}(\zeta - e^{-u_n}, \zeta + e^{-u_n}) \not\subseteq f^{-3}((f(\zeta) - u_n^{-2}, f(\zeta) + u_n^{-2})) \not\subseteq (f^3(\zeta) - u_n^{-2}, f^3(\zeta) + u_n^{-2}).$$

The extremal index: Noting that $m_1 = 0, m_2 = 1, m_3 = 3$, then $I_0 = \{2, 3\}$ and $j_2 =$

$3, j_3 = 5$. So, by Corollary 5.6 we have that

$$\begin{aligned} \theta &= \lim_{n \rightarrow \infty} \left(\frac{\left[\mu(U_n^{(0)}(\xi_i)) - \frac{1}{Df^{m_3-m_2}(f^{m_3}(\zeta))} \mu(U_n^{(0)}(f^{m_3}(\zeta))) \right]}{\sum_{i=1}^N \mu(U_n^{(0)}(\xi_i))} \right. \\ &\quad \left. + \frac{\left[\mu(U_n^{(0)}(\xi_i)) - \frac{1}{Df^{m_5-m_3}(f^{m_5}(\zeta))} \mu(U_n^{(0)}(f^{m_5}(\zeta))) \right]}{\sum_{i=1}^N \mu(U_n^{(0)}(\xi_i))} \right) \\ &= \lim_{n \rightarrow \infty} \frac{u_n^{-2} - \frac{1}{Df^2(f^3(\zeta))} u_n^{-2} + u_n^{-2} - \frac{1}{Df^3(f^2(\zeta))} u_n^{-2}}{e^{-u_n} + u_n^{-2} + u_n^{-2}} = \frac{2 - \left(\frac{1}{3}\right)^2 - \left(\frac{1}{3}\right)^3}{2} = \frac{25}{27}. \end{aligned}$$

The multiplicity distribution: To compute the multiplicity we will use Corollary 5.7.

First, we need to determine the sets I_k , for all $k \in \mathbb{N}$. It is not difficult to see that, in this case, $I_k = \{2, 3\}$, for all $k \in \mathbb{N}$. $m_{i+N} := m_i + p$, then $m_{3l-2} = 5l, m_{3l-1} = 5l+1, m_{3l} = 5l+3$. We now need to determine the value of $j_{i,k}$, for all $k \in \mathbb{N}$ and all $i \in I_k$. By definition $j_{i,0} = i$, for all $i \in \mathbb{N}$. We can verify by induction that

$$\begin{aligned} j_{2,2l} &= 3l - 2, & j_{2,2l-1} &= 3l - 3, \\ j_{3,2l} &= 3l - 3, & j_{3,2l-1} &= 3l - 5. \end{aligned}$$

In order to compute $\mathcal{A}_{q,n}^{(k)}$, note that in this case $q = 5$. So,

$$\begin{aligned}\mu(\mathcal{A}_{5,n}^{(k)}) &= \sum_{i \in I_{k+1}} \left(\alpha_n(m_{j_i,k}, m_i) - \alpha_n(m_{j_i,k+1}, m_i) \right) \\ &= \alpha(m_{j_{2,k}, m_2} - \alpha(m_{j_{2,k+1}, m_2}) + \alpha(m_{j_{3,k}, m_3}) - \alpha(m_{j_{3,k}, m_3}).\end{aligned}$$

Considering first the case $k = 2\ell$, with $\ell \in \mathbb{N}$, we have

$$\begin{aligned}\mu(\mathcal{A}_{5,n}^{(2\ell)}) &= \alpha(m_{j_{2,2\ell}, m_2} - \alpha(m_{j_{2,2\ell-1}, m_2}) + \alpha(m_{j_{3,2\ell}, m_3}) - \alpha(m_{j_{3,2\ell-1}, m_3}) \\ &= \alpha(m_{3\ell-2}, m_2) - \alpha(m_{3\ell-3}, m_2) + \alpha(m_{3\ell-3}, m_3) - \alpha(m_{3\ell-5}, m_3) \\ &= \left(\frac{1}{3}\right)^{5\ell} u_n^{-2} - \left(\frac{1}{3}\right)^{5\ell-2} u_n^{-2} + \left(\frac{1}{3}\right)^{5\ell} u_n^{-2} - \left(\frac{1}{3}\right)^{5\ell-3} u_n^{-2} = \frac{50}{27} \left(\frac{1}{3}\right)^{5\ell} u_n^{-2}.\end{aligned}$$

For the case $k = 2\ell + 1$ with $\ell \in \mathbb{N}_0$, we have

$$\begin{aligned}\mu(\mathcal{A}_{5,n}^{(2\ell+1)}) &= \alpha(m_{j_{2,2\ell-1}, m_2} - \alpha(m_{j_{2,2\ell-2}, m_2}) + \alpha(m_{j_{3,2\ell-1}, m_3}) - \alpha(m_{j_{3,2\ell-2}, m_3}) \\ &= \alpha(m_{3\ell-3}, m_2) - \alpha(m_{3\ell-5}, m_2) + \alpha(m_{3\ell-5}, m_3) - \alpha(m_{3\ell-6}, m_3) \\ &= \left(\frac{1}{3}\right)^{5\ell-2} u_n^{-2} - \left(\frac{1}{3}\right)^{5\ell-5} u_n^{-2} + \left(\frac{1}{3}\right)^{5\ell-3} u_n^{-2} - \left(\frac{1}{3}\right)^{5\ell-5} u_n^{-2} = \frac{34}{243} \left(\frac{1}{3}\right)^{5\ell} u_n^{-2}.\end{aligned}$$

Replacing these in the expression in Corollary 5.6 we obtain

$$\begin{aligned}\pi(2\ell + 1) &= \frac{\mu(\mathcal{A}_{5,n}^{(2\ell)}) - \mu(\mathcal{A}_{5,n}^{(2\ell+1)})}{\mu(\mathcal{A}_{5,n}^{(0)})} = \frac{\frac{50}{27} \left(\frac{1}{3}\right)^{5\ell} u_n^{-2} - \frac{34}{243} \left(\frac{1}{3}\right)^{5\ell} u_n^{-2}}{\frac{50}{27} u_n^{-2}} = \frac{208}{225} \left(\frac{1}{3}\right)^{5\ell}, \text{ for } \ell \in \mathbb{N}_0, \\ \pi(2\ell) &= \frac{\mu(\mathcal{A}_{5,n}^{(2\ell-1)}) - \mu(\mathcal{A}_{5,n}^{(2\ell)})}{\mu(\mathcal{A}_{5,n}^{(0)})} = \frac{\frac{34}{243} \left(\frac{1}{3}\right)^{5(\ell-1)} u_n^{-2} - \frac{50}{27} \left(\frac{1}{3}\right)^{5\ell} u_n^{-2}}{\frac{50}{27} u_n^{-2}} = \frac{868}{50} \left(\frac{1}{3}\right)^{5\ell}, \text{ for } \ell \in \mathbb{N}.\end{aligned}$$

For this particular example, we performed a simulation study and computed a finite sample multiplicity distribution. In this simulation we chose at random an initial condition $x_0 = 0.08918848967830795$ that was iterated 500 000 times by the system. The threshold taken was $u = 44.721359549995796$ which corresponds to the 0.998-quantile of the distribution of X_0 . The results are presented in Figure 2 and reveal a good agreement with the theoretical estimates provided by π computed above.

5.4. Non correlated multiple maxima including periodic points. We discuss briefly the possibility of having multiple maximal points with non-intersecting orbits as in Section 4.4 but now allowing the possibility of having maximal points that are periodic. By [FFT13, Theorem 2] and [AFV14, Theorem A], we have that for every continuous system under Assumption A, for every $\zeta \in \mathcal{X}$ taken as the only maximum of φ , then there exists an EVL with an EI that is 1 if the point is not periodic or is less than 1 (whose value depends on the expansion rate at ζ). Hence, we are going to consider multiple maximal points ξ_1, \dots, ξ_N and to each such point we associate the respective EI guaranteed by [FFT13, Theorem 2] and [AFV14, Theorem A] (see also [Fre13, Theorem 8]), which we denote by θ_i , being that if ξ_i is periodic then $\theta_i < 1$.

Proposition 5.8. *Assume that f is continuous and satisfies Assumption A. Let X_0, X_1, \dots be given by (1), where the observable φ has N maximal points ξ_1, \dots, ξ_N and let θ_i be the corresponding EI. Let p_i be the period of ζ_i (we set $p_i = 0$ when ζ_i is not periodic). Assume that (29) holds and that the following limits exist for every $i = 1, \dots, N$*

$$\lim_{n \rightarrow \infty} \frac{\mu(U_n(\xi_i))}{\mu(U(u_n))} =: \alpha_i,$$

where $U_n(\xi_i)$ is a connected component of $U(u_n)$ as in (24). Then X_0, X_1, \dots satisfies conditions $\mathbb{D}_q(u_n)$ and $\mathbb{D}'_q(u_n)$, for $q = \max_{i=1, \dots, N} p_i$, and consequently for $(u_n)_{n \in \mathbb{N}}$ satisfying (3) then $\lim_{n \rightarrow \infty} \mu(M_n \leq u_n) = e^{-\theta\tau}$, where $\theta = \sum_{i=1}^N \alpha_i \theta_i$.

Proof. Conditions $\mathbb{D}_q(u_n)$ and $\mathbb{D}'_q(u_n)$ follow from the computations in Section 3.3 as long as one check that $\lim_{n \rightarrow \infty} R(\mathcal{A}_{q,n}) = +\infty$. This last statement follows from the definition of $\mathcal{A}_{q,n}$ and a continuity argument very similar to the proof of Lemma 4.1. \square

6. CLUSTERING PATTERNS

As we have seen in Sections 4 and 5, multiple correlated maxima can be used as a mechanism to create clustering, alter the value of EI and produce different multiplicity distributions for the limit of REPP. When compared to the usual method of introducing clustering in a dynamical setting, which was essentially based on considering a global maximum of φ achieved at a unique repelling periodic point, this method of forging clustering by means of considering multiple correlated maxima is much richer.

In fact, we have already seen that, on the contrary to single maximum at a repelling periodic point for which one always obtains a geometric distribution for the multiplicity distribution of the limiting compound Poisson process for the REPP, with multiple correlated maxima we can obtain different multiplicity distributions for the limit of the REPP.

But the richness of this mechanism can also be appreciated by simply looking at the observed data and the corresponding clustering patterns. Observe that the systems we are studying are either uniformly expanding or non-uniformly expanding, which means that the periodic points should be repelling. When the maximum is achieved at a single repelling point the clustering pattern is very simple: it consists of a very large observation corresponding to the first exceedance, which is followed by a strictly decreasing cluster of exceedances observed after a precise number of observations corresponding to the period (see Figure 3). This is because in order to have a cluster of exceedances the orbit needs to enter in a very small neighbourhood of the periodic point ζ of period p . Let us assume we have a very high exceedance, which of course means that the orbit went very close to ζ . Once inside this very small neighbourhood of ζ , the periodicity of ζ forces the appearance of another exceedance after p observations but since ζ is repelling this return to a small neighbourhood of ζ is not that deep, *i.e.*, the orbit is pulled away from ζ which explains the fading of the exceedances.

In order to illustrate how easily one can create a different clustering pattern, where in particular one can observe that the exceedances may grow inside a cluster, we consider the following example.

We consider the system uniformly expanding map

$$\begin{aligned} f : S^1 &\longrightarrow S^1 \\ x &\mapsto 3x \pmod{1} \end{aligned} \tag{32}$$

and the observable $\varphi : \mathcal{S}^1 \rightarrow \mathbb{R}$, where $\varphi(x) = -\log|x - \frac{1}{4}|\mathbf{1}_{[0, \frac{1}{2}]} + |x - \frac{3}{4}|^{-\frac{1}{3}}\mathbf{1}_{(\frac{1}{2}, 1]}$. We notice that for this observable the maximum value is $u_F = \sup_{x \in \mathcal{S}^1} \varphi(x) = \infty$, which is achieved at the points $\zeta = \frac{1}{4}$ and $f(\zeta) = \frac{3}{4}$. As $\zeta = f^2(\zeta)$, the maxima correspond to the points of a periodic orbit of period 2.

Note that for u sufficiently large, $\{X_0 > u\} = (1/4 - e^{-u}, 1/4 + e^{-u}) \cup (3/4 - u^{-3}, 3/4 + u^{-3})$. So an exceedance of u occurs when the orbit enters a ball of radius e^{-u} around $1/4$ or a ball of radius u^{-3} around $3/4$. Note that if we have an exceedance because the orbit hits the point $x = 1/4 + e^{-u}/3$ and $\varphi(x) = u + \log 3 > u$, then in the next iterate the orbit hits the point $f(x) = 3/4 + e^{-u}$ and $\varphi(f(x)) = e^{u/3} \gg u + \log 3 > u$, for u sufficiently large so that we have an exceedance that is followed by another exceedance that is even higher than the first.

This way, we create different clustering patterns which can be fully appreciated by the simulation study we performed. We chose a random point $x \in [0, 1]$ (using the uniform distribution); we iterated the point 2000 times by f ; we considered u to be such that $e^{-u} + \frac{1}{u^3} = \frac{40}{2000}$ (meaning that the expected number of exceedances of u is 40). In Figure 4 we can see the whole time series. In Figure 5 we can see the observations forming a cluster that presents clearly a different pattern. In particular, we observe exceedances being followed by even higher exceedances, which was forbidden in the case of a single maximum as portrayed in Figure 3.

7. COMPETITION BETWEEN DOMAINS OF ATTRACTION

Up until now, we have addressed the issues regarding the effect of having multiple maxima on the existence of EVL, the appearance of clustering, the consequences on the value of the EI and the impact on the limit of REPP. This has been done for normalising sequences $(u_n)_{n \in \mathbb{N}}$ such that condition (3) holds. Typically, these normalising sequences are taken as a one parameter family of linear sequences such that $u_n = y/a_n + b_n$, where $(a_n)_{n \in \mathbb{N}}$, $(b_n)_{n \in \mathbb{N}}$, with $a_n > 0$, and one usually looks for non-degenerate limit distributions for $\mathbb{P}(a_n(M_n - b_n) \leq y)$.

The limit is then $e^{-\tau(y)}$, where $\tau(y)$ is a function of y , which is connected to the type of observable we have. For well behaved measures, if the observable satisfies (20) then one gets that $\tau(y) = e^{-y}$ meaning that the the stochastic process falls in the domain of attraction of a Gumbel law (or type 1); if the observable satisfies (21) then one gets that $\tau(y) = y^{-\alpha}$ (for $y > 0$ and $\alpha > 0$) meaning that the the stochastic process falls in the domain of attraction of a Fréchet law (or type 2); observable satisfies (22) then one gets that $\tau(y) = (-y)^\alpha$ (for $y \leq 0$ and $\alpha > 0$) meaning that the the stochastic process falls in the domain of attraction of a Weibull law (or type 3).

In the independent and identically distributed (i.i.d.) setting the domain of attraction is determined by the tail of the distribution F , where $F(u) = \mu(X_0 \leq u)$. As it was observed in [FFT10, Remark 1], in the dynamical setting (with a single maximum), the shape of the observable determines the type of tail of the distribution F and, once the existence is established by the assumptions on the system, then the domain of attraction is determined by the behaviour of the observable at the maximum.

In here, since we have multiple maxima, then we may have different types of behaviour of the observable at the different maximal points. This creates a sort of competition between different domains of attraction, which in fact is not that difficult to settle. First, we start by noting that if the observable is of type 2 (satisfies (21)) then $\sup \varphi = h(0) = \infty$; if the observable is of type 3 (satisfies (22)) then $\sup \varphi = h(0) = D < \infty$ and if the observable is of type 1 (satisfies (20)) then the maximum value can be either finite or infinite. Hence, since

all maxima achieve the same maximum value, we can only have competitions between Fréchet and Gumbel and between Weibull and Gumbel.

Note that the appearance of clustering and consequently of an EI less than 1 does not change the type of limit law that applies when we consider one parameter linear sequences. This is the statement of [LLR83, Corollary 3.7.3], which also adds that, in fact, we can even perform a linear change of the normalising constants a_n and b_n so that the exact same limit applies as in the corresponding independent case. Hence in what follows we restrict the study to the behaviour of $\mu(U(u)) = 1 - F(u)$ instead of $\mu(A^{(q)}(u))$ as u goes to $\sup \varphi$.

In what follows we establish the reigning domain of attraction in each such situation. For that purpose, we introduce the following notation $\bar{F}(x) = 1 - F(x)$.

7.1. Gumbel versus Fréchet.

Proposition 7.1. *Let $\varphi : \mathcal{X} \rightarrow \mathbb{R} \cup \{\infty\}$ be an observable which has the following form*

$$\varphi(x) = \varphi_1(x)\mathbf{1}_{U(\zeta)}(x) + \varphi_2(x)\mathbf{1}_{U(f^i(\zeta))}(x),$$

where φ_1 is an observable defined in the neighborhood $U(\zeta)$ of ζ , φ_2 is an observable defined in the neighborhood $U(f^i(\zeta))$ of $f^i(\zeta)$ and they satisfy $\max_x \varphi_1(x) = \varphi_1(\zeta) = \infty = \max_x \varphi_2(x) = \varphi_2(f^i(\zeta))$. Let F_i be the distribution function of φ_i . If F_1 and F_2 belong to the domain of attraction for maxima of the Gumbel and the Fréchet distribution, respectively, then the distribution function of φ belongs to the domain of attraction of the Fréchet distribution.

Proof. We start by noting that if F_1 belongs to the domain of attraction of the Gumbel distribution, with infinite right endpoint of the support, then we may write

$$1 - F_1(u) = u^p e^{-u^q} L_1(u), \tag{33}$$

where $p \in \mathbb{R}$, $q > 0$ and L_1 is a function of slow variation at infinity, i.e., $L_1 : (0, +\infty) \rightarrow (0, +\infty)$ and, for all $t > 0$,

$$\lim_{u \rightarrow \infty} \frac{L_1(tu)}{L_1(u)} = 1$$

(cf. [Gom84]).

F_2 belongs to the domain of attraction of the Fréchet distribution if and only if

$$1 - F_2(u) = u^{-\alpha} L_2(u), \tag{34}$$

where $\alpha > 0$ and L_2 is a function of slow variation at infinity (cf. [dH70]).

If

$$\varphi(x) = \varphi_1(x)\mathbf{1}_{U(\zeta)}(x) + \varphi_2(x)\mathbf{1}_{U(f^i(\zeta))}(x),$$

and $X_0 = \varphi$, then

$$\mu(\varphi > u) = \mu(X_0 > u) = u^p e^{-u^q} L_1(u) + u^{-\alpha} L_2(u) = u^{-\alpha} (L_2(u) + u^{\alpha+p} e^{-u^q} L_1(u)).$$

As $\lim_{u \rightarrow \infty} u^{\alpha+p} e^{-u^q} L_1(u) = 0$ we get that the function $L(u) = L_2(u) + u^{\alpha+p} e^{-u^q} L_1(u)$ is a function of slow variation at infinity. Hence, the distribution function of $\varphi(x)$ belongs to the domain of attraction of the Fréchet distribution. \square

7.2. Gumbel versus Weibull.

Proposition 7.2. *Let $\varphi : \mathcal{X} \rightarrow \mathbb{R} \cup \{\infty\}$ be an observable which has the following form*

$$\varphi(x) = \varphi_1(x)\mathbf{1}_{U(\zeta)}(x) + \varphi_2(x)\mathbf{1}_{U(f^i(\zeta))}(x),$$

where φ_1 is an observable defined in the neighborhood $U(\zeta)$ of ζ , φ_2 is an observable defined in the neighborhood $U(f^i(\zeta))$ of $f^i(\zeta)$ and they satisfy $\max_x \varphi_1(x) = \varphi_1(\zeta) = D = \max_x \varphi_2(x) = \varphi_2(f^i(\zeta))$. Let F_i be the distribution function of φ_i . If F_1 and F_2 belong to the domain of attraction for maxima of the Gumbel and the Weibull distribution, respectively, then the distribution function of φ , F , belongs to the domain of attraction of the Weibull distribution.

Proof. If $X_0 = \varphi$, then

$$1 - F(u) = \mu(\varphi > u) = \mu(X_0 > u) = (1 - F_1(u)) + (1 - F_2(u)).$$

So,

$$\begin{aligned} \lim_{s \rightarrow 0} \frac{1 - F(D - sy)}{1 - F(D - s)} &= \lim_{s \rightarrow 0} \frac{1 - F_1(D - sy) + 1 - F_2(D - sy)}{1 - F_1(D - s) + 1 - F_2(D - s)} \\ &= \lim_{s \rightarrow 0} \frac{(1 - F_2(D - sy)) \left(\frac{1 - F_1(D - sy)}{1 - F_2(D - sy)} + 1 \right)}{(1 - F_2(D - s)) \left(\frac{1 - F_1(D - s)}{1 - F_2(D - s)} + 1 \right)}. \end{aligned} \quad (35)$$

As F_2 belongs to the domain of attraction of the Weibull distribution, then

$$\lim_{s \rightarrow 0} \frac{1 - F_2(D - sy)}{1 - F_2(D - s)} = y^\alpha, \quad (36)$$

for some $\alpha > 0$, and

$$1 - F_2(D - s) = s^\alpha L(s), \quad (37)$$

where L is a function of slow variation at zero, i.e., for all $t > 0$,

$$\lim_{s \rightarrow 0} \frac{L(ts)}{L(s)} = 1.$$

We note now that if F_1 belongs to the domain of attraction of the Gumbel distribution, with finite right endpoint of the support D , then, for all $\beta > 0$,

$$\lim_{s \rightarrow 0} \frac{1 - F_1(D - s)}{s^\beta} = 0.$$

Thus, by (37), we have that

$$\lim_{s \rightarrow 0} \frac{1 - F_1(D - s)}{1 - F_2(D - s)} = 0. \quad (38)$$

Using now (36) and (37) in (35), we obtain that

$$\lim_{s \rightarrow 0} \frac{1 - F(D - sy)}{1 - F(D - s)} = y^\alpha,$$

and so F belongs to the domain of attraction of the Weibull distribution. \square

REFERENCES

- [AFV14] Hale Aytac, Jorge Milhazes Freitas, and Sandro Vaienti, *Laws of rare events for deterministic and random dynamical systems*, Trans. Amer. Math. Soc. **Published online with DOI: 10.1090/S0002-9947-2014-06300-9** (2014).
- [BG97] Abraham Boyarsky and Pawel Góra, *Laws of chaos*, Probability and its Applications, Birkhäuser Boston Inc., Boston, MA, 1997, Invariant measures and dynamical systems in one dimension. MR 1461536 (99a:58102)
- [BGTS04] Jan Beirlant, Yuri Goegebeur, Jozef Teugels, and Johan Segers, *Statistics of extremes*, Wiley Series in Probability and Statistics, John Wiley & Sons, Ltd., Chichester, 2004, Theory and applications, With contributions from Daniel De Waal and Chris Ferro. MR 2108013 (2005j:62002)
- [BSTV03] H. Bruin, B. Saussol, S. Troubetzkoy, and S. Vaienti, *Return time statistics via inducing*, Ergodic Theory Dynam. Systems **23** (2003), no. 4, 991–1013. MR MR1997964 (2005a:37004)
- [CC13] J.-R. Chazottes and P. Collet, *Poisson approximation for the number of visits to balls in non-uniformly hyperbolic dynamical systems*, Ergodic Theory Dynam. Systems **33** (2013), no. 1, 49–80. MR 3009103
- [CCC09] Jean-René Chazottes, Zaqueu Coelho, and Pierre Collet, *Poisson processes for subsystems of finite type in symbolic dynamics*, Stoch. Dyn. **9** (2009), no. 3, 393–422. MR 2566908 (2011g:37014)
- [Col01] P. Collet, *Statistics of closest return for some non-uniformly hyperbolic systems*, Ergodic Theory Dynam. Systems **21** (2001), no. 2, 401–420. MR MR1827111 (2002a:37038)
- [dH70] L. de Haan, *On regular variation and its application to the weak convergence of sample extremes*, Mathematical Centre Tracts, vol. 32, Mathematisch Centrum, Amsterdam, 1970. MR 0286156 (44 #3370)
- [FF08] Ana Cristina Moreira Freitas and Jorge Milhazes Freitas, *On the link between dependence and independence in extreme value theory for dynamical systems*, Statist. Probab. Lett. **78** (2008), no. 9, 1088–1093. MR MR2422964 (2009e:37006)
- [FFR] Ana Cristina Moreira Freitas, Jorge Milhazes Freitas, and Fagner Bernardini Rodrigues, *The speed of convergence of rare events point processes in non-uniformly hyperbolic systems*, in preparation.
- [FFT10] Ana Cristina Moreira Freitas, Jorge Milhazes Freitas, and Mike Todd, *Hitting time statistics and extreme value theory*, Probab. Theory Related Fields **147** (2010), no. 3-4, 675–710. MR 2639719 (2011g:37015)
- [FFT11] ———, *Extreme value laws in dynamical systems for non-smooth observations*, J. Stat. Phys. **142** (2011), no. 1, 108–126. MR 2749711 (2012a:60149)
- [FFT12] ———, *The extremal index, hitting time statistics and periodicity*, Adv. Math. **231** (2012), no. 5, 2626–2665. MR 2970462
- [FFT13] ———, *The compound Poisson limit ruling periodic extreme behaviour of non-uniformly hyperbolic dynamics*, Comm. Math. Phys. **321** (2013), no. 2, 483–527. MR 3063917
- [FFT15] ———, *Speed of convergence for laws of rare events and escape rates*, Stochastic Process. Appl. **125** (2015), no. 4, 1653–1687.
- [FFTV15] Ana Cristina Moreira Freitas, Jorge Milhazes Freitas, Mike Todd, and Sandro Vaienti, *Rare events for the manneville-pomeau map*, Preprint arXiv:1503.01372, March 2015.
- [FP12] Andrew Ferguson and Mark Pollicott, *Escape rates for gibbs measures*, Ergod. Theory Dynam. Systems **32** (2012), 961–988.
- [Fre13] Jorge Milhazes Freitas, *Extremal behaviour of chaotic dynamics*, Dyn. Syst. **28** (2013), no. 3, 302–332.
- [Gom84] M. Ivette Gomes, *Penultimate limiting forms in extreme value theory*, Ann. Inst. Statist. Math. **36** (1984), no. 1, 71–85. MR 752007 (85m:60065)
- [Hir93] Masaki Hirata, *Poisson law for Axiom A diffeomorphisms*, Ergodic Theory Dynam. Systems **13** (1993), no. 3, 533–556. MR MR1245828 (94m:58137)
- [HNT12] Mark Holland, Matthew Nicol, and Andrei Török, *Extreme value theory for non-uniformly expanding dynamical systems*, Trans. Amer. Math. Soc. **364** (2012), no. 2, 661–688. MR 2846347 (2012k:37064)
- [HSV99] Masaki Hirata, Benoît Saussol, and Sandro Vaienti, *Statistics of return times: a general framework and new applications*, Comm. Math. Phys. **206** (1999), no. 1, 33–55. MR MR1736991 (2001c:37007)

- [HV09] Nicolai Haydn and Sandro Vaienti, *The compound Poisson distribution and return times in dynamical systems*, Probab. Theory Related Fields **144** (2009), no. 3-4, 517–542. MR MR2496441
- [HWZ14] Nicolai Haydn, Nicole Winterberg, and Roland Zweimüller, *Return-time statistics, hitting-time statistics and inducing*, Ergodic theory, open dynamics, and coherent structures, Springer Proc. Math. Stat., vol. 70, Springer, New York, 2014, pp. 217–227. MR 3213501
- [Kal86] Olav Kallenberg, *Random measures*, fourth ed., Akademie-Verlag, Berlin, 1986. MR 854102 (87k:60137)
- [Kel12] Gerhard Keller, *Rare events, exponential hitting times and extremal indices via spectral perturbation*, Dyn. Syst. **27** (2012), no. 1, 11–27. MR 2903242
- [KL09] Gerhard Keller and Carlangelo Liverani, *Rare events, escape rates and quasistationarity: some exact formulae*, J. Stat. Phys. **135** (2009), no. 3, 519–534. MR 2535206 (2011a:37012)
- [LLR83] M. R. Leadbetter, Georg Lindgren, and Holger Rootzén, *Extremes and related properties of random sequences and processes*, Springer Series in Statistics, Springer-Verlag, New York, 1983. MR MR691492 (84h:60050)
- [LSV99] Carlangelo Liverani, Benoît Saussol, and Sandro Vaienti, *A probabilistic approach to intermittency*, Ergodic Theory Dynam. Systems **19** (1999), no. 3, 671–685. MR MR1695915 (2000d:37029)
- [PS14] Françoise Pène and Benoît Saussol, *Poisson law for some nonuniformly hyperbolic dynamical systems with polynomial rate of mixing*, Preprint arXiv:1401.3599, 2014.
- [Ryc83] Marek Rychlik, *Bounded variation and invariant measures*, Studia Math. **76** (1983), no. 1, 69–80. MR MR728198 (85h:28019)
- [Sau00] Benoît Saussol, *Absolutely continuous invariant measures for multidimensional expanding maps*, Israel J. Math. **116** (2000), 223–248. MR 1759406 (2001e:37037)
- [Sau09] Benoît Saussol, *An introduction to quantitative Poincaré recurrence in dynamical systems*, Rev. Math. Phys. **21** (2009), no. 8, 949–979. MR MR2568049
- [Smi88] Richard L. Smith, *A counterexample concerning the extremal index*, Adv. in Appl. Probab. **20** (1988), no. 3, 681–683. MR 955511 (89g:60136)




FIGURE 1. The histogram gives in the y -axis the frequency of the cluster size appearing in the x -axis. Each dot depicted corresponds to the estimate obtained from the distribution π .




FIGURE 2. The histogram gives in the y -axis the frequency of the cluster size appearing in the x -axis. Each dot depicted corresponds to the estimate obtained from the distribution π .




FIGURE 3. Cluster of observations for an observable with global maximum achieved at the fixed point $1/2$ of the system f given by (32)




FIGURE 4. Time series for 2000 runs of the orbit of the point $x = 0.7756592465669858$ with the level $u = 4.619613119957849$ depicted in red




FIGURE 5. Cluster occurring between the observations 1446 and 1450 of the time series in Figure 4

DAVIDE AZEVEDO, CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL
E-mail address: davidemsa@gmail.com

ANA CRISTINA MOREIRA FREITAS, CENTRO DE MATEMÁTICA & FACULDADE DE ECONOMIA DA UNIVERSIDADE DO PORTO, RUA DR. ROBERTO FRIAS, 4200-464 PORTO, PORTUGAL
E-mail address: amoreira@fep.up.pt

JORGE MILHAZES FREITAS, CENTRO DE MATEMÁTICA & FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL
E-mail address: jmfreira@fc.up.pt
URL: <http://www.fc.up.pt/pessoas/jmfreira>

FAGNER BERNARDINI RODRIGUES, INSTITUTO DE MATEMÁTICA - UNIVERSIDADE FEDERAL DO RIO GRANDE DO SUL, AV. BENTO GONÇALVES, 9500 - PRÉDIO 43-111 - AGRONOMIA, CAIXA POSTAL 15080 91509-900 PORTO ALEGRE - RS - BRASIL
E-mail address: fagnerbernardini@gmail.com