

Integrated Master in Bioengineering

**Antimicrobial activity of selected phytochemicals
against *Escherichia coli* and *Staphylococcus aureus*
cells and biofilms**

**Dissertation for Master Degree in Bioengineering – Specialization in Biological
Engineering**

Joana Isabel Carvalho Monte

Supervisor:
Manuel José Vieira Simões (PhD)



Universidade do Porto

Faculdade de Engenharia

FEUP

June 2013

The present thesis was developed for the obtention of Master degree in Bioengineering, in the Faculty of Engineering of University of Porto.

The work was carried out at LEPAE during 6 months. The main objective of the thesis was the evaluation of the efficacy of phytochemicals against *Escherichia coli* and *Staphylococcus aureus* planktonic cells and biofilms.

The judge that approved the present document was composed by three elements: Luís de Melo (Cathedratric Professor), Maria da Conceição Fernandes (PhD), Manuel Simões (PhD).

The author

(Joana Isabel Carvalho Monte)

The supervisor

(Manuel Simões)

“Try to learn something about everything,
and everything about something.”

Thomas H. Huxley

Acknowledgments

Em primeiro lugar, gostaria de agradecer ao meu orientador, o Professor Doutor Manuel Simões, por toda a dedicação, disponibilidade e empenho ao longo da duração deste projeto.

Gostaria também de agradecer à Ana Abreu, pela incansável ajuda e auxílio na realização deste trabalho. Estou também grata à Anabela e à Joana pelo apoio que me deram nalguns dos ensaios realizados no laboratório.

Quero agradecer ainda a todos do laboratório E007, Carla, Luciana, Ritas, Catarina, Paula, Joanas, Madalena e Renato pela disponibilidade e simpatia.

A todos os meus amigos, em especial às minhas colegas de laboratório Inês, Helena e Carolina pelos ótimos momentos proporcionados e pela sua companhia. Quero agradecer ainda ao João Manuel, ao Francisco, à Catarina e ao João Paulo, por estarem incondicionalmente do meu lado durante os últimos cinco anos. Um especial obrigada ao Nelson pela companhia, apoio e gargalhadas que me proporcionou. À Ana e à Cató por serem as minhas amigas, companheiras e confidentes e por sempre estarem do meu lado.

Finalmente, de modo muito especial, um obrigada à minha família, mãe, pai, Helena e Marta, pelo incentivo, paciência e apoio incondicional durante a realização deste projeto.

Abstract

Antimicrobial resistance is one of the biggest problems facing global public health. The effectiveness of antimicrobial drugs has been lost due to the evolution of pathogen resistance. Plants are considered the greatest source to obtain new antimicrobials. They produce secondary metabolites, phytochemicals, which protect the plant against pathogens.

The aim of this study was to assess the antimicrobial activity of four phytochemicals - 7-hydroxycoumarin (7-HC), indole-3-carbinol (I3C), salicylic acid (SA) and saponin (SP) – against *Escherichia coli* and *Staphylococcus aureus* and also understand their ability to control biofilm formation.

Several experiments were carried out in order to: i) test the ability of phytochemicals to control planktonic bacteria growth through the measurement of minimal inhibitory concentration (MIC) and the minimal bactericidal concentration (MBC); ii) evaluate the phytochemicals action in the control of biofilms; iii) understand aspects of the phytochemicals mode of action against the bacteria.

Results have shown that MIC values were higher for *E. coli* than for *S. aureus*. The 7-HC and I3C were the most effective, with MICs of 200 and 400 µg/mL for *S. aureus*, respectively, and 800 µg/mL against *E. coli*. Regarding MBC, 1600 and 5000 µg/mL were obtained for I3C and SA, respectively. It was also observed that 7-HC and SP has no significant effect in surface charge of *E. coli*; in contrast, I3C and SA make the membrane more and less negative, respectively. *S. aureus* surface charge was changed in contact with SA and SP. It was observed that phytochemical concentration did not affect the biofilm removal for both bacteria. *E. coli* biofilms are more susceptible to phytochemicals comparing to *S. aureus* biofilms.

SA and SP promoted the increase and decrease of hydrophilic properties of *E. coli*, respectively. *S. aureus* became less hydrophilic in contact with 7-HC and SA. *E. coli* showed the highest motility and also an increasing in swimming and swarming motility over time. Motility was mostly affected when I3C was added. Swimming and sliding motilities were completely inhibited and swarming motility was not affected by I3C. The quorum-sensing results indicated that inhibition of violacein production was detectable with 7-HC, I3C and SA, with halos ranging from 5 to 19 mm. I3C was also the most effective phytochemical. The increasing of concentration resulted in an

increasing of pigment inhibition zone. The OMPs expression in *E. coli* was not affect after the exposure to phytochemicals. Dual combinations between antibiotics and I3C produced synergistic effects against *S. aureus* resistant strains.

This study suggests that 7-HC and I3C are the most important phytochemicals against *E. coli* and *S. aureus*. Both phytochemicals affected the motility and QS activity, which means that they can play an important role in the interference of cell-cell interactions and in biofilm control.

Resumo

A resistência antimicrobiana é um dos maiores problemas enfrentados pela saúde pública global. A eficácia dos agentes antimicrobianos é cada vez mais inferior devido à evolução de mecanismos de resistência a infecções e microrganismos patogénicos. As plantas são consideradas a melhor fonte para a obtenção de novos agentes antimicrobianos. Estas produzem metabolitos secundários, os fitoquímicos, que fazem parte de mecanismos de proteção da planta contra agentes invasores e patogénicos.

O principal objetivo foi a avaliação da actividade antimicrobiana de 4 fitoquímicos – 7-hidroxycumarina (7-HC), indol-3-carbinol (I3C), ácido salicílico (SA) e saponina (SP) – nas estirpes *Escherichia coli* e *Staphylococcus aureus* e também compreender a sua capacidade de controlo na formação de biofilmes.

Vários estudos foram realizados de modo a: i) testar a capacidade dos fitoquímicos no controlo do crescimento bacteriano de células planctónicas através da deteção da concentração mínima inibitória (CMI) e concentração mínima bacteriana (CMB); ii) avaliar a ação dos fitoquímicos no controlo de biofilmes; iii) entender os aspetos dos fitoquímicos no modo de ação contra as bactérias.

Os resultados mostraram que os valores de CMI são mais elevados para *E. coli* do que para *S. aureus*. A 7-HC e I3C foram os mais eficazes, obtendo-se valores de CIM de 200 e 400 µg/mL para *S. aureus*, respectivamente, e 800 µg/mL para *E. coli*. Relativamente a CMB, obtiveram-se valores de 1600 e 5000 µg/mL para I3C e SA, respectivamente. Foi também observado que 7-HC e SP não tiveram efeito significativo na carga da membrana da *E. coli*; ao contrário de I3C e SA que tornaram a carga da membrana mais e menos negativa, respectivamente. A carga da membrana de *S. aureus* foi alterada em contato com SA e SP. Foi verificado que a concentração de fitoquímico não influencia a remoção de biofilmes para ambas as bactérias. Os biofilmes de *E. coli* são mais susceptíveis aos fitoquímicos comparativamente aos biofilmes de *S. aureus*.

O SA e a SP promoveram o aumento e diminuição das propriedades hidrofílicas da *E. coli*, respectivamente. *S. aureus* tornou-se menos hidrofílico em contato com 7-HC e SA. *E. coli* apresentou a mobilidade mais elevada e também um aumento na mobilidade *swimming* e *swarming* ao longo do tempo. A mobilidade foi

maioritariamente afetada pela adição de I3C. A mobilidade *swimming* e *sliding* foi completamente inibida e *swarming* foi afetada após a adição de I3C. Os resultados de quorum-sensing indicaram que a inibição da produção de violaceína foi detetada com 7-HC, I3C e SA, com halos compreendidos entre 5 e 19 mm. I3C foi também o fitoquímico mais eficaz. O aumento da concentração resultou num aumento da zona de inibição de pigmento. A expressão das OMPs na *E. coli* não foi afetada após a exposição aos fitoquímicos. A combinação dupla entre os antibióticos e I3C produziu efeitos sinérgicos contra estirpes resistentes de *S. aureus*.

Este estudo sugere que 7-HC e I3C são os fitoquímicos mais promissores contra *E. coli* e *S. aureus*. Ambos os fitoquímicos afetam a mobilidade e a actividade de quorum-sensing, o que significa que apresentam um papel determinante na interferência de interações célula-célula e no controlo de biofilmes.

Content List

Acknowledgments	i
Abstract	iii
Resumo	v
Content List	vii
Figures List	ix
Tables List	xi
Glossary	xiii
Chapter 1	1
Work Outline	1
1.1. Background and Project Presentation.....	1
1.2 Main objectives.....	2
1.3 Thesis Organization.....	3
Chapter 2	5
Literature Review	5
2.1 Antimicrobial resistance and phytochemicals.....	5
2.2. A new therapy against resistance – Phytotherapy.....	7
2.3. Phytochemicals and their classes	9
2.4. Mode of Action of Phytochemicals	12
2.5. Biofilms and phytochemicals:	12
Chapter 3	17
Activity of Selected Phytochemical Products as Antimicrobials and in Biofilm Control	17
3.1 Introduction	17
3.2 Matherial and methods	19
3.2.1. Bacterial Strains	19
3.2.2. Phytochemicals.....	19
3.2.3. Determination of Minimum Inhibitory Concentration.....	19
3.2.4. Determination of Minimum Bactericidal Concentration.....	20
3.2.5. Biofilm formation and control in sterile 96-well polystyrene microtiter plates	20
3.2.6. Biofilm analysis	21
3.2.6.1 Crystal Violet method.....	21
3.2.6.2 Resazurin Method.....	21
3.2.7. Classification of biofilm producer bacteria	22
3.2.8. Statistical analysis.....	22
3.3 Results and Discussion.....	23
3.3.1. Antimicrobial activity of phytochemicals and biofilm control potential.....	23
Chapter 4	29
Aspects Underlying the Antibacterial and Biofilm Control Action of Phytochemicals	29
4.1 Introduction	29
4.2 Materials and methods.....	32
4.2.1. Bacterial Strains	32
4.2.2. Phytochemicals and antibiotics.....	32

Master Thesis

4.2.3. Determination of <i>Zeta</i> Potential.....	32
4.2.4. Physico-chemical charecterization of bacterial surface	33
4.2.5. Free energy of adhesion	34
4.2.6. Motility	34
4.2.7. Detection of quorum-sensing inhibition	35
4.2.8. Outer Membrane Proteins	36
4.2.8.1. Extraction	36
4.2.8.2. OMP Analysis	36
4.2.9. Antibiotic-Phytochemical Dual Combinations Assay – Efflux Pumps Inhibition.....	37
4.2.9.1. Classification of dual combinations	37
4.2.10. Statistical analysis	38
4.3 Results and Discussion	39
4.3.1. Surface characterization of cells in presence of phytochemicals	39
4.3.2. Analysis of <i>E. coli</i> and <i>S. aureus</i> free energy of adhesion.....	41
4.3.3. Motility assays	43
4.3.4. Quorum-sensing assays	45
4.3.5. Characterization of cell membranes	48
4.3.5.1. Outer Membrane Proteins.....	48
4.3.5.1. Efflux pumps	50
Chapter 5	53
Concluding remarks and perspectives for further research	53
5.1. Conclusions	53
5.2. Perspectives for further research.....	55
References	57
Appendix	I
A. Experimental data.....	I

Figures List

Figure 1. Traditional targets of antimicrobial compounds.....	7
Figure 2. Model of biofilm development. Planktonic cells contact with surface and cells resulting in the formation of microcolonies. Cells in the biofilm can return to a planktonic lifestyle to complete the cycle of biofilm development.....	14
Figure 3. OD _{570 nm} as a measure of biofilm mass (a) and fluorescence (λ_{ex} : 570 nm and λ_{em} : 590 nm) results as a measure of biofilm viability (b) for <i>E. coli</i> and <i>S. aureus</i> . The mean \pm standard deviation values resultant from three independent experiments are depicted.....	25
Figure 4. Examples of results obtained with quorum-sensing assay: (a) no bioactivity (SA at 5000 $\mu\text{g/mL}$); (b) antibacterial and anti-QS halos are observed (I3C at 5000 $\mu\text{g/mL}$) with addition of phytochemicals.....	46
Figure 5. OMPs profiles of <i>E. coli</i> . The profile of molecular weight standards (1), control (2), 7-HC (3), I3C (4), SA (5), SP (6) are presented.....	49

Tables List

Table 1. MIC for <i>E. coli</i> and <i>S. aureus</i>	23
Table 2. MBC values for <i>E. coli</i> and <i>S. aureus</i>	24
Table 3. Percentages of biofilm removal and inactivation by the selected phytochemicals against <i>E. coli</i> and <i>S. aureus</i>	26
Table 4. Zeta potential (mV) results of suspensions of <i>E. coli</i> and <i>S. aureus</i> in contact with phytochemicals at the MIC.	39
Table 5. Hydrophobicity (ΔG^{TOT}), and apolar (γ^{LW}) and polar (γ^{AB}) components of the surface tension of untreated and treated cells. The means \pm SDs are illustrated.	40
Table 6. Free energy of adhesion (ΔG^{TOT}_{bws}) of bacterial cells to polystyrene, treated and untreated with phytochemicals.	42
Table 7. Motility results for bacteria with and with phytochemicals. The drop baseline was 6mm which was subtracted from the results presented.	44
Table 9. Antimicrobial activity of antibiotics. The means (mm) \pm standard deviation for at least three replicates are illustrated.	51
Table 10. Classification of the effect of dual combinations of phytochemicals and antibiotics.	52
Table A.1. Physico-chemical characterization of polystyrene (PS).	I

Glossary

Indexes and parameters:

AB – Lewis acid-base component

B – Bacteria

B - α -bromonaphtalene

F – Formamide

LW – Lifshitz-van der Waals component

S – Substratum

TOT – Total

W – Water

sws- Between two entities on a given surface when immersed in water

bws – Between one bacteria and a substratum that are immersed or dissolved in water

bwb – Between two bacterial surfaces, when immersed in water

bw – Between bacteria and substratum

sw – Between substratum and bacteria

TOT – Total

ΔG – Free energy of interaction (mJ/m^2)

γ - Surface free energy (mJ/m^2)

θ - Contact angle ($^\circ$)

+ - Electron acceptor parameter of Lewis acid-base component

- - Electron donor parameter of Lewis acid-base component

Abbreviations:

AHL - N-acyl-homoserines

AI – Autoinducers

CIP – Ciprofloxacin

CV – Crystal Violet

DMSO – Dimethyl sulfoxide

EPS – Extracellular polymeric substances

ERY - Erythromycin

I3C – Indole-3-carbinol

MIC – Minimum Inhibitory Concentration

Master Thesis

MBC – Minimum Bactericidal Concentration

MDR – Multi-drug resistant

OD – Optical Density

OD_C – Cut-off optical density

OMPs – Outer membrane proteins

p – Statistical significance level

PS – Polystyrene

QS – Quorum-sensing

QSI – Quorum-sensing inhibition

RMAs – Resistance-modifying agents

SA – Salicylic Acid

SDS-PAGE – Sodium dodecyl sulfate polyacrylamide gel electrophoresis

SP – Saponin

SPSS - Statistical Package for the Social Sciences

TET - Tetracycline

WHO – World Health Organization

7-HC – 7-hydroxycoumarin

Chapter 1

Work Outline

1.1. Background and Project Presentation

Since the discovery of the first antibiotic, penicillin, the employment of any novel antibiotic has been followed by the appearance of bacterial resistance to that antibiotic in as little time as a few years. Antibiotics have the ability to kill bacteria or inhibit their growth.

Resistance to antibiotics is one of the biggest problems that global public health is facing. Antimicrobial resistance is a natural consequence of the adaption of pathogens to the exposure to antimicrobials used in medicine, food, crop production and to disinfectants in farms and households. Resistant organisms cause infections that are more difficult to treat and more expensive; some strains have become resistant to all available antimicrobial agents (Byarugaba, 2004). Resistant infections affect treatment costs, disease spread and duration of illness (Okeke et al., 2005)

In order to find novel antimicrobial agents with new modes of action, plants have been explored as sources for the identification of new and effective antibacterials. An endless number of plant species have been reported to act against several bacteria *in vitro*, and many medicinal plants produce secondary metabolites (phytochemicals) capable of inhibiting the growth of a wide range of microorganisms including fungi, yeasts and bacteria. Phytochemicals have been studied for the treatment of microbial infections since 1990, due to the increasing inefficacy of conventional antibiotics (Simões et al., 2009).

1.2 Main objectives

The main aim of this work was to assess the antimicrobial efficacy of selected phytochemicals against *Escherichia coli* and *Staphylococcus aureus* planktonic cells and also to evaluate them on biofilm control.

In the present study, four different phytochemicals – 7-hydroxycoumarin (7-HC), indole-3-carbinol (I3C), salicylic acid (SA) and saponin (SP) – were tested against *E. coli* and *S. aureus* in both planktonic and sessile states. The strains tested are considered the most clinical significant bacteria due to their capacity to resist against several antibiotics (Simões et al., 2008; Xu et al., 2006).

To evaluate the antimicrobial activity of the several phytochemicals two experiments were performed to determine the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC). The biofilm control was also performed to understand the efficacy of phytochemicals to remove biomass and inactivate biofilm cells. The biofilm control assay was studied in 24 h aged biofilms after 1 h in contact with the phytochemicals. The biomass removal and metabolic inactivation were calculated through the optical density (OD) and fluorescence measurements.

Several aspects of planktonic cells were evaluated to understand the mode of action of the selected phytochemicals. The surface charge of bacteria was studied through the measurement of Zeta potential and the hydrophobicity of cells was also assessed. To evaluate the potential activity of the phytochemicals to prevent *E. coli* and *S. aureus* adhesion to polystyrene (PS), the prediction of theoretical adhesion

through the measurement of contact angles was performed. The phytochemicals were also studied on the ability to interfere with bacterial motility and quorum-sensing (QS), two microbial aspects involved in biofilm formation.

The OMPs expression of *E. coli* was studied in contact with phytochemicals through a sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) in order to detect the possible expression of resistance proteins. Regarding *S. aureus* resistant strains, dual combinations of phytochemicals and antibiotics – tetracycline (TET), ciprofloxacin (CIP) and erythromycin (ERY) - were tested to understand their ability to act in efflux pumps inhibition.

1.3 Thesis Organization

In chapter 1, the context, motivations and main goals for the development of this thesis are explained. This chapter is also a guideline to the overall work, composed by 5 chapters.

Chapter 2 includes the literature review about the main subjects of this work. In this chapter, the main problems associated with the appearance of bacterial resistance are highlighted. Plant products, especially secondary metabolites, are introduced as one of the solution for the antimicrobial resistance. The mode of action of phytochemicals is also developed in this chapter. Finally, it is reported the problem of the higher resistance associated to bacterial growing in biofilms and quorum-sensing inhibition (QSI) is presented as one possible solution for the prevention of biofilm formation.

In the third chapter are described the results of the activity of phytochemicals as antimicrobial agents against *E. coli* and *S. aureus*. The MIC and MBC of the selected phytochemicals are presented. In this chapter are also studied the phytochemicals in the control of biofilm, showing the biofilm removal and metabolic inactivation for each one of the phytochemicals at the MIC and $5 \times \text{MIC}$.

Chapter 4 describes the study of surface charge and hydrophobicity characteristics of *E. coli* and *S. aureus* when exposed to phytochemicals. The influences of phytochemicals in motility and QS are also presented in this chapter. To finalize the chapter, OMPs expression of *E. coli* are studied when exposed to

phytochemicals; and dual combination of phytochemicals and antibiotics are described in order to study the antimicrobial activity and their synergistic effects in *S. aureus* efflux pump inhibition.

Finally, in chapter 5 the main conclusions of the work are referred and some recommendations for future research are given.

Chapter 2

Literature Review

2.1 Antimicrobial resistance and phytochemicals

There is a continuous search for new drugs and antibiotics in order to heal the main infectious diseases. However, the microorganisms have become resistant to most of the antibiotics. The microorganisms are successful when facing adverse conditions, because they seem to sense and respond to the external environment and modulate gene expression accordingly. Antimicrobial resistance is one of the biggest problems threatening global public health (Byarugaba, 2004; Okeke et al., 2005). This problem is a natural consequence of the adaption of infectious agents to antimicrobials used in several areas, including medicine, food animals, crop production and disinfectants in farms, hospitals and households (Bloomfield, 2002; McEwen and Fedorka-Cray, 2002; Vidaver, 2002; Wise and Soulsby, 2002). Resistance allows microorganisms to survive in the presence of toxic conditions. The effectiveness of many antimicrobial drugs has been lost due to the evolution of pathogen resistance. Many of the microorganisms are no longer susceptible to most of the existing antibiotics and therapeutic agents (Byarugaba, 2004). Bacteria generally

acquire drug resistance in a multitude of ways (Saleem et al., 2010). The acquisition can be done by *de novo* mutation or acquisition of resistance genes from other microorganisms and passed on during replication (Fajardo et al., 2008; McManus, 1997; Sefton, 2002; Smith and Lewin, 1993). Resistance genes are able to act in different ways enabling bacterium to: produce the enzymes that inactivate the antimicrobial agent; modify the target site; produce an alternative metabolic pathway that inhibits the antimicrobial agent; express efflux mechanisms, preventing the antimicrobials to reach intracellular targets (Spratt, 1994; Webber and Piddock, 2003; Woodford and Ellington, 2007).

There are several factors influencing the efficacy of antimicrobial agents. Examples include the use of an inefficient product, in other words, of an antimicrobial product that presents an incomplete spectrum of activity; the application of antimicrobial agents at sublethal concentrations, which can allow the adaptation of microorganisms to these new conditions; or an insufficient contact time between the antimicrobial product and microorganisms (Bessems, 1998; Heinzl, 1998; Russel 2003).

Beyond all the environmental factors affecting the resistance explained before, there are cellular mechanisms influencing this process. The Gram-negative bacteria are less susceptible to antimicrobial agents than Gram-positive bacteria, because they present a thick cell wall, more difficult to entry, and an outer membrane. The waxy envelope presented in mycobacteria inhibits the uptake of antimicrobial agents, so they are even more resistant (McDonnell and Russell, 1999). Efflux is another process related with the increasing of resistance. Through efflux pumps, Gram-negative bacteria pump out the antimicrobial, detergents, organic solvents and disinfectant agents, contributing to the resistance (Beumer et al., 2000; Cloete, 2003; Kumar and Schweizer, 2005).

The resistant microorganisms can cause infections that are more difficult to treat, and it is necessary drugs that are more expensive and toxic and also less available (Howard et al., 2001). The effective microbial therapy is usually delayed because of the acquisition of resistance from microorganisms (Ibrahim et al., 2000; Kollef, et al., 1999; Lautenbach et al., 2001). Bacteria have shown resistance with increasing trends. Consequently, the rate of discovery of new antimicrobial agents has decreased since the 1970s (Byarugaba, 2004). The traditional antibiotics have been recognized because they are able to kill bacteria or inhibit their growth, through

inhibition of bacterial functions, such as: cell wall synthesis, DNA replication, RNA transcription and protein synthesis, which are essential for cell growth (Figure 1) (Clatworthy et al., 2007). In order to treat infections caused by antimicrobial-resistant organisms, it is necessary a more effective therapy (Cosgrove and Carmeli, 2003).




Figure 1. Traditional targets of antimicrobial compounds.

The methods to evaluate antimicrobial activity are essential for the selection of a new drug. However, there are several factors that can affect the antimicrobial susceptibility test: pH, moisture, effects of thymidine or thimine, effects of variation in divalent cations and growth conditions (Lalitha, 2004). Magnesium and calcium are divalent cations that affect the results of aminoglycosides and tetracycline. Finally, the components needed for growth should be fulfilled (Lalitha, 2004). The antimicrobial susceptibility tests are divided into 3 principles: diffusion, dilution and diffusion and dilution. The most popular methods are: the Kirby-Bauer, E-Test, agar dilution, Stokes, microdilution and macrodilution (Anvisa, 2012; Lalitha, 2004; Othman et al., 2011).

2.2. A new therapy against resistance – Phytotherapy

Natural new antimicrobial can be present in several products used by humankind. For example, aromatic plants and spices, used as flavour enhancers, have already been demonstrated to have significant antimicrobial activity (Christaki et al., 2012). Also, antimicrobials from vegetal origin can be obtained from essential oils, seeds, flowers, fruits and roots. These kind of antimicrobial systems have been used

centuries ago, in the preservation of food. Plants produce a wide range of organic compounds. However, the secondary metabolites present special interest because of their importance as pharmaceuticals, venoms, fragrances, industrial materials and cosmetics (Amaral and Silva, 2003; Mendes, 2007). The discovery of all the potential of plants brings a great interest in phytotherapy.

The World Health Organization (WHO) defines medicinal plants those that present traditional use as therapeutic compounds (Severiano et al., 2010). According to the WHO, medicinal plants would be the greatest source to obtain a wide range of drugs with antimicrobial properties (Nascimento et al., 2000). Consequently, interest in medicinal plants has increased in recent years. Indeed, plants were reported to be used by about three quarter of the world population (Rao et al., 2004). The story of medicinal plants says that they are part of the humankind evolution and were the first therapeutic sources used. However, the use of medicinal plants, phytotherapy, had major impact in the beginning of XIX century. There are 250.000-500.000 plant species, however just a small percentage has been investigated phytochemically (Mahesh and Satish, 2008).

It is interesting to realize that most of the medicines used nowadays were identified in plants, and some of the most important pharmaceutical drugs are natural drugs and linked directly to natural product research (Bighetti et al., 2005). Also, most of the commercial drugs were used before in crude form in traditional and folk medicine, suggesting their potential biological activity (Ciocan and Bara, 2007). Also, plants present numerous constituents and their extracts can show synergistic effects between the different active principles due to the presence of classes of compounds or different structures contributing for the same activity (Maciel et al., 2002). The plants can prevent cardiovascular diseases, carcinogenesis, inflammation, atherosclerosis, and others health problems (Albano and Miguel, 2001).

Some plants have potential as remedies for diseases. Examples include the use of bear-berry (*Arctostaphylos uvaursi*) and cranberry juice (*Vaccinium macrocarpon*) to treat urinary tract infections, essential oils of tea tree (*Melaleuca alternifolia*) as active ingredients in many topical formulations to treat cutaneous infections and *Hydrastis canadensis* and *Echinacea* species for tuberculosis infections (Abreu et al., 2012).

Regarding the plants, they can be divided in several families, for example Asteraceae, Lamiaceae, Apeaceae, Euphorbiaceae, Rutaceae and Fabaceae. Regarding

to these families, some of them are more incident in Portugal. Lamiaceae is one of the families found in our country and frequently found in Mediterranean region, Middle East and tropical mountains (Severiano et al., 2010). Currently, there are 3.500 species of this family (Severiano et al., 2010). Lamiaceae are composed of herbs, shrubs and trees. Plants belonging to this family can be applied in medicine, cosmetics and food industry (Mendes, 2007). Asteraceae is another plant family present in Portugal and includes around 23.000 species and 1.500 genus. They can be found in temperate and subtropical climates. Herbs, subshrubs and vines belong to this family (Severiano et al., 2010). Finally, Apiaceae is another family present in Portugal and is composed for 3.000 species and 300 genus (Moreno-Dorado et al., 2000). The plants that belong to this family can be found in tropical or temperate regions (Judd et al., 2002). Such as Lamiaceae, plants of this family are commonly used in medicine and food industry.

2.3. Phytochemicals and their classes

The increasing of antimicrobial resistance has led to the study of plants products for searching new antimicrobials (Clardy et al., 2006). The chemical diversity and structure are the main causes for the success of phytochemicals (Gibbons, 2004). The phytochemicals have become interesting since the 1990's because of the increasing dissemination of bacterial resistance mechanisms worldwide, due to the excessive and inappropriate use of antimicrobials.

Plant products are divided into two classes: phytoanticipins and phytoalexins. Molecules that are present in an inactive form (example: glucosides) belong to the first group; the second group is composed by molecules whose levels increase in response to microbial attack or is generated in response to a specific infection (Tegos et al., 2002). Phytoalexins are a large group of chemically diverse molecules, including: simple phenylpropanoids derivatives, alkaloids, glycosteroids, flavonoids, isoflavonoids, sulfur products, terpenes and polyketides (Hammerschmidt, 1999). The same molecule can be a phytoalexin or a phytoanticipin in different organs of the same plant. Examples of phytoanticipin are terpenoids, quinones and tannins (Abreu et al., 2012).

There are several phytochemical classes with antimicrobial properties; however, the medical community does not recognize them as therapeutic agents. This is mainly explained because the majority of phytochemicals have weak spectra of activities (Tegos et al., 2002) and the concentrations required are too high to be clinically significant (Aeschlimann et al., 1999; Schmitz et al., 1998). The major problem for the identification of antimicrobial agents from plants is the variability in the extraction methods and antibacterial tests used (Simões et al., 2009).

Beyond the antimicrobial properties, phytochemicals are able to present antiviral (Jassim and Naji, 2003; Muhtar et al., 2008; Naithani et al., 2008), antiparasitic (Atawodi and Alafiatayo, 2007; Chan-Bacab and Peña-Rodríguez, 2001; Sriram et al., 2004) and antifungal (Morel et al., 2002; Rahman and Moon, 2007; Treyvaud Amiguet et al., 2006) activities and also exert cytotoxic activity against tumor cells (Rimando and Suh, 2008; Suffredini et al., 2006; Udenigwe et al., 2008).

The groups of secondary metabolites produced by plant include: terpenoids, phenolics, alkaloids, essential oils constituents, lectins polypeptides and polyacetylenes (Mendes, 2007; Simões et al., 2009). The secondary compounds are responsible for defending against external aggressions and the main subclasses are: simple phenols and phenolic acids, quinines, flavones, flavonoids, tannins, coumarins, and others (Simões et al., 2009). Usually, the plants belonging to one family produce the same type of compounds. Normally, alkaloids are produced from the Apocinaceae and Solanaceae families; flavonoids are produced from Rutaceae plants; and terpenoids are produced from Asteraceae and Lamaceae families (Amaral and Silva, 2003). According to the classes that substances belong, they can produce different effects. Flavonoids present anti-inflammatory action, protecting the blood vessels, and are hypotensive and sedative; digestion action is guaranteed by antraquinones; bronchodilator action is made by coumarins; bactericidal action is achieved with tannins and essential oils (Severiano et al., 2010); and essential oils presents, apart from the others already mentioned, sedative, stimulating, analgesic and expectorant properties (Souza et al., 2011). Antimicrobial properties are expressed by most of plants that are composed by: tannins, terpenoids, glycosides, alkaloids, saponins, flavonoids, polyphenols and coumarins (Das et al., 2010; Hill, 1952; Neelima et al., 2011; Padmini et al., 2010).

The phytochemical compounds are mostly secondary metabolites of plants and deposit in specific parts of the plant (Ciocan and Bara, 2007). Regarding phytochemicals, the main classes with health benefits are present below.

Phenolics and polyphenols form one of the simplest groups of bioactive phytochemicals, consisting of a single substituted phenolic ring (Das et al., 2010). This group seems to be toxic to microorganisms because of the site(s) and number of hydroxyl groups present on the phenolic ring (Scalbert et al., 1991; Urs et al., 1975). Increasing the hydroxylation results in the increasing of toxicity (Ciocan and Bara, 2007). Quinones are composed by an aromatic ring with two ketones substitutions (Das et al., 2010). These molecules react with nucleophilic amino acids in proteins, causing the inactivation of the protein or loss of function (Ciocan and Bara, 2007). Flavones, flavonoids and flavonols belong to a group of molecules with phenolic structures containing one carbonyl group (Ciocan and Bara, 2007; Das et al., 2010). They are active against a wide range of microorganisms, probably because they can react with extracellular and soluble proteins and also react with cell walls (Bennet and Wallsgrove, 1994; Ciocan and Bara, 2007). Tannins are another group that can be divided into two groups: hydrolysable and condensed. The first is based on gallic acid; the second group is composed by numerous condensed tannins derived from flavonoid monomers. The antimicrobial mode of action is related with their ability to inactivate microbial adhesin, enzymes and cell envelope transport proteins (Ciocan and Bara, 2007). Coumarins are derivatives of cinnamic acid and comprise a large class of phenolic substances (Hoult and Payát, 1996; Thuong et al., 2009). These compounds are the simplest members of the group of oxygen heterocyclic, also known as 1,2-benzopyrene, consisting of fused benzene and α -pyrone ring (Hoult and Payát, 1996). More than 1300 coumarins have been studied from natural sources and several properties are related with them, such as: antimicrobial, anti-inflammatory, antioxidant, anticoagulation, antiestrogenic and sedative (Hoult and Payát, 2012; Mello, 2009; Paramjeet et al., 2012). Essential oils and terpenoids are compounds based on an isoprene structure. They occur as diterpenes, triterpenes and tetraterpenes (C_{20} , C_{30} , C_{40}), they are called terpenoids when contain other elements, such as oxygen. These compounds are active against bacteria, fungi, viruses and protozoa (Ciocan and Bara, 2007; Das et al., 2010).

The alkaloids are heterocyclic nitrogenous compounds that present analgesic, antispasmodic and bactericidal action (Ciocan and Bara, 2007; Das et al., 2010; Stary, 1996).

2.4. Mode of Action of Phytochemicals

Bacterial growth can be inhibited by phytochemicals through several mechanisms. These plant products can act on various biochemical targets on the bacterial cells. The mode of action of phytochemicals is not completely understood, neither the phytochemical antibacterial specificity (Simões et al., 2009).

Some experiments have been done to study the mode of action of several phytochemicals. The chemical structure and properties influence the site of action of phytochemicals. The mechanism of action of essential oils against bacteria involves membrane disruption through the lipophilic structure (Griffin et al., 1999; Mendoza et al., 1997). Alkaloids, such as berberine and piperine, interact with bacterial cytoplasmic membrane, intercalate with DNA or inhibit efflux pumps in *S. aureus* (Khan et al., 2006). Phenols act by interruption of energy production due to enzyme inhibition by the oxidized products, which react with sulfhydryl groups or non-specific interaction with proteins (Mason and Wasserman, 1987). In the case of flavonoids, they inhibit the synthesis of nucleic acids of Gram-negative and Gram-positive bacteria (Cushnie and Lamb, 2005; Mori et al., 1987). Other authors, shown that glycoside saponins are able to induce pore-like structures that change the membrane permeability; they can also interfere with energy metabolism (Mandal et al., 2005; Melzig et al., 2001; Sinha Babu et al., 1997).

2.5. Biofilms and phytochemicals:

Natural products have been isolated from plants for usage in biodeterioration control. Biodeterioration is the chemical and physical alteration resulting from

biological activity. The microorganisms associated to biodeterioration grow as biofilms that adhere to substrates. The treatment for this problem involved the use of biocides, however these are chemical agents and most of them are cytotoxic. So, natural compounds from plants with biocidal activity have emerged, and they are promising alternative for the control of biodeterioration without negative impacts of the environment (Guiamet et al., 2006). The chemical biocides are toxic and difficult to degrade, being persistent in the environment, causing chemical contaminations and the spread of resistance. In contrast, natural biocides, extracted from plants, are biodegradable and environmental friendly (Guiamet and Saravia, 2005). The phytochemicals can act as control agents on the bacterial biofilm formation and development (Simões et al., 2009).

Guiamet and co-workers (2006) performed a study with *Cichorium intybus*, *Arctium lappa* and *Centaurea cyanus* from Asteraceae family to test their ability to be used as antimicrobial agents against different microorganisms associated with biodeterioration. These plants showed moderate activity against two species of *Pseudomonas* and no activity against *Bacillus cereus*. *Rosmarinus officinalis* L. extracts act as an antimicrobial agent against *S. aureus*. *Allium sativum* produce allicin, which is one of the most effective antimicrobial products isolated from garlic (Abreu et al., 2012).

Biofilm formation is a feature closely related to pathogenicity (Ren et al., 2005). A biofilm is formed by planktonic bacteria that adhere to a surface and initiate the development of sessile microcolonies surrounded by an extracellular matrix (Otto, 2009) (Figure 2). Bacteria form complex surface-attached communities, also called biofilms (Hentzer and Givskov, 2003). Biofilms develop structures that are morphologically and physiologically differentiated from free-living bacteria (Davies et al., 1998).

The process of biofilm formation includes several steps: preconditioning of the adhesion surface; planktonic cells are transferred from the bulk liquid to the surface; adsorption of cells at the surface; desorption of reversible adsorbed cells; the bacterial cells are adsorbed irreversibly at the surface; transport of substrates to the biofilm; substrate metabolism by the biofilm cells and transport of products out of the biofilm; finally, biofilm is removed by detachment or sloughing (Simões, 2005). The formation of biofilm is dependent of several parameters. Regarding to the surfaces, the attachment is easier on rough, hydrophobic and coated surfaces (Donlan, 2002;

Pereira, 2001). Parameters like flow velocity, water temperature and nutrient concentration also influence the biofilm attachment (Pereira, 2001; Vieira, 1995).




Figure 2. Model of biofilm development. Planktonic cells contact with surface resulting in the formation of microcolonies. Cells in the biofilm can return to a planktonic lifestyle to complete the cycle of biofilm development.

Biofilms are an example of physiological modifications and they also increase the tolerance to antimicrobial therapies and to the host immune response (Hentzer and Givskov, 2003; Simões et al., 2009). Most of the bacterial infections detected in human body involve the formation of biofilms. The biofilm mode of growth permits an increased bacterial survival in hostile conditions, such as in the presence of antibiotics and disinfectants (Hentzer and Givskov, 2003; Trentin et al., 2011). There is an increasing interest in preventing, controlling and eradicating biofilms. Biofilms by bacterial cells are thought to be regulated by autoinducer molecules; in a process called quorum-sensing (Ren et al., 2002). The impairment of bacterial adhesion and biofilm formation by a pathway that does not affect the bacterial death is an important feature of the new concept in antivirulence therapies (Trentin et al., 2011). This alternative should maintain the cells in a planktonic state, switching off the virulence expression and attenuating the pathogen, making the microorganisms more susceptible to antimicrobial agents and immune system (Macedo and Abraham, 2009; Martin et al., 2008).

Bacteria in biofilms present a reduced susceptibility to antimicrobial agents caused by a variety of factors, such as: nutrient depletion within the biofilm, reduced access to cells in the biofilm, production of degradative enzymes and neutralizing

chemicals, between others (Brown and Gilbert, 1993). Biofilms have been reported to be 100-1000 more protectors to bacteria than populations of planktonic cells (Gilbert et al., 2002; Mah and O'Toole, 2001; Stewart and Costerton, 2001). The main difference between planktonic cells and biofilms is the presence of a polysaccharide matrix, delaying the diffusion of antimicrobials into the biofilm (Brooun et al., 2000).

Traditional treatment of infectious diseases is related with compounds that inhibit the growth of bacteria. But, it has been observed that they develop resistance to antimicrobial compounds. So, quorum-sensing seems to be the next opportunity to improve bacterial infection. Quorum-sensing influences bacterial biofilm growth and biofilm development that is related with cell-cell interactions (Simões et al., 2009). Quorum-sensing inhibitory compounds are the new line of antimicrobial agents and can be applied in several areas: medicine, agriculture and aquaculture (Hentzer and Givskov, 2003).

Several biotechnology companies have already tried to develop some strategies to interrupt the bacterial quorum-sensing, such as: inhibition of N-acyl-homoserines (AHL) signal recognition, signal dissemination and signal reception (Hentzer and Givskov, 2003). By interfering with cells communication, it is possible to interfere also in the resistance of biofilms and their ability to form resistant structures, causing cell dispersion. Quorum-sensing systems are involved in a wide range of microbial activities: extracellular enzyme biosynthesis, biofilm development, antibiotic biosynthesis, biosurfactant production, extracellular polymeric substances (EPS) synthesis and production of extracellular virulence factors (Chatterjee et al., 1995; Davies et al., 1998; Daniels et al., 2004; Fux et al., 2005; McGowan et al., 1995; Passador et al., 1993; Pearson et al., 1995; von Bodman and Farrand, 1995).

Autoinducers (AI) are molecules to perceive the size of bacterial population and AHL are the major AI molecules. Several Gram-negative bacteria use AHL signals to coordinate the behaviour of cells in a population. An important achievement was the discovery of molecules produced by plants that mimic AHL signals, affecting quorum-sensing behavior. Several plants of medicinal use demonstrated potential to inhibit quorum-sensing (Hentzer and Givskov, 2003; Waters and Bassler, 2005).

Studying the grapefruit and its furanocoumarins as inhibitors of biofilm formation, it has been shown that dihydroxybergamottin and bergamottin exhibit strong inhibition of both AI-1 and AI-2 activities even at concentrations as low as 1

$\mu\text{g/mL}$ (Girenavar et al., 2008). Furanocoumarins are able to interfere with cell-cell signalling and also inhibit biofilm formation. However, the mechanisms of action are not completely understood (Girenavar et al., 2008). *Delissea pulchra*, an Australian macroalga, produces halogenated furanones, which are inhibitors of AHL, inhibiting bacterial quorum-sensing and biofilm formation (McLean et al., 2004). Auraptene and lacinartin, two compounds belonging to coumarin family, are promising natural compounds that can be used to prevent and treat periodontal diseases. They were also evaluated on the growth, biofilm formation/desorption, and adherence to human oral epithelial cells of *Porphyromonas gingivalis*. Lacinartin was able to inhibit biofilm formation and to cause desorption of a pre-formed biofilm of *P. gingivalis*. This suggests that coumarins may contribute to reducing tissue destruction (Marquis et al., 2012). Some information about the structure-activity relationship of coumarins showed that the group on position 5 and position 2'/3' of the isoamylene chain can affect the antifouling activities against both *Balanus albicostatus* and *Bugula neritina* (Wang et al., 2013).

Therefore, phytochemicals are not only important for the antimicrobial response and the substitution of antibiotics because of bacterial resistance, but also for the control of biofilms formation. Natural products are important sources of bioactive compounds and the medicinal plants used in folk medicine can facilitate the search of new agents.

Chapter 3

Activity of Selected Phytochemical Products as Antimicrobials and in Biofilm Control

3.1 Introduction

Microbiologists have learned to assess antibiotic effects *in vivo* by evaluating the MIC and MBC *in vitro*. These methods assess the influence of antibiotics against planktonic microorganisms in the exponential phase of growth and predict the efficacy of antibiotics against bacteria in infections (Ceri et al., 1999; Fux et al., 2005).

Staphylococcus aureus and *Escherichia coli* are two human pathogens that can cause a variety of infections. The main characteristic of these infections is the formation of biofilms (Beenken et al., 2004).

In order to prevent biofilm formation, several studies have been performed to find antimicrobial agents that affect the viability of bacteria in biofilms. Natural products from plants have been shown to influence microbial biofilm formation (Rasooli et al., 2008). Plants are an important source of disinfection compounds as they produce a wide range of phytochemicals with antimicrobial properties, most of them against microorganisms, insects, nematodes and other plants (Abreu et al., 2013a). Phytochemicals are able to inhibit peptidoglycan synthesis, damage microbial

membrane structures, modify bacterial membrane surface hydrophobicity and also modulate quorum-sensing (Rasooli et al., 2008). These processes can also inhibit or affect biofilm formation.

To reach high killing rates and to avoid resistance and adaption, disinfectants are used at very high concentrations relative to their MICs (Abreu et al., 2013a). So, after the detection of MIC and MBC for the selected phytochemicals, they were tested at MIC and 5×MIC. The effects of phytochemicals on *E. coli* and *S. aureus* biofilms were evaluated through biomass prevention and metabolic inactivation. The quantification of biofilm removal takes into account both live and dead cells assessed by the Crystal Violet method (CV). CV is a dye which binds to negatively charged surfaces in the extracellular matrix (Extremina et al., 2011; Peeters et al., 2008). Resazurin is used to quantify the viability of cells, based on the live cells. This compound is a blue redox indicator and it reduces to pink by contact with viable bacteria in the biofilm (Extremina et al., 2011).

Pathogens are increasing their capacity to survive after contact with antimicrobials and disinfectants. Many of the existing antibiotics are ineffective due to their extensive and inappropriate use (Abreu et al., 2013a). Plants are important sources for the development of antimicrobials and strategies to control growth and biofilm formation (Abreu et al., 2013a).

The purpose of this study was to assess the antimicrobial efficacy of selected phytochemicals (7-HC, I3C, SA and SP) against *E. coli* and *S. aureus* planktonic cells. Moreover, the effects of these phytochemicals were assessed on biofilm control.

3.2 Material and methods

3.2.1. Bacterial Strains

The bacteria used in this study were obtained from the Spanish Type Culture Collection (CECT): the Gram-negative bacterium *Escherichia coli* (CECT 434) and the Gram-positive bacterium *Staphylococcus aureus* (CECT 976). The bacteria were distributed over the surface of Plate Count Agar (PCA – Merck) and incubated for 24 h at 30 ± 3 °C.

3.2.2. Phytochemicals

The phytochemicals used were: 7-hydroxicoumarin (7-HC), indol-3-carbinol (I3C), salicylic acid (SA) and saponin (SP). These compounds were obtained from Sigma-Aldrich (Portugal) and prepared in dimethyl sulfoxide (DMSO, Sigma).

3.2.3. Determination of Minimum Inhibitory Concentration

The MIC is considered the lowest concentration of an antimicrobial that will maintain or reduce the growth of a microorganism after 24 h incubation (Andrews, 2001). MIC of phytochemicals was determined by microdilution method in sterile 96-wells microtiter plates (McBain et al., 2004). The cell suspensions of *S. aureus* and *E. coli* were obtained by overnight cell cultures in Mueller-Hinton broth (MHB) (Fluka, Portugal) and were adjusted to a OD_{620nm} at 0.1 ± 0.02 (corresponding to approximately 1×10^6 cells/mL) in the spectrophotometer (VWR V-1200). The suspension cells were added to sterile 96-well polystyrene microtiter plates (Orange Scientific) with different phytochemicals in several concentrations (25, 50, 100, 200, 400, 800, 1600, 3200 $\mu\text{g/mL}$) in a final volume of 200 μL . The volume of 7-HC, I3C, SA and SP added to each well was 10 μL . DMSO was used as a negative controls. No antimicrobial activity was detected by DMSO at this concentration (data not shown). After 24 h at 30 °C, the MIC of each sample was determined by measuring the optical density in the spectrophotometer (620 nm) (SpectraMax M2E, Molecular devices). MIC corresponds to the concentration in which the final OD is inferior or equal to the initial OD.

3.2.4. Determination of Minimum Bactericidal Concentration

The MBC is defined as the lowest concentration of antimicrobial that will prevent the growth of an organism after subculture on to antibiotic-free media (Andrews, 2001). MBC of phytochemicals was determined by the drop method. After measuring the MIC, the wells corresponding to the phytochemicals concentrations equal and above the MIC were added (10 μ L) to PCA plates. The drops were drained along the plate. After 24 h at 30 °C, the plates were analysed and the MBC of each phytochemical corresponding to the concentration which inhibited the growth of the bacteria.

3.2.5. Biofilm formation and control in sterile 96-well polystyrene microtiter plates

Biofilms were developed according to the modified microtiter plate test proposed by Stepanović et al. (2000). For both bacteria, at least 6 wells of a 96-well polystyrene microtiter plate were filled with 200 μ L of overnight batch cultures in MHB ($OD_{620nm} = 0.04 \pm 0.02$). The plates were incubated overnight at 30 °C and 150 rpm. The negative control wells were also placed on the plates, being sterile water and medium. Plates were incubated for 24 h at 30 °C and agitated at 150 rpm. In order to test the effects of phytochemicals in several steps of the process there are various treatments that can be applied. However, in this case, the treatment applied consisted in incubating overnight the strains without phytochemicals. The biofilms were formed in microtiter plates for 24 h, and subsequently, were incubated with phytochemicals for 1 h.

After biofilm development, the content of wells was removed and the wells were washed three times with 200 μ L of NaCl (8.5 g/L) to remove reversibly adherent bacteria. The phytochemicals were added to the wells at the MIC and $5 \times$ MIC. The microtiter plates were incubated for 1 hour. The remaining attached bacteria were analysed by using crystal violet and resazurin.

3.2.6. Biofilm analysis

3.2.6.1 Crystal Violet method

Before phytochemicals application, the inoculum in the wells was removed and the wells were washed with 200 μL of sterile water. Later, 250 μL of ethanol were loaded for 15 minute to promote biofilm fixation. The supernatant was removed and the plates were air-dried. Subsequently, 200 μL of CV solution (Gram color staining set for microscopy, Merck) was added for 10 minutes to stain the fixed bacteria. After washing in water, the plates were dried and finally, the wells were loaded with 200 μL of aceditic acid 33% (v/v) (Merck) to release and dissolve the stain. To analyse the biofilm, the OD of the solutions was measured at 570 nm using a microtiter plates reader (SpectraMax M2E, Molecular Devices). After obtaining the values of absorbance, the percentage of biomass removal is obtained according equation 1:

$$\% \text{ Biomass removal} = \frac{OD_{control_{570}} - OD_{phytochemical_{570}}}{OD_{control_{570}}} \times 100 \quad (1)$$

where $OD_{control_{570}}$ represents the optical density of the control at 570 nm, and $OD_{phytochemical_{570}}$ is the optical density of the phytochemical.

3.2.6.2 Resazurin Method

In this method, a commercially available resazurin solution (Sigm) was used. The plates were loaded with 190 μL of sterile MH medium and 10 μL of resazurin solution. After 20 minutes of incubation at room temperature, fluorescence (λ_{ex} : 570 nm and λ_{em} : 590 nm) was measured using the microtiter plates reader. After measuring the fluorescence, it is possible to calculate the percentage of metabolic inactivation:

$$\% \text{ Metabolic inactivation} = \frac{FLUO_{control} - FLUO_{phytochemical}}{FLUO_{control}} \times 100 \quad (2)$$

where $FLUO_{control}$ represents the fluorescence intensity of biofilms not exposed to phytochemicals and $FLUO_{phytochemical}$ represents the fluorescence intensity value for biofilms exposed to phytochemicals.

3.2.7. Classification of biofilm producer bacteria

According to Stepanović et al. (2000), the bacterial strains can be classified using the following groups:

- Non-producer (-): $OD \leq OD_C$;
- Weak producer (+): $OD_C < OD \leq 2 \times OD_C$;
- Moderate producer (++) : $2 \times OD_C < OD \leq 4 \times OD_C$;
- Strong producer (+++) : $4 \times OD_C < OD$.

where OD_C represents the cut-off OD for the microtiter plates test as three standard deviations above the mean OD of the negative control. The negative control wells contained broth only.

3.2.8. Statistical analysis

The data was analyzed using One-Way Anova and the statistical program SPSS 21.0 (Statistical Package for the Social Sciences). The results were presented as the means \pm standard deviation. Significance level for the differences was set at $p < 0.05$ and the calculations were based on confidence level equal or higher than 95%.

3.3 Results and Discussion

3.3.1. Antimicrobial activity of phytochemicals and biofilm control potential

The MIC of 7-OH, I3C, SA and SP were measured for both bacteria (*E. coli* and *S. aureus*). The MIC value was considered to be the lowest concentration of phytochemical able to totally inhibit microbial growth. Table 1 shows the results obtained.

Table 1. MIC for *E. coli* and *S. aureus*.

Strains	Phytochemicals ($\mu\text{g/mL}$)			
	7-HC	I3C	SA	SP
<i>E. coli</i>	800	800	3200	ND
<i>S. aureus</i>	200	400	1600	ND

ND – Not detectable

The values of MIC for *E. coli* are higher than those for *S. aureus*. This result can be explained because *E. coli* is a Gram-negative bacterium and it is less susceptible to antimicrobials than Gram-positive bacteria. The Gram-negative bacteria tend to be more resistant to lipophilic and amphiphilic inhibitors than those Gram-positive, including dyes, detergents, free fatty acids, antibiotics and chemotherapeutics agents (Nikaido, 1996). In other study, antibiotics from natural origin showed activity against Gram-positive bacteria, but more than 90% of them have no useful effect against *E. coli* (Vaara, 1993). This increased resistance of Gram-negative bacteria can be attributed to the presence of the outer membrane. The porin channels slow down the penetration of small hydrophilic solutes and the low fluidity of the lipopolysaccharide layer decreases the rate of transmembrane diffusion of lipophilic solutes (Nikaido and Vaara, 1985; Plésiat and Nikaido, 1992).

Regarding to the different phytochemicals tested, SP was the unique compound that had no detectable MIC for concentrations lower than 3200 $\mu\text{g/mL}$. The 7-HC and I3C were the most effective compounds against both bacteria, since they presented the lowest values of MIC.

The literature is full of reports describing natural products and extract with antimicrobial activity with MIC values over 1000 $\mu\text{g/mL}$, which has little relevance for clinical application (Gibbons, 2004).

After the determination of MIC values for each phytochemical, MBC values were measured, using several concentrations sub and above-MIC. Table 2 presents the values of MBC for both bacteria. Regarding SP, the MBC was not possible to be quantified due to the absence of a MIC value.

Table 2. MBC values for *E. coli* and *S. aureus*.

Phytochemical	Concentration ($\mu\text{g/mL}$)	
	<i>E. coli</i>	<i>S. aureus</i>
7-HC	ND	ND
I3C	1600	800
SA	5000	3200

ND – Not detectable

Similarly to the results of MIC, the values of MBC were higher for *E. coli* than for *S. aureus*. MBC represents the lowest concentration of antimicrobial product necessary to kill a bacterium; so Gram-negative need a higher concentration of antimicrobial, because these bacteria are more resistant (Nikaido, 1996). Regarding to 7-HC, the MBC was not defined for both bacteria, until the maximum value tested (10000 $\mu\text{g/mL}$). I3C seems to be the most effective phytochemical against both bacteria.

After testing the effects of the selected phytochemicals against the *E. coli* and *S. aureus* strains, the effects of the same phytochemicals on their biofilms were tested. Biofilm formation was performed in sterile 96-well polystyrene microtiter plates.

In this case, the bacteria were incubated overnight without phytochemicals. Biofilms were formed for 24 hours and after that, biofilms were incubated with phytochemicals at the MIC and $5 \times \text{MIC}$ for 1 hour. According this procedure, it would be possible to conclude about the efficacy of phytochemicals in control (removal and inactivation) of biofilms. Figure 3 shows the results obtained from this assay.




Figure 3. OD_{570 nm} as a measure of biofilm mass (a) and fluorescence (λ_{ex} : 570 nm and λ_{em} : 590 nm) results as a measure of biofilm viability (b) for *E. coli* and *S. aureus*. The mean \pm standard deviation values resultant from three independent experiments are depicted.

According to the classification proposed by Stepanović et al. (2000), both strains are weak biofilm producers (+).

In the Table 3 are presented the percentages of biofilm removal and inactivation with the selected phytochemicals at different concentrations. The ability of phytochemicals to control 24 h aged biofilms was analysed, based on their effects on biomass and metabolic activity.

Table 3. Percentages of biofilm removal and inactivation by the selected phytochemicals against *E. coli* and *S. aureus*.

Phytochemicals	<i>E. coli</i>		<i>S. aureus</i>	
	% Biomass removal	% Metabolic inactivation	% Biomass removal	% Metabolic inactivation
7-OH (MIC)	51.03±2.53	73.01±1.48	9.63±2.30	39.79±4.15
7-OH (5 × MIC)	46.67±4.47	75.50±0.70	1.68±0.32	47.30±4.59
I3C (MIC)	43.63±6.47	74.67±4.45	0.14±0.04	42.03±5.61
I3C (5 × MIC)	43.98±7.95	74.68±5.78	4.02±0.09	44.99±4.68
SA (MIC)	33.36±5.15	74.24±4.86	0.00±0.00	42.21±5.34
SA (5 × MIC)	34.19±6.52	78.75±4.57	0.00±0.00	48.26±8.55
SP (MIC)	31.80±6.92	73.86±4.62	0.00±0.00	44.71±7.65
SP (5 × MIC)	23.86±5.80	75.09±1.37	0.00±0.00	43.54±0.29

The results show the capacity of phytochemicals to remove/inactivate the biofilms of *E. coli* and *S. aureus*. The biomass removal is dependent on the phytochemical and on its concentration. Comparing the values obtained, it is concluded that the phytochemical concentration (MIC and 5 × MIC) did not influence the removal of the biofilm ($p>0.05$), for both bacteria. Probably, the cells are able to adapt to the phytochemicals and with a higher concentration, the effect is not so noticed. Other possibility is the occurrence of changes in the biofilm phenotype (Cerca et al., 2006). However, the type of bacteria influences the values of biofilm removal. The same phytochemical, at the same concentration, presents a different behaviour in Gram-negative and Gram-positive bacteria ($p<0.05$). The percentages of biomass removal were always higher for *E. coli* than *S. aureus* with all the phytochemicals and concentrations performed. Total biofilm removal was not achieved with any of the selected phytochemicals. The highest reduction in biomass was found for *E. coli* with 7-HC. All the phytochemicals were more active in removing *E. coli* biofilms than *S. aureus* biofilms.

In terms of metabolic activity, the phytochemicals promoted higher reduction for *E. coli* biofilms than *S. aureus* biofilms. The inactivation percentages refer to the quantity of cells not viable present in the biofilm. The selected phytochemicals induced a higher quantity of cells not viable in biofilms of *E. coli* comparing to those of *S. aureus*. The biofilms of *E. coli* treated with phytochemicals at different concentrations did not show significative differences in the percentage of viable cells ($p>0.05$). Probably, the same effect explained before, for the percentage of removal, occurred in this case and the cells are able to adapt or the biofilm phenotype is

changed. However, in the case of *S. aureus*, the biofilms treated with 7-HC and I3C, at different concentrations, were significantly different ($p < 0.05$). In these cases, *S. aureus* probably did not develop any mechanism to protect the cells with the lowest concentration, and when it was applied the highest concentration, the cells were inactivated, and the percentage of nonviable cells increased significantly. The inactivation was more significant in the case of *E. coli* biofilm than for *S. aureus* biofilms with all the concentrations performed ($p < 0.05$).

Phytochemicals had greater effect in *E. coli* than *S. aureus*. This result is contradictory with the results obtained for MIC and MBC. Gram-negative bacteria should be more resistant to phytochemicals and the removal and inactivation should be higher for *S. aureus* than *E. coli*. According Simões et al. (2008), *E. coli* is less susceptible to antimicrobials because of its outer membrane, which constitutes a barrier for the entrance of molecules. However, the number of resistance mechanisms in biofilms increase significantly and one cannot infer on the biofilm susceptibility based on the results of planktonic cells. Furukawa and co-workers (2010), studied the control of *S. aureus* and *E. coli* biofilms on stainless steel, concluding that the last ones are more susceptible to cleaning by five types of cleaning agents, food additives and other compounds on stainless steel. The morphology of the tested biofilms is known to be different. *S. aureus* biofilms are denser than that of *E. coli* biofilm (Furukawa et al., 2010).

Biofilm eradication is a difficult aspect to be solved due to their inherent resistance to biocides. There are several mechanisms explaining the biocides resistance of biofilms: penetration of chemical agents is delayed in the biofilm matrix; the reaction between the biocides and matrix components; the altered growth rate of biofilm organisms and physiological changes due to the biofilm mode of growth (Sillankorva et al., 2004). So, it is difficult to predict the behaviour of the cells presents in biofilms, because there are several components affecting their resistance. The biofilms also present a different mode of growth from planktonic cells: biofilms individual bacteria are enclosed in a matrix of extracellular polymeric substances; bacteria are not distributed uniformly throughout a biofilm; the cell growth can be lower because of the limited access to nutrients and oxygen (Sillankorva et al., 2004).

Chapter 4

Aspects Underlying the Antibacterial and Biofilm Control Action of Phytochemicals

4.1 Introduction

The use of dual combinations of antimicrobial drugs with positive *in vitro* interactions has become an important parameter to evaluate in clinical applications. The appearance of this practice starts to prevent the emergence and widespread multi-drug resistant (MDR) infections (Sopirala et al., 2010). Several antibiotics have been analysed for their action as resistance-modifying agents (RMAs), i.e. compounds able to modify or inhibit the bacterial resistance, so the antibiotics kill efficiently the resistant bacteria (Abreu et al., 2013b).

The clinical effects and synergism of dual combinations have to be tested *in vivo*. Determination of synergy *in vitro* might not be reflected *in vivo*, due to the achievement of synergic levels of drugs in the tissues, differences in plasma protein binding and drug metabolism (Kalan and Wright, 2011). Dual combinations of phytochemicals and antibiotics were tested to understand their ability to act in efflux pumps of *S. aureus*.

The elucidation of the molecular details of drug resistance is a crucial area of research that crosses many subjects since an understanding of the mechanisms by which drug resistance develops leads to enhancements in extending the efficacy of current antibiotics. Cell envelope is the main target for drug-resistant mechanisms of various pathogens because many drugs need to rapidly diffuse into a cell to meet their targets (Nikaido, 1994; Savage, 2001; Zhang et al., 2001).

Biofilms constitute a threat in the clinical environment by acting as pools of multidrug resistant pathogenic bacteria. Biofilm may be formed in a variety of surfaces including living tissues, indwelling medical devices, portable water system piping or natural water system piping (Kokare et al., 2009).

Diverse mechanisms allow microorganisms to come into closer contact with a surface, attach to it, promote cell-cell interactions and grow as a 3-D structure (Bryers, 2000). There are several mechanisms influencing the attachment of biofilm. The properties of the surface of attachment influence the process. In the case of a rougher, more hydrophobic, and coated surface, the attachment will occur efficiently (Donlan, 2002; Pereira, 2001). Other variable can affect the attachment: flow velocity, water temperature or nutrient concentration (Pereira, 2001; Vieira, 1995). Maximum attachment depends upon high surface free energy or wettability of surfaces. Surfaces with high surface free energies are more hydrophilic and generally show greater bacterial attachment than hydrophobic surfaces.

Quorum sensing, related with cell-cell signalling, play a role in cell attachment and detachment from biofilms (Simões, 2005). This mechanism induces and increased in the intrinsic antimicrobial resistance of biofilm (Brooun et al., 2000). QS regulates a wide number of physiological activities, such as motility, conjugation, competence, sporulation, virulence and biofilm formation. The signal of QS may alter distribution of bacterial species in the biofilm, alter protein expression, introduce new genetic trait and incorporate bacteria in biofilm. The properties of cells, including cell surface hydrophobicity, presence of fimbriae and flagella, and production of EPS influence the attachment of microbial cell (Watnick and Kolter, 2000).

One key element of the adaptability of bacterial cells is their ability to be in a niche where they can propagate. Various mechanisms of motility have been described for bacteria. The most common is flagellar motility, but other mechanisms can be highlighted: twitching, gliding, darting, sliding, swimming and swarming (Davey and O'Toole, 2000).

The aim of this study was to assess aspects of the mode of action of the selected phytochemicals on planktonic cells and in the early stages of biofilm formation. The physicochemical properties of bacterial surface (zeta potential and hydrophobicity) were performed with and without phytochemicals. The evaluation of phytochemicals to inhibit efflux pumps was studied with *S. aureus* while the effects of these products were tested on *E. coli* outer membrane proteins (OMPs) expression. The free energy of adhesion, quorum-sensing inhibition and bacterial motility were assessed in order ascertain the role of phytochemicals on the early stages of biofilm development.

4.2 Materials and methods

4.2.1. Bacterial Strains

Escherichia coli (CECT 434) *Staphylococcus aureus* (CECT 976) were used according to section 3.2.1.. The *Chromobacterium violaceum* (ATCC 12472) were distributed over the surface of Luria–Bertani Agar (Merck) and incubated for 24 h at 30 ± 3 °C. *S. aureus* RN4220 containing plasmid pUL5054, which carries the gene encoding the MsrA macrolide efflux protein; *S. aureus* SA1199B, which overexpresses the NorA MDR efflux protein and *S. aureus* XU212, which possesses the TetK efflux pump and is also a MRSA strain, were kindly provided by S. Gibbons (University College London, UCL) (Gibbons et al., 2003; Oluwatuyi et al., 2004; Gibbons and Udo; 2000; Smith et al., 2007). Prior to use, these strains at -80°C were transferred onto Mueller-Hinton (Merck, Germany) agar plate, grown overnight, and inoculated into MH broth at 30 °C and under agitation (150 rpm).

4.2.2. Phytochemicals and antibiotics

The phytochemicals used were the same referred before and prepared as explained in section 3.2.2. Ciprofloxacin, erythromycin and tetracycline were obtained from Sigma (Portugal). Every antibiotic were prepared in DMSO, which was filtrated before utilization to avoid contamination. After preparation, antibiotics were frozen.

4.2.3. Determination of Zeta Potential

The overnight cultures of *E. coli* and *S. aureus* were centrifuged (Eppendorf centrifuge 5810R) at 3777 g for 10 min and washed twice with sterile water. The OD_{620 nm} of strains was adjusted to 0.2 ± 0.02 and samples were incubated with phytochemicals for 30 minutes at 30 °C. Phytochemical concentration used was the MIC. Cells suspensions without phytochemicals were used as control. The zeta potential experiments were performed using a Malvern Zetasizer instrument (Nano Zetasizer, Malvern instruments, UK). All experiments were carried out in triplicate at room temperature and were repeated at least at three different occasions.

4.2.4. Physico-chemical characterization of bacterial surface

The physico-chemical properties were measured using the sessile drop contact angle method. The bacteria cultures were grown overnight in MH medium. The cells suspensions were washed with NaCl (8.5 g/L) and centrifuged (10 minutes at 3777 g) twice. The OD_{620nm} was adjusted to 0.4 ± 0.02. After that, the biocides were applied (at MIC concentration) during 30 minutes. The solutions were filtrated (0.45 µm, Whatman) and placed in microscope slides. The contact angle was measured with 3 different liquids: water, formamide (polar) and α-bromonaphtalene (nonpolar) (Sigma, Portugal).

The measurement of contact angles was performed using a model OCA 15 Plus (Dataphysics, Germany) video based optical contact angle measuring instrument, allowing image acquisition and data analysis. The degree of hydrophobicity of a surface is expressed as the free energy of interaction between entities of that surface, when immersed in water. ΔG_{s_{ws}} (mJ/m²) can be positive or negative according of the interaction between the surfaces. In the case of ΔG_{s_{ws}} > 0, the material is considered hydrophilic, because the interaction between the two surfaces is weaker than the interaction of each entity with water. In contrast, when ΔG_{s_{ws}} < 0, the interaction between the surfaces is stronger than the interaction of each entity with water and the material is hydrophobic. Hydrophobicity was evaluated after contact angles measurements, following the van Oss approach (van Oss *et al.*, 1987; 1988; 1989).

The degree of hydrophobicity can be calculated through the surface tension components of interacting entities, according to:

$$\Delta G_{s_{ws}} = -2\left(\sqrt{\gamma_s^{LW}} - \sqrt{\gamma_w^{LW}}\right)^2 + 4\left(\sqrt{\gamma_s^+ \gamma_w^-} + \sqrt{\gamma_s^- \gamma_w^+} - \sqrt{\gamma_s^+ \gamma_s^-} - \sqrt{\gamma_w^+ \gamma_w^-}\right) \quad (3)$$

Where, γ^{LW} is the Lifshitz-van der Waals component of the surface free energy and γ^+ and γ^- are the electron acceptor and donor, respectively, of the Lewis acid-base parameter (γ^{AB}), being $\gamma^{AB} = \sqrt{\gamma^+ \gamma^-}$.

The analysis was performed at room temperature using the three liquids referred before. The surface tension components of liquids were obtained from literature (Janczuk *et al.*, 1993). Subsequently, three equations can be solved:

$$(1 + \cos\theta)\gamma_1^{Tot} = 2\left(\sqrt{\gamma_s^{LW} \gamma_w^{LW}} + \sqrt{\gamma_s^+ \gamma_w^-} + \sqrt{\gamma_s^- \gamma_w^+}\right) \quad (4)$$

Where θ is the contact angle and $\gamma^{TOT} = \gamma^{LW} + \gamma^{AB}$.

These measurements were performed at least with 12 determinations for each liquid and microorganism.

4.2.5. Free energy of adhesion

The free energy of adhesion between the bacterial cells and polystyrene surfaces was calculated through the surface tension components of the entities involved in the process using the Dupré equation and the procedure described by Simões et al. (2010). The total interaction energy (ΔG_{bws}^{TOT}) is studied by the interaction between one bacteria (b) and a substratum (s) that are immersed or dissolved in water (w) and is expressed by the interfacial tension components:

$$\Delta G_{bws}^{TOT} = \gamma_{bs} - \gamma_{bw} - \gamma_{sw} \quad (5)$$

The thermodynamic theory of the interfacial tension of one system of interaction (for example, bacteria/surface - γ_{bs}) can be defined by the following equations:

$$\gamma_{bs} = \gamma_{bs}^{LW} + \gamma_{bs}^{AB} \quad (6)$$

$$\gamma_{bs}^{LW} = \gamma_b^{LW} + \gamma_s^{LW} - 2 \times \sqrt{\gamma_b^{LW} \times \gamma_s^{LW}} \quad (7)$$

$$\gamma_{bs}^{AB} = 2 \times \left(\sqrt{\gamma_b^+ \times \gamma_b^-} + \sqrt{\gamma_s^+ \times \gamma_s^-} - \sqrt{\gamma_b^+ \times \gamma_s^-} - \sqrt{\gamma_b^- \times \gamma_s^+} \right) \quad (8)$$

The two other interfacial tension components, γ_{bw} and γ_{sw} , were calculated in the same way, which permits the assessment of thermodynamic energy of adhesion. The bacterial adhesion to the substratum can be favorable or is not expected to occur, according to the values of ΔG_{bws}^{TOT} if are negative or positive, respectively (Simões et al., 2010).

4.2.6. Motility

Plates containing 1% tryptone (Merck), 0.25% NaCl (Merck) and 0.25% or 0.7% (w/v) agar (Merck) were prepared for swimming/sliding or swarming motilities,

respectively (Butler et al., 2010; Stickland et al., 2010). Phytochemicals at MIC concentration were incorporated in the growth medium after sterilize and cooling the medium, to avoid the deterioration. Overnight cultures of *E. coli* and *S. aureus* grown on LB broth (Merck, Germany) were used to determine bacterial motility. A volume of 15 μL of cell suspension ($\text{OD}_{620\text{nm}} = 0.4 \pm 0.02$) was placed in the center of the plates. Then, plates were incubated at 30 °C and the diameter (mm) of the bacterial motility halos were measured at 24, 48 and 72 h (Borges et al., 2012). All experiments were carried out in triplicate. The negative control was performed with DMSO.

4.2.7. Detection of quorum-sensing inhibition

The culture of *Chromobacterium violaceum* (CV12472) was grown overnight in Luria-Bertani (LB) broth at 30 °C. The $\text{OD}_{620\text{nm}}$ of the strain was adjusted to 0.1 ± 0.02 (1.4×10^8 CFU/mL). For each phytochemical a sterile 96-well polystyrene microtiter plates were filled with cells (180 μL) and phytochemicals (20 μL). As a negative controls were used cell suspension with DMSO and without phytochemicals. The microtiter plates were incubated during 24 h at 30 °C in an orbital shaker (150 rpm). Subsequently, the absorbance at 620 nm was measured using a Microplate Reader (Spectramax M2e, Molecular Devices, Inc.). MIC values were determined using the microdilution method, explained above. All tests were performed in triplicate. All the further experiments were performed at sub-MIC concentrations of phytochemicals (Packiavathy et al., 2012).

After these experiments, *C. violaceum* was used to perform the detection of quorum-sensing inhibition (QSI) by the disc diffusion method. The detection of QS activity of phytochemicals was performed at the range of sub-MIC. LB agar (LBA) plates were spread with 100 μL ($\text{OD}_{620\text{nm}}$ of 0.1 ± 0.02) (1.4×10^8 CFU/mL) of overnight culture of *C. violaceum* CV12472. Sterile paper disks (6 mm diameter) were placed in the plates and impregnated with various concentrations of each phytochemical (15 μL). DMSO was used as a negative control. The plates were incubated at 30 °C for 24 h to check the inhibition of pigment production around the disc. The growth inhibition was also recorded. The zones of QS inhibition were measured from the disks to the edges (Adonizio et al., 2006; Khan et al. 2009). Bacterial growth inhibition by the phytochemicals was measured as radius (r_1) in mm while phytochemicals showing both growth and pigment inhibition was measured as

radius (r_2) in mm. The pigment inhibition (QS) was determined by subtracting bacterial growth inhibition radius (r_1) from the total radius (r_2) thus QS inhibition = (r_2-r_1) in mm (Zahin et al. 2010).

4.2.8. Outer Membrane Proteins

4.2.8.1. Extraction

The Outer Membrane Protein extraction was performed for *E. coli*. The bacterial culture was grown overnight at 30 °C in an orbital shaker (120 rpm). The inoculum was centrifuged at 3777 g for 15 minutes and washed with twice NaCl (8.5 g/L). After that, the OD was measured and adjusted to 0.4 ± 0.02 ($\lambda = 620$ nm). The cell suspension was incubated for 1h with the different phytochemicals at MIC.

After that, the cells suspension were centrifuged and the pellet was resuspended in 25 mM Tris and 1 mM MgCl₂ (Merck) buffer (pH 7.4), twice. Behind this process, each sample was sonicated for 5 times (20 seconds) (Vibracell, 60 W) on ice to promote cell lysis. After sonication the solution was centrifuged (7000 g, 10 min, 4°C) (Beckman Avanti J25 centrifuge) in order to remove non-lysed cells. The supernatant was collect and 1 mL of Sarcoisine (Sigma) solution was added to obtain a final concentration of 2% (w/v), in order to solubilize the inner membrane proteins. The samples were left on ice for 20 minutes. After incubation time, 25 mL of Tris-HCl 25 mM and 1mM MgCl₂ buffer. The solution was centrifuged (13000 g, 1 h, 4 °C) (Avantis J-25) two times, to recover the pellet containing the OMP. The pellet was resuspended in 300 μ L of 25 mM Tris-HCl buffer (pH 7.4) and stored at -20 °C until required.

4.2.8.2. OMP Analysis

The protein content of OMP samples was determined using Bicinchoninic Acid Protein Assay Kit (BCA) (BCA-PIERCE Cat. No. 23225) with BSA as standard. This procedure was applied in order to insert in the gel cassettes the same OMP concentration for the several samples (3 μ g protein in each well).

The OMP samples obtained were subjected to SDS-PAGE, as reported by Laemmli (1970) with 12% (w/v) acrylamide (Bio-rad). Electrophoresis was

performed at a constant current of 10 mA. After electrophoresis, the gel was stained with Coomassie blue (Bio-rad) for protein profile detection.

4.2.9. Antibiotic-Phytochemical Dual Combinations Assay – Efflux Pumps Inhibition

To study the antimicrobial effects of phytochemicals conjugated with antibiotics, they were inserted in MH agar medium (at MIC). The phytochemical was inserted after sterilize and cooling the medium, to avoid the deterioration. Colonies of bacteria were picked from overnight PCA cultures (log phase cultures) in solid medium (Merck, Portugal). The suspension of bacteria was prepared with 0.9% NaCl, which was adjusted to match to 0.5 McFarland turbidity standards. The suspension was spread with a sterile cotton swap into Petri dish (90 mm of diameter) containing 20 ml of Mueller-Hinton Agar. Sterile filter paper discs (with 6 mm in diameter), impregnated with 15 μ L of antibiotics, were placed on the agar plate seeded with the respective bacteria. Discs of ciprofloxacin, erythromycin and tetracycline were used as positive controls and discs impregnated with DMSO were used as negative controls. The concentration of antibiotics used was according to Clinical and Laboratory Standards Institute (2005): Performance Standards for Antimicrobial Susceptibility Testing (Fifteen Informational Supplement): ciprofloxacin – 5 μ g/disc; erythromycin – 15 μ g/disc; and tetracycline – 30 μ g/disc. The plates were incubated at 30 °C for 24 hours. After incubation, zones of growth inhibition were measured. All tests were performed in triplicate and the antibacterial activity was expressed as the mean of inhibition diameters (mm).

4.2.9.1. Classification of dual combinations

The effect of dual combinations of antibiotics and phytochemicals can be classified according Saavedra et al. (2010):

- Antagonism (-) – $[\text{inhibition halo} - (\text{antibiotic inhibition halo} + \text{phytochemical inhibition halo})/2] < 0$
- Indifference (+) – $0 \leq [\text{inhibition halo} - (\text{antibiotic inhibition halo} + \text{phytochemical inhibition halo})/2] < \text{antibiotic inhibition halo}$ or $\text{phytochemical inhibition halo}$

- Additive (++) – antibiotic inhibition halo < [inhibition halo – (antibiotic inhibition halo + phytochemical inhibition halo)/2] < 2x antibiotic inhibition halo or phytochemical inhibition halo
- Synergy (+++) – inhibition halo > 3x antibiotic inhibition halo or phytochemical inhibition halo.

For this classification the highest inhibition halos caused by the antibiotic or phytochemical were used.

In the case of no effect of phytochemicals, the classification of the dual combination is different (Abreu et al., 2013b):

- Additive (++) – 4 mm ≤ inhibition zone diameter combination – inhibition zone diameter most active agent) < 6 mm
- Synergistic (+++) – inhibition zone diameter combination – inhibition zone diameter most active agent ≥ 6 mm.

4.2.10. Statistical analysis

The data was analyzed using One-Way Anova and the statistical program SPSS 21.0. The results were presented as the means ± SEM (standard error of the mean). Significance level for the separation was set at $p < 0.05$ and the calculations were based on confidence level equal or higher than 95%.

4.3 Results and Discussion

4.3.1. Surface characterization of cells in presence of phytochemicals

The surface charge of the cells was measured as its zeta potencial, calculated from the mobility of cells in the presence of an electrical field under well-known conditions (Ferreira et al., 2010). It is possible to observe the variations of zeta potencial with several phytochemicals in Table 4.

Table 4. Zeta potencial (mV) results of suspensions of *E. coli* and *S. aureus* in contact with phytochemicals at the MIC.

	Zeta Potencial (mV)	
	<i>E. coli</i>	<i>S. aureus</i>
Control	-13.0 ± 1.4	-29.8 ± 1.3
7-HC	-13.5 ± 1.8	-26.0 ± 6.2
I3C	-21.0 ± 5.7	-27.3 ± 6.3
SA	-1.8 ± 0.3	1.8 ± 0.5
SP	-12.7 ± 1.6	-19.4 ± 2.2

Both bacterial strains present a negative surface charge. Most of the microorganisms are negatively charged due to the presence of anionic groups, as carboxyl and phosphate, in the membranes (Ferreira et al., 2010).

E. coli presents a less negative surface charge. The constitutive molecules of the membrane are responsible for the surface properties and their thermodynamic and bacterial adhesion. A thin layer of peptidoglycan composes the Gram-negative bacteria and proteins, lipopolysaccharides, constitute their external surface. The charge of the bacteria is negative due to carboxyl and phosphoryl groups. So, it is highly charged and interactive with cations. However, the Gram-positive bacteria are composed by a thick layer of peptidoglycan, which is covered for neutral and acid polysaccharides, proteins and teichoic acids (Araújo et al., 2010). The magnitude of negative charge on the cell wall varies from strain to strain (Dickson and Koohmaraie, 1989).

Comparing the values of the control with those caused by the phytochemicals exposure, it is possible to see that 7-HC and SP have no significant effect in the charge of *E. coli* ($p > 0.05$). In contrast, I3C and SA seem to change the surface charge ($p < 0.05$). The contact of I3C with *E. coli* surface makes it more negative, which means that negative charges are transferred for the surface when in contact with this

phytochemicals. In the case of SA, positive charges are transferred for the membrane surface, because *E. coli* surface is less negative. The positive charge of the phytochemical may explain this change in the charge.

The charge of *S. aureus* surface was changed when in contact with SA and SP ($p < 0.05$). The first one is able to make the membrane surface positive, which means that positive charges are moved to the surface. The positive charge of the phytochemical may explain the change in the charge, as referred above. The exposure to SP changes the surface charge of cells to less negative values.

The hydrophobicity has been characterized as one important aspect in bacterial adhesion (Chaves, 2004). Hydrophobicity can be studied through the surface tension by measurement of contact angles. The contact angles were measured using the sessile drop method. van Oss and Giese (1995) developed a methodology to calculate the hydrophobicity of material surfaces through the surface tension parameters. Table 5 shows the variations in the hydrophobicity parameter according the different phytochemicals tested at the MIC.

Table 5. Hydrophobicity (ΔG^{TOT}), and apolar (γ^{LW}) and polar (γ^{AB}) components of the surface tension of untreated and treated cells. The means \pm SDs are illustrated.

Bacteria	Phytochemical	Surface Tension Parameters (mJ/m ²)				Hydrophobicity (mJ/m ²)
		γ^{LW}	γ^{AB}	γ^+	γ^-	ΔG^{TOT}
<i>E. coli</i>	Control	33.6 \pm 5.0	22.4 \pm 5.4	2.6 \pm 0.5	52.0 \pm 4.8	28.9 \pm 7.1
	7-HC	30.7 \pm 4.8	20.8 \pm 4.7	1.5 \pm 0.3	44.9 \pm 7.1	21.0 \pm 5.1
	I3C	20.2 \pm 4.8	37.1 \pm 7.3	7.8 \pm 1.9	55.8 \pm 6.7	20.6 \pm 3.8
	SA	31.0 \pm 5.2	2.07 \pm 6.0	1.5 \pm 0.3	59.7 \pm 11.7	37.6 \pm 18.8
	SP	21.0 \pm 1.8	40.1 \pm 3.1	7.8 \pm 1.3	52.1 \pm 2.8	19.7 \pm 3.3
<i>S. aureus</i>	Control	35.4 \pm 5.4	19.7 \pm 4.6	2.0 \pm 0.4	53.5 \pm 3.8	30.2 \pm 3.2
	7-HC	36.2 \pm 3.4	21.1 \pm 3.8	2.7 \pm 0.3	47.8 \pm 4.2	22.4 \pm 4.8
	I3C	34.5 \pm 4.2	20.4 \pm 4.8	2.2 \pm 0.5	55.4 \pm 5.1	32.2 \pm 7.3
	SA	37.4 \pm 3.0	15.2 \pm 3.5	1.5 \pm 0.3	44.8 \pm 7.7	22.6 \pm 5.6
	SP	36.1 \pm 4.4	18.3 \pm 4.4	2.1 \pm 0.5	54.4 \pm 1.5	30.4 \pm 2.9

The hydrophilic and hydrophobic character of surfaces can be determined by the surface tension components:

- γ^{LW} represents the apolar interactions that are established, meaning the higher it is the value, more apolar is the surface.
- γ^{AB} measures the polar interactions and can be attractive or repulsive.

- γ^- is the donor of electrons.
- γ^+ is the electrons receptor.

The hydrophobicity can also be calculated through the energy of hydrophobic attraction (ΔG^{TOT}). If $\Delta G^{TOT} < 0$ the interaction between the surface molecules is attractive, means that molecules have less affinity for water than among themselves, and it is considered hydrophobic. If $\Delta G^{TOT} > 0$ the surface is considered hydrophilic, and the interaction between the surface molecules is repulsive. So, the more negative the value of ΔG^{TOT} , more hydrophobic is the surface; and the more positive the value of ΔG^{TOT} , more hydrophilic is the surface (Araújo et al., 2010; Chaves, 2004).

Regarding the several parameters obtained, the apolar interactions (γ^{LW}) are influenced when *E. coli* is treated with I3C and SP ($p < 0.05$), making the molecules of surface less apolar. The same phytochemicals are also able to change the molecules surface by making them more hydrophilic (γ^{AB}) ($p < 0.05$). *E. coli* is mainly electron donor. However, this ability to give electrons increases with the treatment with SA.

In the case of *S. aureus*, the treatment with the several phytochemicals did not permit variations on the polarity of surface molecules ($p > 0.05$). Except in the case of SA, the value of γ^{AB} was lower than the observed in the control, which means that the surface of molecules was less hydrophilic. Regarding their capacity of giving electron, 7-HC and SA varied significantly ($p < 0.05$). These phytochemicals decrease the capacity of supply electrons.

Finally, comparing the values obtained for *E. coli* and *S. aureus* of ΔG^{TOT} it is concluded that they have hydrophilic character ($\Delta G^{TOT} > 0 \text{ mJ/m}^2$). The interaction of bacteria in the surfaces is thermodynamically not favourable. The application of SA and SP influences their hydrophilic character, in the case of *E. coli* ($p < 0.05$). SA promotes the increase and SP promotes the decrease of hydrophilic properties. For *S. aureus*, 7-HC and SA decrease their hydrophilic character ($p < 0.05$). These phytochemicals significantly interact with bacterial surface constituents, modifying their physico-chemical properties.

4.3.2. Analysis of *E. coli* and *S. aureus* free energy of adhesion

The bacterial adhesion is a complex process that can be influenced by several factors: physico-chemical characteristics of bacteria (hydrophobicity and surface charge), material surface properties and environmental factors (temperature, pH,

exposure time, concentration of bacteria, chemical treatment or the presence of antimicrobials and fluid flow conditions) (Simões et al., 2010). Behind the physic conditions, biological properties of bacteria also influence the attachment to surface, such as: presence of fimbriae and flagella, and the production of EPS (Simões et al., 2010).

The PS microtiter plates are usually used as the standard bioreactor system for adhesion and biofilm formation bacteria. According to the PS surface properties, this material was characterized for being hydrophobic ($\Delta G^{\text{TOT}} = -44 \text{ mJ/m}^2$) (Simões et al., 2010) (Appendix - A). So, in order to predict the ability of microorganisms to adhere to PS surfaces, the free energy of interaction between the bacterial surface and the PS surface was calculated according to the thermodynamic approach. Table 6 shows the thermodynamical ability of bacteria to adhere to PS.

Table 6. Free energy of adhesion ($\Delta G^{\text{TOT}}_{\text{bws}}$) of bacterial cells to polystyrene, treated and untreated with phytochemicals.

Strain	Phytochemical	Free Energy of Adhesion (mJ/m^2)
<i>E. coli</i>	Control	4.4 ± 1.2
	7-HC	3.5 ± 0.5
	I3C	15.3 ± 3.3
	SA	10.3 ± 2.1
	SP	13.5 ± 2.1
<i>S. aureus</i>	Control	5.7 ± 1.2
	7-HC	1.4 ± 0.2
	I3C	6.4 ± 1.2
	SA	-3.2 ± 0.4
	SP	5.4 ± 0.5

According the values of free energy, it is possible to conclude that only *S. aureus* treated with SA have theoretical thermodynamic ability to adhere to PS ($\Delta G^{\text{TOT}} < 0 \text{ mJ/m}^2$).

Comparing the free energy of adhesion for both bacteria, it is possible to conclude that *S. aureus* and *E. coli* are not significantly different in terms of capacity to adhere to PS ($p > 0.05$).

The addition of I3C influences adhesion of *E. coli* to PS ($p < 0.05$), the value of free energy increases, so the adhesion is not expected to be favorable. In contrast, 7-HC decreases the value of free energy, meaning that the theoretical adhesion to PS is

more favorable. In the case of *S. aureus* with 7-HC and SA, the bacteria have more ability to adhere to PS (ΔG^{TOT} is inferior than control) ($p < 0.05$).

It is important to refer that microorganisms have different mechanisms of adhesion and retention, influenced by the substrata, nutrients, ionic strength, pH values and temperatures, and also their phenotype and genotype (Simões et al., 2010).

The ability of microorganisms to attach to the surfaces is crucial for the beginning of bacterial surface colonization. Adhesion is mediated by two mechanisms. Specific bacterial colonization factors, or adhesins, can act as ligands to mediate attachment of the organisms to specific receptors. Bacteria can also bind by a less specific mechanism mediated by hydrophobic domains (Drumm et al., 1989)

The thermodynamic theory of adhesion permits the quantification of the free energy of adhesion and predicts the possibility of establishment of an interface between the surface and microorganism. However, this theory does not quantify the electrostatic interactions and microbiologic aspects of adhesion (Chaves, 2004).

The hydrophobicity is a surface property that plays an important role in the adhesion phenomena. Several authors explained that increasing the hydrophobicity produces an increasing extension of adhesion (Ghannoum, 1992; Masuoka and Hazen, 1997; Sinde et al., 2000).

4.3.3. Motility assays

Motility is one of the most important features in microbial physiology. The bacteria show different ways of motility. In the case of swimming and swarming, motility is dependent on flagella. These types of motility contribute to the virulence of pathogens through adhesion and biofilm formation on biotic and abiotic surfaces (Borges et al., 2012). Swarming motility is also important for the early stages of biofilm formation. Sliding motility is produced by the expansive forces of a growing colony in combination with reduced surface tension. This type of motility is important in bacterial surface colonization (Harshey, 2003). The phytochemicals at the MIC were tested for their ability to act in motility inhibition. The results obtained are presented in Table 7.

Table 7. Motility results for bacteria with and with phytochemicals. The drop baseline was 6mm which was subtracted from the results presented.

Time/ Phytochemical	<i>E. coli</i>		<i>S. aureus</i>
	Swimming (mm)	Swarming (mm)	Sliding (mm)
24h			
Control	79.0 ± 1.2	8.7 ± 0.6	7.0 ± 0.0
7-HC	7.0 ± 1.0	7.7 ± 1.5	5.0 ± 0.0
I3C	4.7 ± 0.6	7.7 ± 2.9	0.0 ± 0.0
SA	3.3 ± 0.9	2.0 ± 0.9	7.7 ± 0.6
SP	80.0 ± 0.0	56.0 ± 2.0	84.0 ± 0.0
48h			
Control	84.7 ± 0.6	13.7 ± 3.8	8.0 ± 1.0
7-HC	43.3 ± 2.9	8.7 ± 1.2	8.3 ± 0.6
I3C	0.0 ± 0.0	10.0 ± 7.8	0.0 ± 0.0
SA	0.0 ± 0.0	55.0 ± 8.7	0.0 ± 0.0
SP	84.0 ± 0.0	61.7 ± 9.1	56.7 ± 5.8
72h			
Control	84.0 ± 0.0	64.3 ± 7.6	7.7 ± 0.6
7-HC	51.3 ± 2.3	8.3 ± 0.6	8.3 ± 0.6
I3C	0.0 ± 0.0	8.7 ± 5.5	0.0 ± 0.0
SA	0.0 ± 0.0	54.3 ± 6.4	2.0 ± 0.6
SP	84.0 ± 0.0	13.3 ± 3.2	55.0 ± 8.7

Swimming and swarming motilities have been documented as two forms of surface motility for *E. coli*, and sliding for *S. aureus* (Borges et al., 2012; Harshey, 2003; Pratt and Kolter, 1998). *E. coli* exhibits flagella; in contrast, *S. aureus* is a non-flagelated bacterium with a motility phenomenon defined as colony spreading (Borges et al., 2012).

The Gram-negative bacteria increased their growth over the time and the Gram-positive bacteria maintained its growth, without the addiction of phytochemicals. In this study, it was verified that *E. coli* presented the highest motility and also showed an increasing in swimming and swarming motility, without the addiction of phytochemicals. However, *S. aureus* had more capacity to adhere to PS, concluding that motility does not regulate adhesion.

Regarding the several phytochemicals performed, motility was mostly affected when I3C is added. In the case of swimming and sliding, the motility was completely inhibited with this phytochemical ($p < 0.05$). However, I3C did not influence swarming motility. SA was also able to stop swimming and sliding motilities. Swarming motility was very low in the first 24h, and in the last 48h, increased. Probably, *E. coli* is able to adapt to SA after a long period of exposure. Swimming motility increased with the addiction of SP after 72h of exposure ($p < 0.05$); however, with swarming and

sliding motility, the bacteria showed an increase in the first hours, but after a long period of exposure, the motility decreased. Finally, 7-HC influenced swarming and swimming motility and it was not able to change sliding motility. In the case of swimming motility, the value was very low in the first 24h, after that, it started to increase.

Borges et al. (2012) also studied *E. coli* and *S. aureus* motilities in contact with ferrulic and gallic acids. They concluded that both compounds show potential to inhibit cell motility. Swarming was completely inhibited in *E. coli*, after 24h; Swimming motility of *E. coli* was reduced by the addition of both compounds. *S. aureus* motility was also inhibited by ferrulic acid. Although the compounds tested are not the same, it is possible to conclude that the motility can be changed.

The motility is apparently related with the cell state of the bacteria. The changes in the motility are explained by cell stimuli to alter the functions of its motility machinery to improve or decrease its chances of migrating to a better location (Jarrell and McBride, 2008). Morphological differentiation in bacteria, which distinguishes them from their planktonic state, occurs as a response of motility (Julkowska et al., 2004).

A relationship between cells surface motility and biofilm formation has been reported, especially in the case of swarming motility. Both processes, biofilm formation and swarming, require production of flagella and surface polysaccharides (Borges et al., 2012). Several authors have been reported mutants with altered swarming motility that present difficulties in biofilm formation, concluding that they can play a role in biofilm development (Pratt and Kolter 1998; Shrouf et al., 2006).

4.3.4. Quorum-sensing assays

QS is a mechanism by which a bacterial population senses its cell density (Khan et al., 2009). This mechanism influences bacterial biofilm growth and development and it is related to cell-cell interactions (Simões et al., 2009). This cell-cell communication is dependent of several factors: synthesis, exchange and perception of small signal molecules between bacteria (Khan et al., 2009). The 4 phytochemicals was tested as QS inhibitors, at several concentrations. The Table 8 shows the results obtained.

The MIC of the phytochemicals tested against *C. violaceum* CV12472 ranged from 25 to 3200 $\mu\text{g/mL}$. This means that phytochemicals are able to inhibit the bacteria growth. QS results show the effect of the phytochemicals in the bacteria growth (inhibition halo) and also the effect of phytochemical in quorum-sensing, through the detection of pigment inhibition (QS halo). Inhibition of pigment production was detected with some phytochemicals at different concentrations. Figure 4 shows several agar plates with different results.




Figure 4. Examples of results obtained with quorum-sensing assay: (a) no bioactivity (SA at 5000 $\mu\text{g/mL}$); (b) antibacterial and QSI halos are observed (I3C at 5000 $\mu\text{g/mL}$) with addition of phytochemicals.

Of the 4 phytochemicals tested, inhibition of pigment production was detected with 7-HC, I3C and SA with zones of pigment inhibition ranging 5 to 19 mm. No effect on the pigment was observed with SA at the concentrations tested.

Table 8. Quorum sensing results for several phytochemicals at different concentrations.

Phytochemical Concentration ($\mu\text{g/mL}$)	7-HC			I3C			SA			SP		
	QSI pigment	Inhibition halo (mm)	QS halo (mm)	QSI pigment	Inhibition halo (mm)	QS halo (mm)	QSI pigment	Inhibition halo (mm)	QS halo (mm)	QSI pigment	Inhibition halo (mm)	QS halo (mm)
Control	+	5	n.d.									
DMSO	+	15	n.d.									
250	+/-	12	n.d.	+	11	n.d.	+	14	n.d.	+	10	n.d.
500	+/-	11	n.d.	+/-	11	5	+	14	n.d.	+	11	n.d.
750	+/-	10	n.d.	+/-	11	6	+	19	n.d.	+	12	n.d.
1000	+/-	11	7	+/-	11	14	+	16	n.d.	+	14	n.d.
1500	+/-	11	7	+/-	12	15	+/-	14	6	+	9	n.d.
2000	+/-	10	5	-	16	9	-	16	8	+	10	n.d.
3000	+/-	10	8	-	20	5	-	16	8	+	10	n.d.
4000	+/-	12	10	-	20	11	-	15	12	+	10	n.d.
5000	+/-	11	5	-	25	9	-	18	9	+	11	n.d.

(+) - There was formation of purple pigment in the plate; (-) - Purple pigment was not formed in the plate; (+/-) - The pigment formed was clearer.
n.d. - not detected.

The I3C is the most effective to inhibit production of pigment. At 500 µg/mL, the inhibition is low, but by increasing the concentration the zone of pigment inhibition is also increasing. Regarding 7-HC and SA, at low concentrations there is no inhibition of pigment production; but from 1000 and 1500 µg/mL of 7-HC and SA, respectively, the QSI halo is detectable. Although, SA shows antimicrobial activity, it was not possible to observe an effect on pigment inhibition at the concentrations tested. Regarding QS halos, the concentration of I3C, 7-HC and SA added influences significantly the quantity of pigment production. So, QS activity of phytochemicals is concentration dependent, as reported by other authors (Khan et al., 2009; Zahin et al., 2010). The same authors identified one compound (clove oil) able to inhibit pigment production with 19 mm of pigment inhibition zone against *C. violaceum* (CV12472). Also cinnamon, peppermint and lavender present zones of pigment inhibition against the same bacteria (Khan et al., 2009). Al-Hussaini and Mahasneh (2009) reported *T. capensis*, *Sonchus oleraceus*, *Pityriasis alba*, *Pinus nigra*, *Jasminum sambac*, *Rosmarinus officinalis*, *Lavandula angustifolia* and *Laurus nobilis* as great sources of microbial growth and QS inhibitors.

The phytochemicals and other compounds that affect QS can interfere at different levels: inhibition of signal biosynthesis or inhibition of activity of AHL-producing enzymes, enzymatic signal degradation and inhibition of reception signal molecules (Khan et al., 2009).

4.3.5. Characterization of cell membranes

In order to characterize both bacterial cell membranes, two different techniques were performed: outer membrane protein extraction and analysis for *E. coli*, and study of efflux pumps for *S. aureus*.

4.3.5.1. Outer Membrane Proteins

The OMPs are key molecules that are the interface between the cell and the environment. *E. coli* comprises three different layers, as explained before. The external layer consists of lipids, polysaccharides and proteins. Some bacterial proteins are expressed in high copy number, which aid their detection and characterization. These include porins (OmpC and OmpF). Porins are a protein family of OMPs that

form a hydrophilic channel, permitting nonspecific diffusion of small molecules across the outer membrane (Molloy et al., 2000). Due to their location, OMPs are important candidate antigens for the development of strategies to protect against bacterial pathogens (Kawahara et al., 1994a; Kawahara et al., 1994b; Negm and Pistole, 1998; Pages et al., 1987). The dominant OMPs of *E. coli* are OmpA, OmpX, OmpF and OmpC (Molloy et al., 2000).

The planktonic cells of *E. coli* were characterized phenotypically in terms of OMPs. The OMPs of *E. coli* strains as planktonic cells were isolated and analysed by SDS-PAGE. The OMP profiles obtained with different phytochemicals (at the MIC) are presented in Figure 5.




Figure 5. OMPs profiles of *E. coli*. The profile of molecular weight standards (1), control (2), 7-HC (3), I3C (4), SA (5), SP (6) are presented.

According to Figure 3, the OMP profiles obtained with cells and phytochemicals do not differ considerably from the ones obtained with *E. coli* without phytochemicals (well 2). The application of several phytochemicals does not seem to affect OMP expression since, for all lanes, the protein expression is similar. Nevertheless, the outer membrane of cells with SA exhibits a higher intensity than the other OMP. The profile of OMPs shows different proteins. According to the results the phytochemicals applied do not act on membrane proteins.

Regarding the OMPs analysed, there are three proteins that were isolated with all the phytochemicals and also at the control. The molecular weights of them are 55, 38 and 35 kDa. The 35 kDa OMP it was identified as OmpA (Hellman et al., 2000; Molloy et al., 2000), one of the most important OMPs, as referred before. Other studies have been done to study the OMPs of *E. coli*. Proteins with 38 and 55 kDa were identified as being OmpC and ATP synthase alfa subunit (Molloy et al., 2000; Xu et al., 2006)

OMPs have a significant role in the context of biofilm eradication. The bacterial proteins form an adaptative barrier to the external environment, protecting cells from damaging substances, such as biocides and antimicrobial agents, when procedures of disinfections are applied. This permits the selective uptake of nutrients (Simões, 2005). Due to their location, OMPs are important candidates antigens for the development of strategies to protect bacterial cells against pathogens (Molloy et al., 2000).

4.3.5.1. Efflux pumps

Efflux mechanisms are implicated in antimicrobial. Efflux pumps contribute to the resistance of bacteria by pumping out a wide variety of products: dyes, detergents and antibiotics (Simões, 2005). However, efflux pumps have been studied as membrane components in all cell types, from prokaryotes to eukaryotes (van Bambeke et al., 2007). The role of efflux pumps in bacteria has been related to the elimination of metabolites that are poisonous to the cell and in cell stress responses (Costa et al., 2013). The association of antibiotics with phytochemicals can create a synergistic effect against resistant bacteria, creating new choices for the treatment of infectious diseases.

S. aureus genome reveals high potential multidrug efflux-pump-encoding genes (Huet et al., 2008). Several efflux resistance mechanisms have been described for *S. aureus* such as QacA and NorA, which are multidrug transporters, and the more specific MsrA and TetK transport proteins (Gibbons et al., 2003).

In this study, the antimicrobial activity of several phytochemicals was tested in combination with three antibiotics. For this experiments 4 different strains of *S. aureus* were tested. Table 9 shows the antimicrobial activity of antibiotics against *S. aureus* strains.

According to the Clinical and Laboratory Standards Institute (2005), the organisms are susceptible, intermediate or resistant to the agents. *S. aureus* CECT 976 is considered susceptible to all antibiotics tested. *S. aureus* XU212, *S. aureus* RN4220 and *S. aureus* SA1199B present resistance to tetracycline, erythromycin and ciprofloxacin, respectively.

The negative control performed with DMSO in the preparation of phytochemical solutions presented no effects on bacterial growth.

Table 9. Antimicrobial activity of antibiotics. The means (mm) \pm standard deviation for at least three replicates are illustrated.

	Diameter of inhibition zone (mm)			
	<i>S. aureus</i> CECT 976	<i>S. aureus</i> XU212	<i>S. aureus</i> RN4220	<i>S. aureus</i> SA1199B
TET	41.5 \pm 9.2	16.0 \pm 4.2	42.5 \pm 0.7	46.5 \pm 2.1
ERY	37.5 \pm 3.5	24.5 \pm 4.9	22.0 \pm 4.2	35.5 \pm 3.5
CIP	40.5 \pm 0.7	26.0 \pm 2.8	31.5 \pm 2.1	18.0 \pm 1.4

The antibiotics (tetracycline, erythromycin and ciprofloxacin) had antimicrobial effect against the bacteria tested. Tetracycline was the most effective against all the bacteria, while erythromycin has the lowest antimicrobial activity ($p < 0.05$). Regarding to phytochemicals, I3C and SA showed antimicrobial activity against *S. aureus*, with inhibition halos of 20 and 14 mm, respectively. In contrast, 7-HC and SP do not demonstrate antimicrobial properties against the same strain. The most effective phytochemicals was I3C ($p < 0.05$).

Dual combinations of antibiotic-phytochemicals were performed. Table 10 shows that the combined application of tetracycline, erythromycin and ciprofloxacin had both negative and positive antimicrobial activities compared to the single application of phytochemicals and antibiotics. The classification presented in Table 10 was done according Saavedra et al. (2010).

The combination of bioactive compounds is expected to exert a synergistic effect or to reduce possible adverse side effects. The development of active compounds, such as phytochemicals, in conjunction with antibiotics could avoid the emergence of resistant variants that might otherwise arise during treatment (Abreu et al., 2013b).

Table 10. Classification of the effect of dual combinations of phytochemicals and antibiotics.

		7-HC	I3C	SA	SP
<i>S. aureus</i> CECT 976	TET	+	+++	+	-
	ERY	-	+++	+	++
	CIP	+	+++	+	-
<i>S. aureus</i> XU212	TET	++	+++	+++	+++
<i>S. aureus</i> RN4220	ERY	-	+++	+++	+++
<i>S. aureus</i> SA1199B	CIP	+	+++	+++	+++

(-) – Antagonist; (+) – Indifference; (++) – Additive; (+++) – Synergistic

The results present dual combinations able to improve the antimicrobial activity against the resistant strains. The combined application of ERY with 7-HC, TET or CIP with SP against *S. aureus* CECT 976 was antagonist. Combination of SA with all antibiotics tested showed to be indifferent against *S. aureus* CECT 976. Regarding 7-HC, the combination of phytochemicals with TET and CIP is also indifferent in the antimicrobial activity when compared with single antibiotic and phytochemicals activities against *S. aureus* CECT 976. The dual combination of ERY and SP showed an additive effect against the growth of *S. aureus* CECT 976. Regarding the resistant strains of *S. aureus*, combined application of TET and 7-HC against *S. aureus* XU212 had an additive effect. The remaining combinations produced a synergistic effect.

Antibiotic synergism occurs when the effects of combination of antimicrobials is greater than the sum of the effects of individual antimicrobials (Saavedra et al., 2010). An additive effect of phytochemical combined with antibiotic may occur due to a double attack of both agents at different target sites of bacteria (Adwan and Mhanna, 2008).

The identification of effective efflux pumps inhibitors of *S. aureus* could restore the clinical utility of pump substrates. Efflux pumps inhibitors could extend the useful lifetime of antibiotics by improving therapeutic efficacy, suppressing the emergence of resistant variants, reducing the effective dose of antibiotics to reduce the adverse toxic effects (Gibbons et al., 2003; Saavedra et al., 2010). Combination therapy with two or more antimicrobials is used to prevent the emergence of resistant strains, to treat emergency cases and to take advantage of antimicrobial products synergy (Hemaiswarya and Doble, 2009).

Chapter 5

Concluding remarks and perspectives for further research

5.1. Conclusions

To find new antimicrobial agents, plant products have been studied as substituents of antibiotics for which bacteria already acquired resistance. So, in this work, the antimicrobial effect of four phytochemicals - 7-HC, I3C, SA and SP – was evaluated. After the development of the experiments, it was possible to conclude about the antimicrobial activity of the selected phytochemicals and their ability to control biofilms of two important pathogens – *E. coli* and *S. aureus*.

The 7-HC was one of the most effective phytochemicals tested against *E. coli* and *S. aureus*. The values of MIC obtained were 800 and 200 $\mu\text{g/mL}$ for *E. coli* and *S. aureus*, respectively. However, MBC was not detected for the concentrations tested. Regarding the biofilm control, the exposure of *S. aureus* biofilms to 7-HC at different concentrations, produced significantly different percentages of inactivation. The same phytochemical had no effect in the surface charge of *E. coli*, meaning that this phytochemical did not act in the surface cell membrane. The same phytochemical

decreased the hydrophilic character of *S. aureus* and increase the ability of *S. aureus* to adhere to PS. Swimming and swarming motilities were affected when in contact with 7-HC. However, sliding motility did not change. 7-HC was also able to inhibit the formation of QS pigment (violacein), with halos of 10 mm.

I3C was also effective against both bacteria. The MIC values were 800 and $\mu\text{g/mL}$ for *E. coli* and *S. aureus*, respectively. Regarding the MBC, I3C was also the most effective phytochemicals, with values of 1600 and 800 $\mu\text{g/mL}$ for *E. coli* and *S. aureus*, respectively. By measuring the zeta potential of bacterial strains in contact with I3C, it was concluded that I3C affected the charge of *E. coli*, making the membrane more negative. The values of free energy of adhesion decreased by exposing *E. coli* to I3C, so the adhesion was expected to be more favourable. This phytochemical affected the motility; swimming and sliding motilities were completely inhibited. QSI halos were observed due to I3C, presenting the highest halos (25 mm). Dual combinations of all the antibiotics and I3C presented a synergistic effect against *S. aureus* resistant strains.

SA and SP were the least effective phytochemicals. In the case of SP, the MIC value was not detected for concentrations below to 3200 $\mu\text{g/mL}$.

SA affected the surface charge of *E. coli* and *S. aureus* making the membrane less negative and positive, respectively. This phytochemical influenced the hydrophilic character of *E. coli* and *S. aureus*, increasing and decreasing the character, respectively. SA was the only phytochemical, that when applied to *S. aureus*, was able to make bacteria to adhere to PS, when assessing the free energy of adhesion. Swimming and sliding motilities were inhibited when exposed to SA. Regarding QS, SA presented inhibition zone of pigment production.

Regarding SP, this phytochemical had no significant effect in *E. coli* charge, but influenced the charge of *S. aureus*, making it less negative. The same phytochemical decreased the hydrophilic character of *E. coli* and increased the value of the free energy of adhesion, meaning that the adhesion of *E. coli* to PS was less favourable.

Multi-target antimicrobial products are more effective against Gram-positive bacteria than Gram-negative bacteria (Aires et al., 2009). So, it is important to find antimicrobial products/phytochemicals with significant activity against the more resistant bacteria, which can be applied as therapeutic agents to control infections

caused by them.

This study suggests that 7-HC and I3C are the most promising phytochemicals against *E. coli* and *S. aureus*. Both phytochemicals affected the motility and QS activity, which means that they can play an important role in biofilm prevention and interference with cell-cell interactions. These phytochemicals presented MIC values lower than 1000 µg/mL. The literature has been reported several natural products and extracts with antimicrobial activity with MIC values below 1000 µg/mL, which has important relevance for clinical applications (Gibbons, 2004). In order to apply these phytochemicals to human use, it is necessary the evaluation of their toxicity against mammalian cells. Also, *in vivo* confirmation of the efficacy of these compounds and their side effects would be necessary for a full evaluation of their practical usefulness in the clinical field.

5.2. Perspectives for further research

During this work, the capacity of the selected phytochemicals to control biofilm formation was studied. However, it would be also interesting to study the influence of the selected phytochemicals in biofilm prevention. The same assays could be done, but instead of adding the phytochemicals to the 24-aged biofilms (for 1h), the biofilms should be formed in the presence of the phytochemicals during 24 h.

It would be interesting to develop an assay to evaluate if the phytochemicals can have any influence on the efflux pumps in Gram-negative resistant bacteria and study the dual combinations as it was performed for *S. aureus* resistant strains. So, it would be possible to understand if there are any synergistic effect between the selected antibiotics and phytochemicals against *E. coli* strains. To study the effects of the phytochemicals in the membrane proteins of *S. aureus*, it would be important the development of a protocol to study their membrane and the influence of phytochemicals. After the performance of these studies, it would be possible to conclude more precisely about the mechanisms of action of the phytochemicals.

Although 7-HC and I3C seems to be important plant compounds able to inhibit microbial growth, these phytochemicals should be tested clinically to evaluate their efficacy and cytotoxicity *in vitro* and *in vivo*.

References

- Abreu, A. C., Borges, A., Simões, L. C., Saavedra, M. J. and Simões, M., 2013a, Antibacterial activity of phenyl isothiocyanate on *Escherichia coli* and *Staphylococcus aureus*, *Medicinal Chemistry*, Vol. 9, no. 5, p: 756-761.
- Abreu, A. C., McBain, A. J. and Simões, M., 2012, Plants as sources of new antimicrobials and resistance-modifying, *Natural Product Reports*, Vol. 29, p: 1007-1021.
- Abreu, A. C., Serra, S., Borges, A., Saavedra, M. J., Salgado, A. and Simões, M., 2013b, Synergism between selected alkaloids and antibiotics against *Staphylococcus aureus*, *ChemMedCom* (Submitted).
- Adwan, G. and Mhanna, M., 2008, Synergistic effects of plant extracts and antibiotics on *Staphylococcus aureus* strains isolated from clinical specimens, *Middle-East Journal of Scientific Research*, Vol. 3, no. 3, p: 134-139.
- Adonizio, A. L., Downum, K., Bennett, B. C. and Mathee, K., 2006, Anti-quorum sensing activity of medicinal plants in southern Florida, *Journal of Ethnopharmacology*, Vol. 105, no. 3, p: 427-435.
- Aeschlimann, J. R., Dresser, L. D., Kaatz G. W. and Rybak, M. J., 1999, Effects of NorA inhibitors on in vitro antibacterial activities and postantibiotic effects of levofloxacin, ciprofloxacin, and norfloxacin in genetically related strains of *Staphylococcus aureus*, *Antimicrobial Agents and Chemotherapy*, Vol. 52, p: 198-203.
- Aires, A., Mota, V. R., Saavedra, M. J., Rosa, E. A. S. and Bennett, R. N., 2009, The antimicrobial effects of glucosinolates and their respective enzymatic hydrolysis products on bacteria isolated from the human intestinal tract, *Journal of Applied Microbiology*, Vol. 106, no. 6, pp 2086-2095.
- Albano, S. M. and Miguel, M. G., 2011, Biological activities of extracts of plants grown in Portugal, *Industrial Crops and Products*, Vol. 33, p: 338-343.
- Al-Hussaini, R. and Mahasneh, A. M., 2009, Microbial growth and quorum sensing activities of herbal plants extracts, *Molecules*, Vol. 14, p: 3425-3435.
- Amaral, C. L. F. and Silva, A. B., 2003, Melhoramento Biotecnológico de Plantas Mediciniais, *Revista Biotecnologia Ciência e Desenvolvimento*, Vol. 30, p: 55-59.
- Andrews, J. M., 2001, Determination of minimum inhibitory concentrations, *Journal of Antimicrobial Chemotherapy*, Vol. 48, no. 1, p: 5-16.
- Anvisa, consulted on website: http://www.anvisa.gov.br/servicos/controle/rede_rm/cursos/boas_praticas/modulo5/objetivos.htm accessed on 22nd December 2012.
- Araújo, E. A., Andrade, N. J., Carvalho, A. F., Ramos, A. M. and Silva, A. S., 2010, Aspectos coloidais da adesão de micro-organismos, *Química Nova*, Vol. 33, no. 9, p: 1940-1948.

- Atawodi, S.E. and Alafiatayo, A.A., 2007, Assessment of the phytochemical and antitrypanosomal properties of some extracts of leaves, stem and root bark of *Landolphia* sp., P . Beauv., *Journal of Ethnopharmacology*, Vol. 114, p: 207-211.
- Beenken, K. E., Dunman, P. M., McAleese, F., Macapagal, D., Murphy, E., Projan S. J., Blevins, J. S. and Smeltzer, M. S., 2004, Global gene expression in *Staphylococcus aureus* biofilms, *Journal of Bacteriology*, Vol. 186, no. 14, p: 4665-4684.
- Bennett, R. N. and Wallsgrave, R. M., 1994, Secondary metabolites in plant defense mechanisms, *Phytotherapy Research*, Vol. 127, p: 617-633.
- Bessemis, E., 1998, The effect of practical conditions on the efficacy of disinfectants, *International Biodeterioration & Biodegradation*, Vol. 41, p: 177-183.
- Beumer, R., Bloomfield, S. F., Exner, M., Fara, G. M., Nath, K. J. and Scott E., 2000, Microbial resistance and biocides, *International Scientific Forum on Home Hygiene*.
- Bighetti, A. E., Antônio, M. A., Kohn, L. K., Rehder, V. L. G., Foglio, M. A., Possenti, A., Vilela, L. and Carvalho, J. E., 2005, Antiulcerogenic activity of a crude hydroalcoholic extract and coumarin isolated from *Mikania laevigata* Schultz Bip, *Phytomedicine*, Vol. 12, p: 72-77.
- Bloomfield, S. F., 2002, Significance of biocide usage and antimicrobial resistance in domiciliary environments, *Symposium series (Society for Applied Microbiology)*, Vol. 31, p: 144S-57S.
- Borges, A., Saavedra, M. J. and Simões M., 2012, The activity of ferulic and gallic acids in biofilm prevention and control of pathogenic bacteria, *Biofouling*, Vol. 28, no. 7, p: 755-767.
- Brooun, A., Liu, S. and Lewis, K., 2000, A dose-response study of antibiotic resistance in *Pseudomonas aeruginosa* biofilms, *Antimicrobial Agents and Chemotherapy*, Vol. 44, p: 640- 646.
- Brown, M. R. W. and Gilbert, P., 1993, Sensitivity of biofilms to antimicrobial agents, *Journal of Applied Bacteriology*, Vol. 74, p: 87S-97S.
- Bryers, J. D., 2000, Biofilm formation and persistence. In: Biofilms II: Process analysis and applications, *Bryers J D. (ed)*, p: 45-88.
- Butler, M. T., Wang, Q. and Harshey, R.M., 2010, Cell density and mobility protect swarming bacteria against antibiotics, *Proceedings of the National Academy of Sciences*, Vol. 107, p: 3776-3781.
- Byarugaba, D. K., 2004, A view on antimicrobial resistance in developing countries and responsible risk factors, *International Journal of Antimicrobial Agents*, Vol. 24, no. 2, p: 105-110.
- Cerca, N. M., 2006, Virulence aspects of *Staphylococcus epidermis*: biofilm formation and Poly-N-Acetyl-Glucosamine production, PhD thesis, University of Minho, Braga.
- Ceri, H., Olson, H. E., Stremick, C., Read, R. R., Morck, D. and Buret, A., 1999, The calgary biofilm device: new technology for rapid determination of antibiotic

- susceptibilities of bacterial biofilms, *Journal of Clinical Microbiology*, Vol. 37, no. 6, p: 1771-1776.
- Chan-Bacab, M. J. and Peña-Rodríguez, L. M., 2001, Plant natural products with leishmanicidal activity, *Natural Product Reports*, Vol., 18, no. 6, p: 674-688.
- Chatterjee, A., Cui, Y., Lui, Y., Dumenyo, C. K. and Chaterjee, A. K., 1995, Inactivation of rsmA leads to overproduction of extracellular pectinases, cellulases, and proteases in *Erwinia carotovora* subsp. *carotovora* in the absence of the starvation/cell density- sensing signal, N-(3-oxohexanoyl)-L-homoserine lactone, *Applied and Environmental Microbiology*, Vol. 61, p: 1959-1967.
- Chaves, L. C. D., 2004, Estudo da cinética de formação de biofilmes em superfícies em contacto com água potável, MSc Thesis, University of Minho.
- Christaki, E., Bonos, E., Giannenas, I. and Florou-Paneri, P., 2012, Aromatic plants as a source of bioactive compounds, *Agriculture*, Vol. 2, p: 228-243.
- Ciocan, I. D. and Bara, I. I., 2007, Plant products as antimicrobial agents, *Molecular Biology*, TOM VIII, p: 151-156.
- Clardy, J., Fischbach, M. A. And Walsh, C. T., 2006, New antibiotics from bacterial natural products, *Nature Biotechnology*, Vol. 24, no. 12, p: 1541-1550.
- Clatworthy, A. E., Pierson, E. and Hung, D. T., 2007, Targeting virulence: a new paradigm for antimicrobial therapy, *Nature Chemical Biology*, Vol. 3, no. 9, p: 541-548.
- Clinical and Laboratory Standards Institute 2005, 'NCCLS M100-S15: Performance standards for antimicrobial susceptibility testing: fifteen informational supplement', vol. 25, no. 1.
- Cloete, T. E., 2003, Resistance mechanisms of bacteria to antimicrobial compounds, *International Biodeterioration & Biodegradation*, Vol. 51, p: 277-282.
- Cosgrove, S. E. and Carmeli, Y., 2003, The impact of antimicrobial resistance on health and economic outcomes, *Antimicrobial Resistance*, Vol. 36, p: 1433-1437.
- Costa, S. S., Viveiros, M., Amaral, L. and Couto, I., 2013, Multidrug efflux pumps in *Staphylococcus aureus*: an update, *The Open Microbiology Journal*, Vol. 7, p: 59-71.
- Cushnie, T. P. T. and Lamb, A. J., 2005, Antimicrobial activity of flavonoids, *International Journal of Antimicrobial Agents*, Vol. 26, p: 343-356.
- Daniels, R., Vanderleyden, J. and Michiels, J., 2004, Quorum sensing and swarming migration in bacteria, *FEMS Microbiology Reviews*, Vol.28, no. 3, p: 261-289.
- Das, K., Tiwari, R. K. S. and Shrivastava, D. K., 2010, Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends, *Journal of Medicinal Plants Research*, Vol. 4, no. 2, p: 104-111.
- Davey, M. E., and O'toole, G. A., 2000, Microbial biofilms: from ecology to molecular genetics, *Microbiology and Molecular Biology Reviews*, Vol. 64, no. 4, p: 847-867.
- Davies, D. G., Parsek, M. R., Pearson, J. P., Iglewski, B. H., Costerton, J. W. and Greenberg, E. P., 1998, The involvement of cell-to-cell signals in the development of a bacterial biofilm, *Science*, Vol. 280, p: 295-298.

- Dickson, J. S. and Koochmaraie, M., 1989, Cell surface charge characteristics and their relationship to bacterial attachment to meat surfaces, *Applied and Environmental Microbiology*, Vol. 55, no. 4, p: 832-836.
- Donlan, R. M., 2002, Biofilms: microbial life on surfaces, *Emerging Infectious Diseases*, Vol. 8, p: 881-890.
- Drumm, B., Neumann, W. N., Policova, Z. and Sherman, P. M., 1989, Bacterial cell surface hydrophobicity properties in the mediation of *In vitro* adhesion by the rabbit enteric pathogen *Escherichia coli* strain RDEC-1, *Journal of Clinical Investigation*, Vol. 84, p: 1588-1594.
- Extremina, C. I., Costa, L., Aguiar, A. I., Peixe, L. and Fonseca, A. P. 2011, Optimization of processing conditions for the quantification of enterococci biofilms using microtitre-plates, *Journal of Microbiological Methods*, Vol. 84, no. 2, p: 167-173.
- Fajardo, A., Martínez-Martín, N., Mercadillo, M., Galán, J. C., Ghysels, B., Mattijs, S., Cornelis, P., Wiehlmann, L., Tümmler, B., Baquero, F. and Martínez, J. L., 2008, The Neglected Intrinsic Resistome of Bacterial Pathogens, *PLoS ONE*, Vol. 3, no. 2, p: 1-6.
- Ferreira, C., Rosmaninho, R., Simoes, M., Pereira, M. C., Bastos, M. M., Nunes, O. C., Coelho, M. and Melo, L. F. 2010, Biofouling control using microparticles carrying a biocide, *Biofouling*, Vol. 26, no. 2, pp 205-212.
- Furukawa, S., Akiyoshi, Y., Komoriya, M., Ogihara, H. and Morinaga, Y., 2010, Removing *Staphylococcus aureus* and *Escherichia coli* biofilms on stainless steel by cleaning-in-place (CIP) cleaning agents, *Food Control*, Vol. 21, no. 5, p: 669-672.
- Fux, C. A., Costerton, J W., Stewart, P. S. and Stoodley, P., 2005, Survival strategies of infectious biofilms, *Trends in Microbiology*, Vol. 13, p: 34-40.
- Ghannoum, M. A., Filler, S. C., Ibrahim, A. S., Fu, Y. and Edwards, J. J. E., 1992, Modulation of Interactions of *Candida albicans* and Endothelial Cells by Fluconazole and Amphotericin B, *Antimicrobial Agents and Chemotherapy*, Vol. 36, p: 2239-2244.
- Gibbons, S., 2004, Anti-staphylococcal plant natural products, *Natural Product Reports*, Vol. 21, p: 263-277.
- Gibbons, S., Oluwatuyi, M. and Kaatz, G.W., 2003, A novel inhibitor of multidrug efflux pumps in *Staphylococcus aureus*, *Journal of Antimicrobial Chemotherapy*, Vol. 51, p: 13-17.
- Gibbons, S., Udo, E. E., 2000, The effect of reserpine, a modulator of multidrug efflux pumps, on the *in vitro* activity of tetracycline against clinical isolates of methicillin-resistant *Staphylococcus aureus* (MRSA) possessing the Tet(K) determinant, *Phytotherapy Research*, Vol. 14, p: 139-40.
- Gilbert, P., Allison, D. G. and McBain, A. J., 2002, Biofilms in vitro and in vivo: do singular mechanisms influx cross-resistance?, *Journal of Applied Microbiology Symposium Supplement*, Vol. 92, p: 98S-110S.
- Girenavar, B., Cepeda, M. L., Soni, K. A., Vikram, A., Jesudhasan, P., Jayaprakasha, G. K., Pillai, S. D. and Patil, B., 2008, Grapefruit juice and its

- furanocoumarins inhibits autoinducer signaling and biofilm formation in bacteria, *International Journal of Food Microbiology*, Vol. 125, p: 204-208.
- Griffin, S.G., Wyllie, S.G., Markham, J.L., Leach, D.N., 1999, The role of structure and molecular properties of terpenoids in determining their antimicrobial activity. *Flavour and Fragrance Journal*, Vol. 14, p: 322–332.
- Guiamet, P. S. and Saravia, S. G., 2005, Laboratory studies of biocorrosion control using traditional and environmentally friendly biocides: An overview, *Latin American Applied Research*, Vol. 35, p: 295-300.
- Guiamet, P. S., Saravia, S. G., Arenas, P., Pérez, M. L., Paz, J. and Borrego, S. F., 2006, Natural products isolated from plants used in biodeterioration control, *Pharmacologyonline*, Vol. 3, p: 537-544.
- Hammerschmidt, R., 1999, Phytoalexins: what have we learned after 60 years?, *Annual Review of Phytopathology*, Vol. 37, p: 285-306.
- Harshey, R. M., 2003, Bacterial motility on a surface: many ways to a common goal, *Annual Review of Microbiology*, Vol. 57, p: 249-273.
- Heinzel, M., 1998, Phenomena of biocide resistance in microorganisms, *International Biodeterioration and Biodegradation*, Vol. 41, p: 225-234
- Hellman, J., Loiselle, P. M., Tehan, M. M., Allaire, J. E., Boyle, L. A., Kurnick, J. T., Andrews, D. M., Kim, K. S. and Warren, H. S., 2000, Outer Membrane protein A., Peptidoglycan-associated lipoprotein, and murein lipoprotein are released by *Escherichia coli* bacteria into serum, *Infection and Immunity*, Vol. 68, no. 5, p: 2566-2572.
- Hemaiswarya, S. and Doble, M., 2009, Synergistic interaction of eugenol with antibiotics against Gram-negative bacteria, *Phytomedicine*, Vol. 16, p: 997-1005.
- Hentzer, M. and Givskov, M., 2003, *Pharmacological inhibition of quorum sensing for the treatment of chronic bacterial infections*, *The Journal of Clinical Investigation*, Vol. 112, p: 1300-1307.
- Hill, A. F., 1952, *Economic Botany: A Textbook of Useful Plant and Plant Products*, 2nd edition, McGraw-Hill Book Company Inc, New York.
- Hoult, J. R. S. and Payát, M., 1996, Pharmacological and Biochemical actions of simple coumarins: natural products with therapeutic potential, *General Pharmacology*, Vol. 27, p: 713-722.
- Howard, D., Cordell, R., McGowan, J. E., Packard, R. M., Scott, R. D. and Solomon, S. L., 2001, Measuring the economic costs of antimicrobial resistance in hospital settings: summary of the Centers for Disease Control and Prevention-Emory Workshop, *Clinical Infectious Diseases*, Vol. 33, p:1573–1578.
- Huet, A. A., Raygada, J. L., Mendiratta, K., Seo, S. M. and Kaatz, G. W., 2008, Multidrug efflux pump overexpression in *Staphylococcus aureus* after single and multiple *in vitro* exposures to biocides and dyes, *Microbiology*, Vol. 154, p: 3144-3153.
- Ibrahim, E. H., Sherman, G., Ward, S., Fraser, V.J. and Kollef, M. H., 2000, The influence of inadequate antimicrobial treatment of bloodstream infections on patient outcomes in the ICU setting, *Chest Journal*, Vol. 118, p: 146–115.

- Janczuk, B., Chibowski, E., Bruque, J. M., Kerkeb, M. L. and Caballero, F. G., 1993, On the consistency of surface free energy components as calculated from contact angles of different liquids: an application to the cholesterol surface, *Journal of Colloid and Interface Science*, Vol. 159, no. 2, p: 421-428.
- Jarrell, K. F. and McBride, M.J., 2008, The surprisingly diverse ways that prokaryotes move, *Nature Reviews Microbiology*, Vol. 6, p: 466–476.
- Jassim, S. A. and Naji, M. A., 2003, Novel antiviral agents: a medicinal plant prespective, *Journal of Applied Microbiology*, Vol. 95, no. 3, p: 412-427.
- Judd, W., Campbell, C., Kellogg, E., Stevens, P. and Donoghue, M., 2002, Plant systematics: a phylogenetic approach, 2nd edition, *Sinauer Associates, Inc. Sunderland*, p: 466-468, 470-473.
- Julkowska, D., Obuchowski, M., Holland, I. B. and Seror, S. J., 2004, Branched swarming patterns on synthetic medium formed by wild-type *Bacillus subtilis* strain 3610: detection of different cellular morphologies and constellations of cells as the complex architecture develops, *Microbiology*, Vol. 150, p: 1839–1849.
- Kalan, L. and Wright, G. D., 2011, Antibiotic adjuvants: multicomponent anti-infective strategies, *Expert Reviews in Molecular Medicine*, Vol. 13, p: null-null.
- Kawahara, M., Human, L. G., Kawahara, K. and Domingue, G. J., 1994a, Electroeluted outer membrane proteins as immunogens, *Immunological Investigation*, Vol. 23, p: 223-230.
- Kawahara, M., Human, L.G., Winningham, J.S. and Domingue, G.J., 1994b, Antibodies to Escherichia coli O6 porins cross-react with urinary pathogens, *Immunobiology*, Vol. 192, p: 65-76.
- Khan, I. A., Mirza, Z. M., Kumar, A., Verma, V. and Qazi, G. N., 2006, Piperine, a phytochemical potentiator of ciprofloxacin against *Staphylococcus aureus*, *Antimicrobial Agents and Chemotherapy*, Vol. 50, no. 2, p:810-812.
- Khan, M. S. A., Zahin, M., Hasan, S., Husain, F.M. and Ahmad, I., 2009, Inhibition of quorum sensing regulated bacterial functions by plant essential oils with special reference to clove oil, *Letters in Applied Microbiology*, Vol. 49, no. 3, p: 354-360.
- Kokare, C. R., Chakraborty, S., Khopade, A. N. and Mahadik, K. R., 2009, Biofilm: Importance and applications, *Indian Journal of Biotechnology*, Vol. 8, p: 159-168.
- Kollef, M. H., Sherman, G., Ward, S. and Fraser, V.J., 1999, Inadequate antimicrobial treatment of infections: a risk factor for hospital mortality among critically ill patients, *Chest Journal*, Vol. 115, p: 462–474.
- Kumar, A. and Schweizer, H.P., 2005, Bacterial resistance to antibiotics: active efflux and reduced uptake, *Advanced Drug Delivery Reviews*, Vol. 57, p: 1486–1513.
- Laemmli, U. K., 1970, Cleavage of structural proteins during the assembly of the head of bacteriophage T4, *Nature*, Vol. 227, p: 680-685.
- Lalitha, M. K., 2004, Manual on Antimicrobial Susceptibility Testing, Christian Medical College, Vellore, Tamil Nadu.

- Lautenbach, E., Patel, J.B., Bilker, W. B., Edelstein, P. H. and Fishman, N. O., 2001, Ex-tended-spectrum b-lactamase–producing *Escherichia coli* and *Klebsiella pneumoniae*: risk factors for infection and impact of resistance on outcomes, *Clinical Infectious Diseases*, Vol. 32, p: 1162–1171.
- Macedo, A.J. and Abraham, W.R., 2009, Can infectious biofilm be controlled by blocking bacterial communication?, *Medicinal Chemistry*, Vol. 5, p: 517–528.
- Maciel, M. A. M., Pinto. A. C. and Veiga, V. F., 2002, Medicinal plants: the need for multidisciplinary scientific studies, *Química Nova*, Vol. 25, p: 429-438.
- Mandal, P., Sinha Babu S. P. and Mandal, N. C., 2005, Antimicrobial activity of saponins from *Acacia auriculiformis*, *Fitoterapia*, Vol. 76, p: 462–465.
- Mah, T.F., and O’Toole, G.A., 2001, Mechanisms of biofilm resistance to antimicrobial agents, *Trends in Microbiology*, Vol. 9, p: 34–39.
- Mahesh, B. and Satish, S., 2008, Antimicrobial activity of some important medicinal plant against plant and human pathogens, *World Journal of Agricultural Sciences*, Vol. 4, p: 839-843.
- Martin, C. A., Hoven, A. D. and Cook, A. M., 2008, Therapeutic frontiers: preventing and treating infectious diseases by inhibiting bacterial quorum sensing, *European Journal of Clinical Microbiology & Infectious Diseases*, Vol. 27, p: 635–642.
- Marquis, A., Genovese, S., Epifano, F. and Grenier, D., 2012, The plant coumarins auraptene and lacinartin as potential multifunctional therapeutic agents for treating periodontal disease, *BioMed Central*, Vol. 12, p: 1-9.
- Mason, T. L., and Wasserman, B. P., 1987, Inactivation of Red Beet Betaglucan Synthase by Native and Oxidized Phenolic Compounds, *Phytochemistry*, Vol. 26, no. 8, p: 2197-2202.
- Masuoka, J. and Hazen, K. C., 1997, Cell Wall Protein Mannosylation Determines *Candida albicans* Cell surface Hydrophobicity, *Microbiology*, Vol. 143, p: 3015-3021.
- McBain, A. J., Ledder, R. G., Sreenivasan, P. and Gilbert, P., 2004, Selection for high-level resistance by chronic triclosan exposure is not universal, *Journal of Antimicrobial Chemotherapy*, Vol. 53, no. 5, p: 772-777.
- McDonnell, G. and Russell, A. D., 1999, Antiseptics and disinfectants: activity, action and resistance, *Clinical Microbiology Reviews*, Vol. 12, p: 147-179.
- McEwen, S. A. and Fedorka-Cray, P. J., 2002, Antimicrobial use and resistance in animals, *Clinical Infectious Diseases*, Vol. 34, no. 3, p: S93–S106.
- McGowan, S., Sebahia, M., Jones, S., Yu, B. and Bainton, N., 1995, Carbapenem antibiotic production in *Erwinia carotovora* is regulated by CarR, a homologue of the LuxR transcriptional activator, *Microbiology*, Vol. 141, p: 541-550.
- McLean, R. J. C., Pierson, L. S. and Fuqua, C., 2004, A simples screening protocol for the identification of quorum signal antagonists, *Journal of Microbiological Methods*, Vol. 58, p: 351-360.
- McManus, M. C., 1997, Mechanisms of bacterial resistance to antimicrobial agents, *American Journal of Health-System Pharmacy*, Vol. 54, p: 1420–33.

- Mello, M. M., 2009, Desenvolvimento de uma metodologia por espectroscopia de fluorescência para quantificação de cumarina e 7-hidroxycumarina em drágeas e soro sintético, Universidade Vale do Paraíba, Brazil.
- Melzig, M. F., Bader, G. and Loose, R., 2001, Investigations of the mechanism of membrane activity of selected triterpenoid saponins, *Planta Medica*, Vol. 67, p: 43–48.
- Mendes, M. D. S., 2007, Caracterização química e molecular de espécies das famílias Lamiaceae e Apiaceae da flora aromática de Portugal, Universidade de Lisboa, Portugal.
- Mendoza, L., Wilkens, M. and Urzua, A., 1997, Antimicrobial study of the resinous exudates and of diterpenoids and flavonoids isolated from some Chilean *Pseudog naphalium* (Asteraceae), *Journal of Ethnopharmacology*, Vol. 58, p: 85-88.
- Molloy, M. P., Herbet, B. R., Slade, M. B., Rabilloud, T., Nouwens, A. S., Williams, K. L. and Gooley, A. A., 2000, Proteomic analysis of the *Escherichia coli* outer membrane, *European Journal of Biochemistry*, Vol. 267, p: 2871-2881.
- Morel, C., Séraphin, D., Teyrouz, A., Larcher, G., Bouchara, J. P., Litaudon, M., Richomme, P. and Bruneton, J., New and antifungal xanthenes from *Clophyllum caledonicum*, *Planta Medica*, Vol. 68, no. 1, p: 41-44.
- Moreno-Dorado, F. J., Guerra, F. M., Aladro F. J., Bustamante, J. M., Jorge, Z. D. and Massanet, G. M., 2000, Synthesis of (±)-11 α -Hydroxy-3-oxo-6 α H,7 α H,10 β Me-endesman-1,2-4,5-dien-6,12-olide, *Journal of Natural Products*, Vol. 63, p: 934-938.
- Mori, A., Nishino, C., Enokib, N. and Tawataa, S., 1987, Antibacterial activity and mode of action of plant flavonoids against *Proteus vulgaris* and *Staphylococcus*, *Phytochemistry*, Vol. 26, p: 2231–2234.
- Muhtar, M., Arshad, M., Ahmad, M., Pomerantz, R. J., Wigdahl, B. and Parveen, Z., 2008, Antiviral potentials of medicinal plants, *Virus Research*, Vol. 131, no. 2, p: 111-120.
- Nascimento, G. G., Locatelli, J., Freitas, P. C. and Silva, G. L. 2000, Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria, *Brazilian Journal of Microbiology*, Vol. 31, no. 4, p: 247-256.
- Naithani, R., Huma, L., Holland, L. E., Shukla, D., McCormick, D. L., Mehta, R. G. and Moriarty, R. M., 2008, Antiviral activity of phytochemicals: A comprehensive review, *Mini-Reviews in Medicinal Chemistry*, Vol. 8, p: 1106-1133.
- Neelima, N., Devidas, N. G., Sudhakar, M. and Kiran, V. J., 2011, A preliminary phytochemical investigation on the leaves of *Solanum xanthocarpum*, *International Journal of Research in Ayurveda and Pharmacy*, Vol. 2, p: 845-850.
- Negm, R.S. and Pistole, T.G., 1998, Macrophages recognize and adhere to an OmpD-like protein of *Salmonella typhimurium*, *FEMS Immunology and Medical Microbiology*, Vol. 20, p: 191-199.

- Nikaido, H., 1994, Prevention of drug access to bacterial targets: Permeability barriers and active efflux, *Science*, Vol. 264, p: 382-388.
- Nikaido, H., 1996, Multidrug efflux pumps of Gram-negative bacteria, *Journal of Bacteriology*, Vol. 178, no. 20, p: 5853-5859.
- Nikaido, H., and Vaara, M., 1985, Molecular basis of bacterial outer membrane permeability, *Microbiology Reviews*, Vol. 49, p: 1-32.
- Okeke, I. N., Laxminarayan, R., Bhutta, Z. A., Duse, A. G. D., Jenkins, P., O'Brien, T. F., Pablos-Mendez, A. and Klugman, K. P., 2005, Antimicrobial resistance in developing countries. Part I: recent trends and current status, *The Lancet Infectious Diseases*, Vol. 5, p: 481-493.
- Oluwatuyi, M., Kaatz, G. W. and Gibbons, S., 2004, Antibacterial and resistance modifying activity of *Rosmarinus officinalis*, *Phytochemistry*, Vol.65, p: 3249-54.
- Othman, H., Loh, H. S., Wiart, C., Khoo, T. J., Lim, K. H. and Ting, K. N., 2011, Optimal methods for evaluating antimicrobial activities from plant extracts, *Journal of Microbiological Methods*, Vol. 83, p: 161-166.
- Otto, M., 2009, Staphylococcus epidermis-the 'accidental' pathogen, *Nature Reviews Microbiology*, Vol. 7, p: 555-567.
- Padmini, E., Valarmathi, A. and Rani, M. U., 2010, Comparative analysis of chemical composition and antibacterial activities of *Mentha spicata* and *Camellia sinensis*, *Asian Journal of Experimental Biological Sciences*, Vol. 1, p: 772-781.
- Pages, C., Prince, P. and Pages, J. M., 1987, Immunological comparison of major outer membrane proteins from different strains of *Escherichia coli*, *Annales de l'Institut Pasteur Microbiology*, Vol. 138, p: 393-406.
- Paramjeet, M., Dipak, S., and Arti, D., 2012, Overview of synthesis and activity of coumarins, *E-International Scientific Research Journal*, Vol. IV, p: 16-35.
- Passador, L., Cook, J. M., Gambello, M. J., Rust, L. and Iglewski, B. H., 1993, Expression of *Pseudomonas aeruginosa* virulence genes requires cell-to-cell communication, *Science*, Vol. 260, p: 1127-1130.
- Pearson, J. P., Passador, L., Iglewski, B. H. and Greenberg, E. P., 1995, A second N-acylhomoserine lactone signal produced by *Pseudomonas aeruginosa*, *Proceedings of the National Academy of Science*, Vol. 92, p: 1490-1494.
- Peeters, E., Nelis, H. J. and Coenye, T., 2008, Comparison of multiple methods for quantification of microbial biofilms grown in microtiter plates, *Journal of Microbiological Methods*, Vol. 72, no. 2, p: 157-165.
- Pereira, M. O., 2001, Comparação da eficiência de dois biocidas (carbamato e glutaraldeído) em sistemas de biofilme, PhD Thesis, University of Minho.
- Plésiat, P. and Nikaido, H., 1992, Outer membranes of Gram-negative bacteria are permeable to steroid probes, *Molecular Microbiology*, Vol. 6, p: 1323-1333.
- Pratt, L. A. and Kolter, R., 1998, Genetic analysis of *Escherichia coli* biofilm formation: roles of flagella, motility, chemotaxis and type I pili, *Molecular Microbiology*, Vol. 30, no. 2, p: 285-293.

- Rahman, M. A. A. and Moon, S. S., 2007, Antimicrobial phenolic derivatives from *Dendranthema zawadskii* var. *Latilobum* Kitamura (Asteraceae), *Archives of Pharmacal Research*, Vol. 30, p: 1374-1379.
- Rao, K. V. K., Schwartz, S. A., Nair, H. K., Aalinkeel, R., Mahajan, S., Chawda, R. and Nair, M. P. N., 2004, Plant derived products as a source of cellular growth inhibitory phytochemicals on PC-3M, DU-145 and LNCaP prostate cancer cell lines, *Current Science*, Vol. 87, p: 1585- 1588.
- Rasooli, I., Shayeh, S., Taghizadeh, M. and Astaneh, S. D. A., 2008, Phytotherapeutic prevention of dental biofilm formation, *Phytotherapy Research*, Vol. 22, p: 1162-1167.
- Ren, D., Sims, J.J. and Wood, T.K., 2002, Inhibition of biofilm formation and swarming of *Bacillus subtilis* by (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone, *Letters in Applied Microbiology*, Vol. 34, p: 293–299.
- Ren, D., Zuo, R., Barrios, A. F. G., Bedzyk, L. A., Eldrige, G. R., Pasmore, M. E. and Wood, T. K., 2005, Differential gene expression for investigation of *Escherichia coli* biofilm inhibition by plant extract ursolic acid, *Applied and Environmental Microbiology*, Vol. 71, p: 4022–4034.
- Rimando, A.M. and Suh, N., 2008, Biological/Chemopreventive Activity of Stilbenes and Their Effect on Colon Cancer, *Planta Medica*, Vol. 74, p: 1635-1643.
- Russell, A. D., 2003, Biocide use and antibiotic resistance: the relevance of laboratory findings to clinical and environmental situations, *The Lancet Infectious Diseases*, Vol. 3, p: 794- 803.
- Saavedra, M. J., Borges, A., Dias, C., Aires, A., Bennett, R. N., Rosa, E. S. and Simões, M., 2010, Antimicrobial activity of phenolics and glucosinate hydrolysis products and their synergy with streptomycin against pathogenic bacteria, *Medicinal Chemistry*, Vol. 6, p: 174-183.
- Saleem, M., Nazir, M., Ali, M. S., Hussain, H., Lee, Y. S., Riaz, N. and Jabbar, A., 2010, Antimicrobial natural products: an update on future antibiotic drug candidates, *Natural Product Reports*, Vol. 27, p: 238-254.
- Savage, P. B., 2001, Multidrug resistant bacteria: overcoming antibiotic permeability barriers of gram-negative bacteria, *Annals of Internal Medicine*, Vol. 33, p: 167-171.
- Saúde-info, consulted on website: <http://saude-info.info/vantagens-e-desvantagens-da-fitoterapia.html> accessed on 22nd December 2012.
- Scalbert, A, 1991, Antimicrobial properties of tannins, *Phytochemistry*, Vol. 30, p: 3875-3883.
- Schmitz, F. J., Fluit, A. C., Lückefahr, M., Engler, B., Hofmann, B., Verhoef, J., Heinz, H. P., Hadding, U. and Jones, M. E., 1998, The effect of reserpine, an inhibitor of multidrug efflux pumps, on the in-vitro activities of ciprofloxacin, sparfloxacin and moxifloxacin against clinical isolates of *Staphylococcus aureus*, *Journal of Antimicrobial Chemotherapy*, Vol. 42, no. 6, p: 807-810.
- Sefton, A. M., 2002, Mechanisms of antimicrobial resistance: their clinical relevance in the new Millennium, *Drugs*, Vol. 62, p: 557–566.

- Severiano, M. V. N., Dantas, I. C., Silva, J. C. and Felismino, D. C., 2010, Survey of medicinal plants cultured in the center of study and research Malaquias da Silva Amorim, *Revista de Biologia e Farmácia*, Vol. 4, p: 91-101.
- Shrout, J. D., Chopp, D. L., Just, C. L., Hentzer, M., Givskov, M. and Parsek, M. R., 2006, The impact of quorum sensing and swarming motility on *Pseudomonas aeruginosa* biofilm formation is nutritionally conditional, *Molecular Microbiology*, Vol. 62, p: 1264–1277.
- Sillankorva, S., Oliveira, R., Vieira, M. J., Sutherland, I. and Azeredo, J., 2004, Bacteriophage ϕ S1 infection of *Pseudomonas fluorescens* planktonic cells versus biofilms, *Biofouling*, Vol. 20, no. 3, p: 133-138.
- Simões, M., 2005, Use of biocides and surfactants to control *Pseudomonas fluorescens* biofilms: role of the hydrodynamic conditions, PhD, University of Minho, Braga.
- Simões, M., Bennett, R. N. and Rosa, E. A. S., 2009, Understanding antimicrobial activities of phytochemicals against multidrug resistant bacteria and biofilms, *Natural Product Reports*, Vol. 26, p: 746-757.
- Simões, M., Rocha, R., Coimbra, M. A. and Vieira, M., 2008, Enhancement of *Escherichia coli* and *Staphylococcus aureus* antibiotic susceptibility using sesquiterpenoids, *Medicinal Chemistry*, Vol. 4, no. 6, p: 616-623.
- Simões, L., Simões, M. and Vieira, M., 2010, Adhesion and biofilm formation on polystyrene by drinking water-isolated bacteria', *Antonie van Leeuwenhoek*, Vol. 98, no. 3, p: 317-329.
- Sinde, E. and Carballo, J., 2000, Attachment of *Salmonella* spp. and *Listeria monocytogenes* to Stainless Steel, Rubber and Polytetrafluorethylene: the Influence of Free Energy and the Effect of Commercial Sanitizers, *Food Microbiology*, Vol. 17, p: 439-447.
- Sinha Babu, S. P., Sarkar, D., Ghosh, N. K., Saha, A., Sukul, N. C. and Bhattacharya, S., 1997, Enhancement of membrane damage by saponins isolated from *Acacia auriculiformis*, *Japanese Journal of Pharmacology*, Vol. 75, p: 451–454.
- Smith, J. T. and Lewin, C. S., 1993, Mechanisms of antimicrobial resistance and implications for epidemiology, *Veterinary Microbiology*, Vol. 35, p: 233–242.
- Smith, E., Williamson, M., Wareham, N., Kaatz, G. W., and Gibbons, S., 2007, Antibacterials and modulators of bacterial resistance from the immature cones of *Chamaecyparis lawsoniana*, *Phytochemistry*, Vol. 68, p: 210-217.
- Sopirala, M. M., Mangino, J. E., Gebreyes, W. A., Biller, B., Bannerman, T., Balada-Llassat, J. M. and Pancholi, P., 2010, Synergy testing by Etest, microdilution checkerboard, and time-kill methods for pan-drug-resistant *Acinetobacter baumannii*, *Antimicrobial Agents and Chemotherapy*, Vol. 54, p: 4678-83.
- Souza, G. S., Castro, E. M., Soares, A. M., Pinto, J. E. B. P., Resende, M. G. and Bertolucci, S. K. V., 2011, Crescimento, teor de óleo essencial e conteúdo de cumarina de plantas jovens de guaco (*Mikania glomerata* Sprengel) cultivadas sob malhas coloridas, *Revista Biotemas*, Vol. 24, p: 1-11.
- Spratt, B. G., 1994, Resistance to antibiotics mediated by target alterations, *Science*, Vol. 264, no. 5157, p: 388-393.

- Sriram, D., Rao, V. S., Chandrasekhara, K. V. and Yogeewari, P., 2004, Progress in the research of Artemisinin and its analogues as antimalarials, an update, *Nature Products Research*, Vol.18, no. 6, p: 503-527.
- Stary, F., 1996, The natural guide to medicinal herbs and plants, *Barnes&Noble Inc.*
- Stewart, P.S. and Costerton, J.W., 2001, Antibiotic resistance of bacteria in biofilms, *Lancet*, Vol. 358, p: 135–138.
- Stepanović, S., Vukovic, D., Dakic, I., Savic, B. and Svabic-Vlahovic, M., 2000, A modified microtiter-plate test for quantification of staphylococcal biofilm formation, *Journal of Microbiological Methods*, Vol. 40, no. 2, p: 175-179.
- Stickland, H.G., Davenport, P.W., Lilley, K.S., Griffin, J.L. and Welch, M., 2010, Mutation of nfxB causes global changes in the physiology and metabolism of *Pseudomonas aeruginosa*, *Journal of Proteome Research*, Vol. 9, p: 2957–2967.
- Suffredini, I. B., Varella, A.D. and Younes, R. N., 2006, Cytotoxic molecules from natural sources: tapping the Brazilian biodiversity, *Anti-cancer Agents in Medicinal Chemistry*, Vol. 6, p: 367-375.
- Packiavathy, I. A. S. V., Agilandeswari, P., Musthafa, K. S., Pandian S. K. and Ravi A. V., 2012. Antibiofilm and quorum sensing inhibitory potential of *Cuminum cyminum* and its secondary metabolite methyl eugenol against Gram negative bacterial pathogens, *Food Research International*, Vol. 45, no. 1, p: 85-92.
- Tegos, G., Stermitz, F. R., Lomovskaya, O. and Lewis, K., 2002, Multidrug pump inhibitors uncover remarkable activity of plant antimicrobials, *Antimicrobial Agents and Chemotherapy*, Vol. 46, no. 10, p: 3133-3141.
- Thoung, P. T., Hung, T. M., Ngoc, T. M., Ha, D. T., Min, B. S., Kwack, S. J., Kang, T. S., Choi, J. S. and Bae, K., 2009, Antioxidant activities of coumarins from Korean medicinal plants and their structure-activity relationship, *Phytotherapy Research*, Vol. 24, p: 101-106.
- Trentin, D. N., Giordani, R. B., Zimmer, K. R., Silva, A. G., Silva, M. V., Correia, M. T. S., Baumvol, I. J. and Macedo, A. J., 2011, Potencial of medicinal plants from the Brazilian semi-arid region (Caatinga) against *Staphylococcus epidermis* planktonic and biofilm lifestyles, *Journal of Ethnopharmacology*, Vol. 137, p: 327-335.
- Treyvaud Amiguet, V., Petit, P., Ta, C.A., Nuñez, R., Sánchez-Vindas, P., Alvarez, L. P., Smith, M. L., Arnason, J. T. and Durst, T., 2006, Phytochemistry and antifungal properties of the newly discovered tree *Pleodendron costaricense*, *Journal of Natural Products*, Vol. 69, no. 7, p: 1005-1009.
- Udenigwe, C. C., Ramprasath, V. R., Aluko, R. E. and Jones, P. J., 2008, Potential of resveratrol in anticancer and anti-inflammatory therapy, *Nutrition Reviews*, Vol. 66, p: 445–454.
- Urs, N. V. R. R. and Dunleavy, J. M., 1975, Enhancement of the bactericidal activity of a peroxidase system by phenolic compounds (*Xanthomonas phaseoli* var. *sojensis*, soybeans), *Phytopathology*, Vol. 65, p: 686-690.
- Vaara, M., 1993, Antibiotic-supersusceptible mutants of *Escherichia coli* and *Salmonella typhimurium*, *Antimicrobial Agents and Chemotherapy*, Vol. 37, p: 2255–2260.

- van Bambeke, F., Glupczynski, Y., Plésiat, P., Pechère, J. C., and Tulkes, P. M., 2003, Antibiotic efflux pumps in prokaryotic cells: occurrence, impact on resistance and strategies for the future of antimicrobial therapy, *Journal of Antimicrobial Chemotherapy*, Vol. 51, p: 1055-1065.
- van Oss, C. J., Chaudhury, M. K. and Good, R. J., 1987, Monopolar surfaces, *Advances in Colloid and Interface Science*, Vol. 28, p: 35-64.
- van Oss, C. J. and Giese, R. F., 1995, The Hydrophilicity and Hydrophobicity of Clay Minerals, *Clay Minerals*, Vol. 43, p: 474-477.
- van Oss, C. J., Good, R. J. and Chaudhury, M. K., 1988, Additive and nonadditive surface tension components and the interpretation of contact angles, *Langmuir*, Vol. 4, no. 4, p: 884-891.
- van Oss, C. J., Ju, L., Chaudhury, M. K. and Good, R. J., 1989, Estimation of the polar parameters of the surface tension of liquids by contact angle measurements on gels, *Journal of Colloid and Interface Science*, Vol. 128, no. 2, p: 313-319.
- Vidaver, A. K., 2002, Uses of antimicrobials in plant agriculture, *Clinical Infectious Diseases*, Vol. 34, no. 3, p: S107–S110.
- Vieira, M. J., 1995, Estudo da Formação de Filmes Biológicos por *Pseudomonas fluorescens* e dos efeitos associados à transferência de massa interna e à incorporação de partículas de caulino, PhD Thesis, University of Minho.
- von Bodman S. B. and Farrand S. K., 1995, Capsular polysaccharide biosynthesis and pathogenicity in *Erwinia stewartii* require induction by an N-acylhomoserine lactone autoinducer, *Journal of Bacteriology*, Vol. 177, p: 5000-5008.
- Wang, Z. C., Feng, D. Q. and Ke, C. H., 2013, Coumarins from the herb *Cnidium monnieri* and chemically modified derivatives as antifoulants against *Balanus albicostatus* and *Bugula neritina* Larvae, *International Journal of Molecular Sciences*, Vol. 14, p: 1197-1206.
- Waters, C. M. and Bassler, B. L., 2005, Quorum-sensing: cell-to-cell communication in Bacteria, *Annual Review of Cell and Developmental Biology*, Vol. 21, p:319-346.
- Watnick, P. and Kolter, R., 2000, Biofilm, city of microbes, *Journal of Bacteriology*, Vol. 182, p: 2675-2679.
- Wise, R. and Soulsby, E. J., 2002, Antibiotic resistance-an evolving problema, *Veterinary Record*, Vol. 151, p: 371–372.
- Webber, M. A. and Piddock, L. J.V., 2003, The importance of efflux pumps in bacterial antibiotic resistance, *Journal of Antimicrobial Chemotherapy*, Vol. 51, p: 9-11.
- Woodford, N. and Ellington, M. J., 2007, The emergence of antibiotic resistance by mutation, *Clinical Microbiology and Infection*, Vol. 13, no. 1, p: 5-18.
- Xu, C., Lin, X., Ren, H., Zhang, Y., Wang, S. and Peng, X., 2006, Analysis of outer membrane proteome of *Escherichia coli* related to resistance to ampicillin and tetracycline, *Proteomics*, Vol.6, p: 462-473.

Master Thesis

Zhang, H. Z., Hackbarth, C. J., Chansky, K. M. and Chambers, H. J., 2001, A proteomic transmembrane signalling pathway and resistance to beta-lactams in staphylococci, *Science*, Vol. 291, p: 1962-1965.

Zahin, M., Hasan, S., Aqil, F., Khan, M. S. A., Husain, F. M. and Ahmad I., 2010, Screening of certain medicinal plants from India for their anti-quorum sensing activity, *Indian Journal of Experimental Biology*, Vol. 48, no. 12, p: 1219-1224.

Appendix

A. Experimental data

Table A.1. describes the physico-chemical characteristics of polystyrene in order to predict cells adhesion to this surface (Simões et al., 2010).

Table A.1. Physico-chemical characterization of polystyrene (PS).

	Contact Angle (°)			Surface tension parameters (mJ/m ²)			Hydrophobicity (mJ/m ²)	Zeta potential (mV)
	θ_w	θ_F	θ_B	γ^{LW}	γ^+	γ^-	ΔG_{sws}^{TOT}	
<i>PS</i>	83±3	71±2	28±1	39	0.0	9.9	-44	-32±2