

CORRECTION

Open Access



Correction to: The relationship between endogenous thymidine concentrations and [^{18}F]FLT uptake in a range of preclinical tumour models

Kathrin Heinzmann^{1,8}, Davina Jean Honess¹, David Yestin Lewis^{1,7}, Donna-Michelle Smith¹, Christopher Cawthorne^{2,9}, Heather Keen³, Sandra Heskamp⁴, Sonja Schelhaas⁵, Timothy Howard Witney^{6,10}, Dmitry Soloviev^{1,7}, Kaye Janine Williams^{2,7}, Andreas Hans Jacobs⁵, Eric Ofori Aboagye⁶, John Richard Griffiths¹ and Kevin Michael Brindle^{1,7,11*}

Correction

Unfortunately, the original version of Figs. 4, 5 and 6b in the article [1] contained errors in the n numbers as indicated on the columns. Please note that column heights and error bars in the original figures and data in the ESM tables are correct and statistical tests are valid. These corrections do not affect any results or conclusions in this article.

The correct n numbers are as follows:

Figure 4 (http://media.springernature.com/full/springer-static/image/art%3A10.1186%2Fs13550-016-0218-3/MediaObjects/13550_2016_218_Fig4_HTML.gif): 'HCT116' (IC) $n = 8$; 'A549' (WWU) $n = 7$; 'HCT116' (WMIC) $n = 3$; 'H1975' (AZ) $n = 6$.

Figure 5 (http://media.springernature.com/full/springer-static/image/art%3A10.1186%2Fs13550-016-0218-3/MediaObjects/13550_2016_218_Fig5_HTML.gif): 'ONU (AZ)' $n = 6$; 'H1975 in ONU (AZ)' $n = 6$; 'A431 in ONU (AZ)' $n = 6$.

Figure 6b (http://media.springernature.com/full/springer-static/image/art%3A10.1186%2Fs13550-016-0218-3/MediaObjects/13550_2016_218_Fig6_HTML.gif): 'PC9' $n = 13$.

Author details

¹Cancer Research UK Cambridge Institute, University of Cambridge, Cambridge, UK. ²Wolfson Molecular Imaging Centre, Manchester Pharmacy School, University of Manchester, Manchester, UK. ³Personalised Healthcare and Biomarkers, AstraZeneca, Alderley Park, Macclesfield, UK. ⁴Department of Radiology and Nuclear Medicine, Radboud University Medical Centre,

Nijmegen, The Netherlands. ⁵European Institute for Molecular Imaging (EIMI), Westfälische Wilhelms-Universität (WWU), University Hospital of Münster, Münster, Germany. ⁶Comprehensive Cancer Imaging Centre, Imperial College London, London, UK. ⁷CRUK-EPSC Cancer Imaging Centre in Cambridge and Manchester, Cambridge, UK. ⁸Comprehensive Cancer Imaging Centre, Imperial College London, London, UK. ⁹Positron Emission Tomography Research Centre, University of Hull, Hull, UK. ¹⁰UCL Centre for Advanced Biomedical Imaging, University College London, London, UK. ¹¹Cancer Research UK Cambridge Institute, Li Ka Shing Centre, Cambridge CB2 0RE, UK.

Received: 30 November 2017 Accepted: 30 November 2017

Published online: 15 December 2017

Reference

1. Heinzmann K, Honess DJ, Lewis DY, Smith D-M, Cawthorne C, Keen H, Heskamp S, Schelhaas S, Witney TH, Soloviev D, Williams KJ, Jacobs AH, Aboagye EO, Griffiths JR, Brindle KM. The relationship between endogenous thymidine concentrations and [^{18}F]FLT uptake in a range of preclinical tumour models. *EJNMMI Res.* 2016;6:63. <https://doi.org/10.1186/s13550-016-0218-3>.

* Correspondence: kmb1001@cam.ac.uk

¹Cancer Research UK Cambridge Institute, University of Cambridge, Cambridge, UK

⁷CRUK-EPSC Cancer Imaging Centre in Cambridge and Manchester, Cambridge, UK